

The Categorical Product of Two 5-Chromatic Digraphs can be 3-Chromatic

Stéphane Bessy, Stéphan Thomassé

► **To cite this version:**

Stéphane Bessy, Stéphan Thomassé. The Categorical Product of Two 5-Chromatic Digraphs can be 3-Chromatic. *Discrete Mathematics*, Elsevier, 2005, 305, pp.344-346. 10.1016/j.disc.2004.01.023 . lirmm-00140331

HAL Id: lirmm-00140331

<https://hal-lirmm.ccsd.cnrs.fr/lirmm-00140331>

Submitted on 5 Apr 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The Categorical Product of two 5-chromatic digraphs can be 3-chromatic.

Stéphane Bessy
and
Stéphan Thomassé

Laboratoire LaPCS, U.F.R. de Mathématiques,
Université Claude Bernard - Lyon 1, 50, avenue Tony Garnier,
Bâtiment RECHERCHE [B], Domaine de Gerland
69367 Lyon Cedex 07, France
email : bessy@univ-lyon1.fr
thomasse@univ-lyon1.fr

Abstract

We provide an example of a 5-chromatic oriented graph D such that the categorical product of D and TT_5 is 3-chromatic, where TT_5 is the transitive tournament on 5 vertices.

The transitive tournament TT_5 is the digraph on the vertex set $\{1, 2, 3, 4, 5\}$ and edge set $\{(i, j) : i < j\}$. Let $D_1 = (V_1, E_1)$ and $D_2 = (V_2, E_2)$ be two digraphs. The *categorical product* of D_1 and D_2 is the digraph $D_1 \times D_2 = (V_1 \times V_2, E)$ where $E = \{((x, y), (z, t)) : (x, z) \in E_1 \text{ and } (y, t) \in E_2\}$. Motivated by Hedetniemi's conjecture (see Sauer [5] for a survey), the following function was defined:

$$\delta(n) := \min\{\chi(D_1 \times D_2) : \chi(D_1) = \chi(D_2) = n \text{ and } D_1, D_2 \in \mathcal{D}\}$$

where \mathcal{D} is the class of finite digraphs and $\chi(D)$ is the chromatic number of D . The first striking result concerning this function was proved by Poljak and Rödl [4]: either δ is bounded by 4 or it tends to infinity. Later on, Poljak [3], and independently Zhu [6] proved that 4 can be replaced by 3. When δ is restricted to undirected graphs, Hedetniemi [2] conjectured that $\delta(n) = n$. El-Zahar and Sauer proved in [1] that $\delta(4) = 4$ for undirected graphs. For directed graphs, it is known that $\delta(4) = \delta(3) = 3$.

Theorem 1 $\delta(5) = 3$

Proof. The oriented graph D depicted in Fig. 1 is 5-chromatic. Observe for this that the twelve vertices with degree 8 induce a graph G which is uniquely 4 colorable up to a rotation. Indeed the stability of G is 3 and the color classes must consist of three consecutive vertices in the cyclic order. Now, one of the three vertices with degree 4 has to have one neighbour in each of the four color classes. Thus D is not 4-colorable. The 3-coloration f of $D \times TT_5$ is given by the label of the vertices of D : when a vertex x of D is labelled by c^3a^2 , for instance, we mean that $f(x, 1) = c, f(x, 2) = c, f(x, 3) = c, f(x, 4) = a, f(x, 5) = a$. To see that f is a good coloration, note that when there is an edge from x to y (both seen as words of length 5 on the alphabet $\{a, b, c\}$), the i^{th} letter of x is not equal to the j^{th} letter of y for every $j > i$. \square




Fig. 1

This example is a subgraph of the graph $K_3^{TT^5}$ which has chromatic number 5 (this is not hard to check). A short analysis also gives that $\chi(K_3^{TT^4}) = 6$ and $\chi(K_3^{TT^k}) = 4$ for all $k > 5$. We were not able to find an infinite family of 5-chromatic digraphs (D_1, D_2) such that $\chi(D_1 \times D_2) = 3$.

References

- [1] E. El-Zahar and N. Sauer, The chromatic number of the product of two 4-chromatic graphs is 4, *Combinatorica*, **5** (1985), 121-126.
- [2] S.T. Hedetniemi, Homomorphisms and graph automata, University of Michigan Technical Report, **03105-44-T** (1966).
- [3] S. Poljak, Coloring digraphs by iterated antichains. *Comment. Math. Univ. Carolin.*, **32** (1991), 209-212.
- [4] S. Poljak and V. Rödl, On the arc-chromatic number of a digraph, *J. Combin. Theory Ser. B*, **31** (1981), 190-198.
- [5] N. Sauer, Hedetniemi's conjecture - a survey, preprint.
- [6] X. Zhu, Multiplicative Structures, Ph. D. thesis (1990), The university of Calgary.