

Evolutionary Dynamics of the Restorer and a non-restorer Allelic Rfo Locus.

José R. Hernández Mora, Eric Rivals, Hakim Mireau, Françoise Budar

► **To cite this version:**

José R. Hernández Mora, Eric Rivals, Hakim Mireau, Françoise Budar. Evolutionary Dynamics of the Restorer and a non-restorer Allelic Rfo Locus.. Société française de biologie végétale. SFBV'07: VIIe Colloque National de la Société Française de Biologie Végétale, Sep 2007, Versailles, France. pp.113, 2007. <lirmm-00203886>

HAL Id: lirmm-00203886

<https://hal-lirmm.ccsd.cnrs.fr/lirmm-00203886>

Submitted on 11 Jan 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Evolutionary dynamics of the restorer and a non-restorer allelic Rfo locus.

J.R. Hernández Moraa, E. Rivalsb, H. Mireaua et F. Budara

aStation de Génétique et d'Amélioration des Plantes - IJPB - INRA, route de Saint Cyr, 78026 Versailles, France

bLaboratoire d'Informatique de Robotique et de Microélectronique de Montpellier (LIRMM), 161 rue Ada, 34392 Montpellier, France

Contact: budar@versailles.inra.fr

The Rfo locus, first described in Asian radish cultivars and later introduced into Brassica, is involved in the restoration of fertility in the Ogura cytoplasmic male sterility system (CMS). CMS systems are constituted of two actors: a mitochondrial protein that causes male sterility and a nuclear gene that encodes a mitochondria-targeted protein able to impair the expression of the sterility gene.

The nuclear restorer gene in the Rfo-Ogura system, as all identified restorer genes (with the only exception of the Rf2 Texas maize restorer), encodes a protein belonging to the pentatricopeptide repeat (PPR) family. The PPR gene family is a very large family in plants, with about 450 members in *A. thaliana*, and their function is largely unknown, although some members have proven to play roles in posttranscriptional organelle gene regulation.

Recent studies analyzing chromosome regions with duplicated PPR genes (allelic variants of a restorer locus in rice and two chromosome regions in *A. thaliana*) suggest revealed high levels of recombination in these regions. In addition, restorer genes (because of their ability to adapt to specificity changing/fast evolving targets) are evolved in a similar way as “resistant genes”. Anyway, evolutionary dynamics of these loci is largely unknown so far.

In the Rfo region, the restorer gene seems to be duplicated too: three genes encoding highly related PPR proteins are present in the Rfo locus (even if just one of the proteins is able to restorer fertility). We recently sequenced a European non-restorer allelic locus. Analysis of this sequence shows the presence of only two PPR genes in an apparently very dynamic region.

In this study, we analyzed the European allelic variant sequence of the Rfo locus and the possible evolutionary events that occurred in both loci, in order to give insight into the dynamics and evolution of the Rfo locus.

Nombre de mots du résumé: 300

Thème: Chloroplastes et Mitochondries