

## An edge variant of the Erdős–Pósa property

Jean-Florent Raymond, Ignasi Sau, Dimitrios M. Thilikos

► **To cite this version:**

Jean-Florent Raymond, Ignasi Sau, Dimitrios M. Thilikos. An edge variant of the Erdős–Pósa property. *Discrete Mathematics*, Elsevier, 2016, 339 (8), pp.2027-2035. <10.1016/j.disc.2016.03.004>. <lirmm-01347430>


**HAL Id: lirmm-01347430**

**<https://hal-lirmm.ccsd.cnrs.fr/lirmm-01347430>**

Submitted on 4 Oct 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# An edge variant of the Erdős–Pósa property<sup>☆</sup>



Jean-Florent Raymond<sup>a,b,c,\*</sup>, Ignasi Sau<sup>b</sup>, Dimitrios M. Thilikos<sup>b,d</sup>

<sup>a</sup> Faculty of Mathematics, Informatics and Mechanics, University of Warsaw, Warsaw, Poland

<sup>b</sup> ALGCo Project Team, CNRS, LIRMM, Montpellier, France

<sup>c</sup> University of Montpellier, Montpellier, France

<sup>d</sup> Department of Mathematics, National and Kapodistrian University of Athens, Athens, Greece

## ARTICLE INFO

### Article history:

Received 2 April 2014

Received in revised form 15 September 2015

Accepted 6 March 2016

Available online 26 April 2016

### Keywords:

Erdős–Pósa property

Packings in graphs

Coverings in graphs

## ABSTRACT

For every  $r \in \mathbb{N}$ , we denote by  $\theta_r$  the multigraph with two vertices and  $r$  parallel edges. Given a graph  $G$ , we say that a subgraph  $H$  of  $G$  is a *model* of  $\theta_r$  in  $G$  if  $H$  contains  $\theta_r$  as a contraction. We prove that the following edge variant of the Erdős–Pósa property holds for every  $r \geq 2$ : if  $G$  is a graph and  $k$  is a positive integer, then either  $G$  contains a packing of  $k$  mutually edge-disjoint models of  $\theta_r$ , or it contains a set  $S$  of  $f_r(k)$  edges such that  $G \setminus S$  has no  $\theta_r$ -model, for both  $f_r(k) = O(k^2 r^3 \text{polylog } kr)$  and  $f_r(k) = O(k^4 r^2 \text{polylog } kr)$ .

© 2016 Elsevier B.V. All rights reserved.

## 1. Introduction

Typically, an Erdős–Pósa property reveals relations between covering and packing invariants in combinatorial structures. The origin of the study of such properties comes from the Erdős–Pósa Theorem [5], stating that there is a function  $f : \mathbb{N} \rightarrow \mathbb{N}$  such that for every  $k \in \mathbb{N}$  and for every graph  $G$ , either  $G$  contains  $k$  vertex-disjoint cycles, or there is a set  $X$  of  $f(k)$  vertices in  $G$  meeting all cycles of  $G$ . In particular, Erdős and Pósa proved this result for  $f(k) = O(k \cdot \log k)$ .

An interesting line of research aims at extending Erdős–Pósa Theorem for packings and coverings of more general graph structures. In this direction, we say that a graph class  $\mathcal{G}$  satisfies the Erdős–Pósa property if there exists a function  $f_{\mathcal{G}} : \mathbb{N} \rightarrow \mathbb{N}$  such that, for every graph  $G$  and every positive integer  $k$ , either  $G$  contains  $k$  mutually vertex-disjoint subgraphs, each isomorphic to a graph in  $\mathcal{G}$ , or it contains a set  $S$  of  $f_{\mathcal{G}}(k)$  vertices meeting every subgraph of  $G$  that is isomorphic to a graph in  $\mathcal{G}$ . When this property holds for a class  $\mathcal{G}$ , we call the function  $f_{\mathcal{G}}$  the *gap of the Erdős–Pósa property for the class  $\mathcal{G}$* . In this sense, the classic Erdős–Pósa Theorem says that the class containing all cycles satisfies the Erdős–Pósa property with gap  $O(k \cdot \log k)$ .

Given a graph  $J$ , we denote by  $\mathcal{M}(J)$  the set of all graphs containing  $J$  as a contraction. Robertson and Seymour proved the following proposition, which in particular can be seen as an extension of the Erdős–Pósa Theorem.

**Proposition 1.** *Let  $J$  be a graph. The class  $\mathcal{M}(J)$  satisfies the Erdős–Pósa property if and only if  $J$  is planar.*

<sup>☆</sup> This work was supported by the ANR project AGAPE (ANR-09-BLAN-0159) and the Languedoc-Roussillon Project “Chercheur d’avenir” KERNEL. The first author was partially supported by the (Polish) National Science Centre grant PRELUDIUM 2013/11/N/ST6/02706. The third author was co-financed by the European Union (European Social Fund ESF) and Greek national funds through the Operational Program “Education and Lifelong Learning” of the National Strategic Reference Framework (NSRF) - Research Funding Program: ARISTEIA II.

\* Corresponding author.

E-mail addresses: [jean-florent.raymond@mimuw.edu.pl](mailto:jean-florent.raymond@mimuw.edu.pl) (J.-F. Raymond), [ignasi.sau@lirmm.fr](mailto:ignasi.sau@lirmm.fr) (I. Sau), [sedthilk@thilikos.info](mailto:sedthilk@thilikos.info) (D.M. Thilikos).




Fig. 1. The graph  $\theta_5$ .

A proof of [Proposition 1](#) appeared for the first time in [18]. Another proof can be found in Diestel's monograph [4, Corollary 12.4.10 and Exercise 40 of Chapter 12]. In view of [Proposition 1](#), it is natural to try to derive good estimations of the gap function  $f_{\mathcal{M}(J)}$  in the case where  $J$  is a planar graph. In this direction, the recent breakthrough results of Chekuri and Chuzhoy imply that  $f_{\mathcal{M}(J)}(k) = k \cdot \text{polylog } k$  [2] when  $J$  is a planar graph and, even more, that  $f_{\mathcal{M}(J)} = (k + |V(J)|)^{O(1)}$  [3]. Before this, the best known estimation of the gap for planar graphs was exponential, namely  $f_{\mathcal{M}(J)}(k) = 2^{O(k \log k)}$ , and could be deduced from [14] using the proof of [18]. Moreover, some improved polynomial gaps have been proven for particular instantiations of the graph  $J$  (see [6–9,15,16]). Another direction is to add restrictions on the graphs  $G$  that we consider, which usually allows to optimize the gap  $f_{\mathcal{M}(J)}$ . In this direction, it is known that  $f_{\mathcal{M}(J)} = O(k)$  in the case where graphs are restricted to some non-trivial minor-closed class [10].

We consider the edge counterpart of the Erdős–Pósa property, where packings are edge-disjoint (instead of vertex-disjoint) and coverings contain edges instead of vertices. We say that a graph class  $\mathcal{G}$  satisfies *the edge variant of the Erdős–Pósa property* if there exists a function  $f_{\mathcal{G}}$  such that, for every graph  $G$  and every positive integer  $k$ , either  $G$  contains  $k$  mutually edge-disjoint subgraphs, each isomorphic to a graph in  $\mathcal{G}$ , or it contains a set  $X$  of  $f_{\mathcal{G}}(k)$  edges meeting every subgraph of  $G$  that is isomorphic to a graph in  $\mathcal{G}$ . Recently, the edge variant of the Erdős–Pósa property was proved in [12] for 4-edge-connected graphs in the case where  $\mathcal{G}$  contains all odd cycles.

In this paper we concentrate on the case where  $\mathcal{G} = \mathcal{M}(J)$  for some graph  $J$ . We find it an interesting question whether an edge-analogue of [Proposition 1](#) exists or not. To our knowledge, the only case for which  $\mathcal{M}(J)$  satisfies the edge variant of the Erdős–Pósa property is when  $J = K_3$ , i.e. when the class of graphs  $\mathcal{G}$  contains all cycles. This result is the edge-counterpart of the Erdős–Pósa Theorem and appears as a (hard) exercise in [4, Exercise 23 of Chapter 7]. For every  $r \geq 2$ , let  $\theta_r$  be the graph containing two vertices and  $r$  multiple edges between them (see [Fig. 1](#)). The results of this paper can be stated as follows:

**Theorem 1.** *The edge variant of the Erdős–Pósa property holds for  $\mathcal{M}(\theta_r)$  with gap  $f_{\mathcal{M}(\theta_r)}$ , with*

$$f_{\mathcal{M}(\theta_r)}(k) = O(k^2 r^3 \text{polylog } kr) \quad \text{and} \quad f_{\mathcal{M}(\theta_r)}(k) = O(k^4 r^2 \text{polylog } kr).$$

[Theorem 1](#) is the edge-counterpart of the main result of [9]. The proof is presented in [Section 3](#) and contains three main ingredients. The first is a reduction of the problem to graphs of bounded degree, presented in [Section 3.1](#). The second is an application of recent results of [2] to obtain bounds on the treewidth of the graphs we consider ([Section 3.2](#)) and the last is an extension of the techniques in [10] fitting our needs, which is presented in [Section 3.3](#). [Section 2](#) contains definitions and preliminary results and [Section 4](#) discusses further research about the problem investigated in this paper.

## 2. Definitions and preliminaries

For any graph  $G$ ,  $V(G)$  (respectively  $E(G)$ ) denotes the *set of vertices* (respectively *edges*) of  $G$ . Even when dealing with multigraphs (i.e. graphs where more than one edge is allowed between two vertices) we will use the term *graph*. A graph  $G'$  is a *subgraph* of a graph  $G$  if  $V(G') \subseteq V(G)$  and  $E(G') \subseteq E(G)$ , and we denote this by  $G' \subseteq G$ . If  $X$  is a subset of  $V(G)$  (respectively  $E(G)$ ), we denote by  $G[X]$  the *subgraph of  $G$  induced by  $X$* , i.e. the graph with vertex set  $X$  (respectively  $\cup_{e \in X} e$ ) and edge set  $\{\{x, y\} \in E(G), x \in X \text{ and } y \in X\}$  (respectively  $X$ ). If  $S$  is a subset of vertices or edges of a graph  $G$ , the graph  $G \setminus S$  is the graph obtained from  $G$  after the removal of the elements of  $S$ . For every vertex  $v \in V(G)$  the neighborhood of  $v$  in  $G$ , denoted by  $N_G(v)$ , is the subset of vertices that are adjacent to  $v$ , and its size is called the *degree* of  $v$  in  $G$ , written  $\deg_G(v)$ . The maximum degree  $\Delta(G)$  of a graph  $G$  is the maximum value taken by  $\deg_G$  over  $V(G)$ . Given a non-negative integer  $k$ , a triple  $(V_1, S, V_2)$  is called a  *$k$ -separation triple* of a graph  $G$  if  $|S| \leq k$  and  $\{V_1, S, V_2\}$  is a partition of  $V(G)$  such that there is no edge between a vertex of  $V_1$  and a vertex of  $V_2$ . Unless otherwise stated, logarithms are binary. For any two integers  $a, b$  such that  $a \leq b$ , the notation  $\llbracket a, b \rrbracket$  stands for the set of integers  $\{a, a + 1, \dots, b\}$ . In a tree  $T$ , rooted at a vertex  $r \in V(T)$ , a vertex  $u \in V(T)$  is said to be a *descendant* of a vertex  $v \neq u$  if the path in  $T$  from  $r$  to  $u$  contains  $v$ . The set of descendants of  $v$  is denoted by  $\text{desc}_T(v)$ . A graph is *biconnected* if the removal of any vertex leaves the graph connected, and a biconnected component of a graph is a maximal biconnected subgraph.

*Minors and models.* In a graph  $G$ , a *contraction* of an edge  $e = \{u, v\} \in E(G)$  is the operation that removes  $e$  from  $G$  and identifies the vertices  $u$  and  $v$ . In this paper, we keep multiple edges that may appear between two vertices after a contraction (for instance, contracting an edge in a triangle gives a graph with two vertices connected by two edges). For any graph  $J$ , let  $\mathcal{M}(J)$  denote the class of *contraction models* (models for short) of  $J$ , i.e. the class of graphs that can be contracted to  $J$ . We say that a graph  $J$  is *minor* of a graph  $G$  (denoted by  $J \leq_m G$ ) if a subgraph of  $G$  is a model of  $J$  ( *$J$ -model* for short), or, equivalently, if  $J$  can be obtained from  $G$  by a series of vertex deletions, edge deletions, and edge contractions.

*Packings and coverings.* Let  $G$  and  $J$  be graphs. We denote by  $\text{pack}_J^v(G)$  the maximum number of vertex-disjoint models of  $J$  in  $G$  and by  $\text{cover}_J^v(G)$  the minimum size of a subset  $S \subseteq V(G)$  (called  $J$ -vertex-hitting set) that meets the vertex sets of all models of  $J$  in  $G$ . These invariants are widely studied in the context of the classic Erdős–Pósa property.

Similarly, we write  $\text{pack}_J^e(G)$  for the maximum number of edge-disjoint models of  $J$  in  $G$  and  $\text{cover}_J^e(G)$  for the minimum size of a subset  $S \subseteq E(G)$  (called  $J$ -edge-hitting set) that meets the edge sets of all models of  $J$  in  $G$ . Obviously, for every two graphs  $G$  and  $J$ , the following inequality holds:

$$\text{pack}_J^e(G) \leq \text{cover}_J^e(G).$$

A graph  $J$  is said to satisfy the (vertex-)Erdős–Pósa property for minors (vertex-Erdős–Pósa property for short) if there is a function  $f_J: \mathbb{N} \rightarrow \mathbb{N}$ , called vertex-Erdős–Pósa gap of  $J$ , such that for every graph  $G$ , the following holds:

$$\text{cover}_J^v(G) \leq f_J(\text{pack}_J^v(G)).$$

The research of this paper is motivated by the course of detecting graphs  $J$  for which there is a function  $h_J: \mathbb{N} \rightarrow \mathbb{N}$  satisfying the following inequality for every graph  $G$ :

$$\text{cover}_J^e(G) \leq h_J(\text{pack}_J^e(G)). \tag{1}$$

Such graphs are said to satisfy the edge variant of the Erdős–Pósa property for minors (or, in short, the edge-Erdős–Pósa property) and the function  $h_J$  is called the gap of the edge-Erdős–Pósa property for  $J$  (edge-Erdős–Pósa gap for short). This definition is an edge-counterpart to the existing Erdős–Pósa property and (vertex-)Erdős–Pósa gap.

*Treewidth.* A tree decomposition of a graph  $G$  is a pair  $(T, \mathcal{V})$  where  $T$  is a tree and  $\mathcal{V}$  a family  $(V_t)_{t \in V(T)}$  of subsets of  $V(G)$  (called bags) indexed by the vertices of  $T$  and such that

- (i)  $\bigcup_{t \in V(T)} V_t = V(G)$ ;
- (ii) for every edge  $e$  of  $G$  there is an element of  $\mathcal{V}$  containing both endpoints of  $e$ ; and
- (iii) for every  $v \in V(G)$ , the subgraph of  $T$  induced by  $\{t \in V(T) \mid v \in V_t\}$  is connected.

The width of a tree decomposition  $T$  is defined as  $\max_{t \in V(T)} |V_t| - 1$  (that is, the maximum size of a bag minus one). The treewidth of  $G$ , written  $\text{tw}(G)$ , is the minimum width of any of its tree decompositions.

A tree decomposition  $(T, \mathcal{V})$  of a graph  $G$  is said to be a nice tree decomposition if

- (i) every vertex of  $T$  has degree at most 3;
- (ii)  $T$  is rooted at one of its vertices  $r$  whose bag is empty ( $V_r = \emptyset$ ); and
- (iii) every vertex  $t$  of  $T$  is
  - either a base node, i.e. a leaf of  $T$  whose bag is empty ( $V_t = \emptyset$ ) and different from the root;
  - or an introduce node, i.e. a vertex with only one child  $t'$  such that  $V_t = V_{t'} \cup \{u\}$  for some  $u \in V(G)$ ;
  - or a forget node, i.e. a vertex with only one child  $t'$  such that  $V_{t'} = V_t \cup \{u\}$  for some  $u \in V(G)$ ;
  - or a join node, i.e. a vertex with two children  $t_1$  and  $t_2$  such that  $V_t = V_{t_1} = V_{t_2}$ .

It is known that every graph has an optimal tree decomposition which is nice [13].

*The graph  $\theta_r$  and the Erdős–Pósa property.* The vertex-Erdős–Pósa property of  $\theta_r$  received some attention, in particular in [7,9,11]. For instance the main result of [9] is the following estimation of the vertex-Erdős–Pósa gap for  $\theta_r$ .

**Proposition 2** ([9]). *For every positive integer  $r$ ,  $\theta_r$  has the vertex-Erdős–Pósa property with gap  $O(k^2)$ .*

However in this estimation the dependency in terms of  $r$  is hidden in the multiplicative constant of the Big-O notation. By a careful analysis of the size of a  $\theta_r$ -hitting set presented in [9] (c.f. Lemma 6), the estimation of the gap of Proposition 2 can be made quadratic in both  $k$  and  $r$ . From this, we can derive an  $O(k^3r^3)$  edge-Erdős–Pósa gap for  $\theta_r$  (Corollary 2) by using our Lemma 7 that makes possible to translate a  $\theta_r$ -vertex-hitting set into a  $\theta_r$ -edge-hitting set.

However, Theorem 1 gives better estimations of this gap, either in  $k$  or in  $r$ .

*Patterns in graphs of big treewidth.* In the following section, we will use several propositions asserting that every graph  $G$  of treewidth at least  $c_H$  contains some fixed graph  $H$  as a minor, where the constant  $c_H$  depends on  $H$ . For instance, we will show in Lemma 6 a simple relation between the constant  $c_{k,\theta_r}$  and the vertex-Erdős–Pósa gap for  $\theta_r$ . These propositions are stated thereafter.

**Proposition 3** ([17, Lemma 3.2]). *For every integer  $r \geq 1$  and graph  $G$ , if  $\text{tw}(G) \geq 2r - 1$  then  $G$  contains a  $\theta_r$ -model.*

**Proposition 4** ([9, Lemma 1], see also [1]). *Let  $k$  and  $r$  be two positive integers. For every graph  $G$ , if  $\text{tw}(G) \geq 2k^2r^2$  then  $G$  contains at least  $k$  vertex-disjoint models of  $\theta_r$ .*

**Proposition 5** ([2, Theorem 1.1]). *There is a function  $f_{\text{Proposition 5}}(t) = O(\text{polylog } t)$  such that, for every graph  $G$  and every positive integers  $h$  and  $p$ , if  $hp^2 \leq \frac{\text{tw}(G)}{f_{\text{Proposition 5}}(\text{tw}(G))}$ , there is a partition  $G_1, \dots, G_h$  of  $G$  into vertex-disjoint subgraphs such that  $\text{tw}(G_i) \geq p$  for each  $i \in \llbracket 1, h \rrbracket$ .*

**Proposition 6** ([2, Theorem 1.2]). *There is a function  $f_{\text{Proposition 6}}(t) = O(\text{polylog } t)$  such that, for every graph  $G$  and every positive integers  $h$  and  $p$ , if  $h^3p \leq \frac{\text{tw}(G)}{f_{\text{Proposition 6}}(\text{tw}(G))}$  then there is a partition  $G_1, \dots, G_h$  of  $G$  into vertex-disjoint subgraphs such that  $\text{tw}(G_i) \geq p$  for each  $i \in \llbracket 1, h \rrbracket$ .*




Fig. 2. A marked tree  $T$  with  $\mu(T) = 5$  and a 3-good partition  $(T_1, T_2)$  of root  $v$ . Marked vertices appear in white.

### 3. The edge-Erdős–Pósa property for graphs $\theta_r$

#### 3.1. Bounding the degree

In the sequel, we deal with graphs in which some vertices are marked. If  $G$  is a graph and  $m: V(G) \rightarrow \{0, 1\}$  is a function, we say that  $(G, m)$  is a graph *marked* by  $m$ . A vertex  $v$  of  $G$  such that  $m(v) = 1$  is said to be *marked*. We denote by  $\mu$  the function that, given a graph, returns its number of marked vertices. We now define an  $r$ -good partition. Given a positive integer  $r$ , a marked tree  $(T, m)$  is said to have an  $r$ -good partition of root  $v$  if there is a pair  $((T_1, m_1), (T_2, m_2))$  of marked trees such that:

- (i)  $T_1$  and  $T_2$  are subtrees of  $T$  such that  $(E(T_1), E(T_2))$  is a partition of  $E(T)$ ;
- (ii)  $r \leq \mu((T_1, m_1)) \leq 2r$ ;
- (iii)  $v \in V(T_2)$ ; and
- (iv) every vertex that is marked in  $(T, m)$  is either marked in  $(T_1, m_1)$  or marked in  $(T_2, m_2)$ , but not in both. In other words, for every  $u \in V(T)$ ,
  - if  $v \in V(T_1) \cap V(T_2)$  then  $m(v) = 1 \Leftrightarrow m_1(v) = 1$  or  $m_2(v) = 1$  but not both;
  - otherwise, let  $i \in \{1, 2\}$  be the integer such that  $v \in V(T_i)$ . Then we have  $m(v) = m_i(v)$ .

We remark that because of (iv),  $\mu(T) = \mu(T_1) + \mu(T_2)$ . If for every  $v \in V(T)$ ,  $(T, m)$  has an  $r$ -good partition of root  $v$ , then  $T$  is said to have an  $r$ -good partition. Examples of a marked tree and of a good partition are given in Fig. 2.

**Lemma 1.** For every integer  $r > 0$  and every marked tree  $(T, m)$ , if  $\mu(T) \geq 2r$  then  $(T, m)$  has an  $r$ -good partition.

**Proof.** Let  $r > 0$  be an integer. We prove this lemma by induction on the size of the tree.

*Base case:*  $|V(T)| = 0$ . Since  $2r \geq 2 > |V(T)|$ ,  $T$  does not have  $2r$  marked vertices and we are done.

*Induction step:* Assume that for every integer  $n' < n$ , every marked tree  $(T', m')$  on  $n'$  vertices and satisfying  $\mu((T', m')) \geq 2r$  has an  $r$ -good partition (induction hypothesis).

Let us prove that every marked tree on  $n$  vertices has a  $r$ -good partition if it has at least  $2r$  marked vertices. Let  $(T, m)$  be a tree on  $n$  vertices and let  $v$  be a vertex of  $T$ . We assume that  $\mu((T, m)) \geq 2r$ . We distinguish two cases.

- $\mu((T, m)) = 2r$ :

Let  $T_1 = T$ , let  $m_1 = m$ , let  $T_2 = (\{v\}, \emptyset)$ , and let  $m_2: V(T_2) \rightarrow \{0, 1\}$  be the function equal to 0 on every vertex of  $T_2$ . Remark that  $(E(T_1), E(T_2)) = (E(T), \emptyset)$  is a partition of  $E(T)$ ,  $T_2$  contains  $v$ , and as  $(T, m)$  contains (exactly)  $2r$  marked vertices, so does  $(T_1, m)$ . Consequently  $((T_1, m_1), (T_2, m_2))$  is an  $r$ -good partition of  $(T, m)$ .

- $\mu((T, m)) > 2r$ :

We distinguish different cases depending on the degree of the root  $v$  in  $T$ .

Case 1:  $\deg(v) = 1$ .

Let  $u$  be the neighbor of  $v$  in  $T$ , let  $T' = T \setminus \{v\}$ , and  $m' = m|_{V(T')}$ . Remark that  $\mu((T', m')) \geq 2r$  and  $|V(T')| = |V(T)| - 1$ . By induction hypothesis,  $(T', m')$  has an  $r$ -good partition  $((T'_1, m'_1), (T'_2, m'_2))$  of root  $u$ . We extend it to  $T$  by setting  $T_1 = T'_1$ ,  $m_1 = m'_1$ ,  $T_2 = (V(T'_2) \cup \{v\}, E(T'_2) \cup \{v, u\})$ , and  $m_2 = m'_2$ . Notice that  $T_2$  contains  $v$ . As the subtree  $T'_1$  contains at least  $r$  and at most  $2r$  marked vertices (induction hypothesis), so does  $T_1$ . Also, remark that  $(E(T_1), E(T_2))$  is a partition of  $E(T)$  and that since  $u \in T'_2$ , the graph  $T_2$  is connected. Therefore  $((T_1, m_1), (T_2, m_2))$  is an  $r$ -good partition of  $T$ .

Case 2:  $\deg(v) = d > 1$ .

Let  $u_1, \dots, u_d$  be the neighbors of  $v$  in  $T$  and for every  $i \in \llbracket 1, d \rrbracket$ , let  $C_i$  be the connected component of  $T \setminus \{v\}$  that contains  $u_i$ . We also define, for every  $i \in \llbracket 1, d \rrbracket$ , the restricted marking function  $w_i = m|_{V(C_i)}$ .

Subcase (a): there exists  $i \in \llbracket 1, d \rrbracket$  such that  $\mu((C_i, w_i)) > 2r$ .

Let  $T' = (V(C_i) \cup \{v\}, E(C_i) \cup \{u, v\})$  and let  $m' = m|_{V(T')}$ . Remark that  $|V(T')| < |V(T)|$  and  $\mu((T', m')) > 2r$ . According to the induction hypothesis,  $(T', m')$  has an  $r$ -good partition  $((T'_1, m'_1), (T'_2, m'_2))$  of root  $v$ . Similarly as before, we can extend it into an  $r$ -good partition  $((T_1, m_1), (T_2, m_2))$  of  $(T, m)$ . This is done by setting:

$$\begin{aligned} T_1 &= T'_1, \\ m_1 &= m'_1, \\ T_2 &= (V(T'_2) \cup (V(G) \setminus V(C_i)), E(G) \setminus E(T'_1)), \quad \text{and} \\ m_2 &: \begin{cases} u \mapsto m'_2(u) & \text{if } u \in V(C_i) \cup \{v\} \\ u \mapsto m(u) & \text{otherwise.} \end{cases} \end{aligned}$$

As  $((T'_1, m'_1), (T'_2, m'_2))$  is an  $r$ -good partition of root  $v$ ,  $v \in V(T'_2)$  and therefore  $T_2$  is connected.

Subcase (b): there exists  $i \in \llbracket 1, d \rrbracket$  such that  $r \leq \mu((C_i, w_i)) \leq 2r$ .

Let  $T_1 = C_i$  and  $T_2 = T[E(T) \setminus E(T_1)]$ . In this case,  $(E(T_1), E(T_2))$  is a partition of  $E(T)$  and  $T_2$  is connected since it contains  $v$ , the vertex which is adjacent to the  $C_j$ 's. Thus, if we set  $m_1 = m|_{V(T_1)}$  and  $m_2 = m|_{V(T_2)}$ ,  $((T_1, m_1), (T_2, m_2))$  is an  $r$ -good partition of  $(T, m)$ .

Subcase (c): for all  $i \in \llbracket 1, d \rrbracket$ ,  $\mu((C_i, w_i)) < r$ .

Let  $j = \min \left\{ j \in \llbracket 2, d \rrbracket, \sum_{i=1}^j \mu((C_i, w_i)) \geq r \right\}$ . Such value exists since  $\mu((T, m)) > 2r$ . We then set:

$$\begin{aligned} T_1 &= (\cup_{i \in \llbracket 1, j \rrbracket} V(C_i) \cup \{v\}, \cup_{i \in \llbracket 1, j \rrbracket} (E(C_i) \cup \{v, u_i\})), \\ m_1 &: \begin{cases} v \mapsto 0 \\ u \in V(T_1) \setminus \{v\} \mapsto m(u), \end{cases} \\ T_2 &= T[E(T) \setminus E(T_1)], \quad \text{and} \\ m_2 &= m|_{V(T_2)}. \end{aligned}$$

By definition of  $j$ ,  $\mu((T_1, m_1)) \geq r$  and as for every  $i \in \llbracket 1, d \rrbracket$ ,  $\mu((C_i, w_i)) < r$  we also have  $\mu((T_1, m_1)) < 2r$ . As before, the pair  $((T_1, m_1), (T_2, m_2))$  is an  $r$ -good partition of  $(T, m)$ .

In conclusion, we proved by induction that for every integer  $r$ , every tree having at least  $2r$  marked vertices has an  $r$ -good partition.  $\square$

In the sequel we will deal with packings of the graph  $\theta_r$ , for  $r > 1$ . The following remark is important.

**Remark 1.** If  $G$  is not biconnected, the number of edge-disjoint models of  $\theta_r$  in  $G$  is equal to the sum of the number of edge-disjoint models of  $\theta_r$  in every biconnected component of  $G$ . This enables us to treat biconnected components separately.

**Lemma 2.** Let  $k > 0, r > 0$  be two integers, and let  $G$  be a biconnected graph with  $\Delta(G) \geq 2kr$ . Then  $\text{pack}_{\theta_r}^e(G) \geq k$ .

**Proof.** As  $G$  is biconnected, the removal of a vertex  $v$  of maximum degree gives a connected graph. Let  $T$  be a minimal tree of  $G \setminus \{v\}$  spanning the neighborhood  $N_G(v)$  of  $v$ . We mark the vertices of  $T$  that are elements of  $N_G(v)$ : this gives the marking function  $m$  for  $T$ . Let us prove by induction on  $k$  that  $(T, m)$  has  $k$  edge-disjoint marked subtrees  $(T_1, m_1), \dots, (T_k, m_k)$ , each containing at least  $r$  marked vertices. If we do so, then we are done because  $\{\{v\}, T_i\}_{i \in \llbracket 1, k \rrbracket}$  is a collection of  $k$  edge-disjoint  $\theta_r$  models. In fact, as for every  $i \in \llbracket 1, k \rrbracket$ ,  $T_i$  contains  $r' \geq r$  vertices adjacent to  $v$  in  $G$ , contracting the edges of  $T_i$  in  $G \setminus \{v\} \cup V(T_i)$  gives the graph  $\theta_{r'}$ . Let  $r > 0$  be an integer.

Base case  $k = 1$ : Clear.

Induction step  $k > 1$ : Assume that for every  $k' < k$ , every tree with at least  $2k'r$  vertices marked has  $k'$  edge-disjoint subtrees, each with at least  $r$  marked vertices. Let  $(T, m)$  be a marked tree such that  $\mu((T, m)) \geq 2kr$ . According to Lemma 1,  $(T, m)$  has an  $r$ -good partition  $((T_1, m_1), (T'_1, m'_1))$  such that  $r \leq \mu((T_1, m_1)) \leq 2r$  and  $\mu((T'_1, m'_1)) = \mu((T, m)) - \mu((T_1, m_1)) \geq 2(k-1)r$ . By induction hypothesis,  $(T'_1, m'_1)$  has  $k-1$  edge-disjoint marked subtrees  $(T_2, m_2), \dots, (T_k, m_k)$  each containing at least  $r$  marked vertices. Remark that as all these trees are subgraphs of  $T'_1$ , which is edge-disjoint from  $T_1$  in  $T$ , they are edge-disjoint from  $T_1$  as well. Consequently,  $(T_1, m_1), (T_2, m_2), \dots, (T_k, m_k)$  is the family of edge-disjoint subtrees we were looking for.  $\square$

### 3.2. Bounding the treewidth

**Lemma 3.** There is a function  $h_r(k) = O(kr^2 \text{polylog } kr)$  such that for every positive integers  $k$  and  $r$  and every graph  $G$ , if  $\text{tw}(G) \geq h_r(k)$ , then  $\text{pack}_{\theta_r}^v(G) \geq k$ .

**Proof.** Let  $G$  be a graph and  $k, r$  be two positive integers. By Proposition 4, if  $\text{tw}(G) \geq 2k^2r^2$ , then  $G$  contains  $k$  vertex-disjoint models of  $\theta_r$ . Therefore, we only have to consider the case where  $\text{tw}(G) < 2k^2r^2$ .

As  $f_{\text{Proposition 5}}(t) = O(\text{polylog } t)$  (cf. Proposition 5 for the definition of  $f_{\text{Proposition 5}}$ ), there are three positive reals  $t_0, A \geq 1$ , and  $\alpha \geq 1$  such that for every real  $t \geq t_0$  we have  $f_{\text{Proposition 5}}(t) \leq A \log^\alpha(t)$ . Let  $B = \max(0, \max_{i \in \llbracket 1, \lceil t_0 \rceil \rrbracket} f_{\text{Proposition 5}}(i))$  and observe that for every positive integer  $i$  we have  $f_{\text{Proposition 5}}(i) \leq A \log^\alpha(i) + B$ .

Let  $h_r(k) = k(2r)^2 \cdot (A \log^\alpha(2k^2r^2) + B)$  for every positive integers  $k$  and  $r$ . Observe that  $h_r(k) = O(kr^2 \text{polylog } kr)$ . We will show that graphs whose treewidth is at least  $h_r(k)$  contain  $k$  vertex-disjoint models of  $\theta_r$ . For every positive integers  $r$  and  $k$ , if  $\text{tw}(G) \geq h_r(k)$  then we have

$$\begin{aligned} \text{tw}(G) &\geq k(2r)^2 \cdot (A \log^\alpha(\text{tw}(G)) + B) && \text{(as we assume } \text{tw}(G) < 2k^2r^2) \\ \frac{\text{tw}(G)}{A \log^\alpha(\text{tw}(G)) + B} &\geq k(2r)^2 && \text{(because } A \log^\alpha(\text{tw}(G)) + B \text{ is positive)} \\ \frac{\text{tw}(G)}{f_{\text{Proposition 5}}(\text{tw}(G))} &\geq k(2r)^2 && \text{(as } \text{tw}(G) \text{ is integer).} \end{aligned}$$

Notice that  $k$  and  $2r$  meet the conditions of Proposition 5. Consequently, there is a partition  $G_1, \dots, G_k$  of  $G$  into vertex-disjoint subgraphs such that  $\forall i \in \llbracket 1, k \rrbracket, \text{tw}(G_i) \geq 2r$ . By Proposition 3, each of these subgraphs contains a model of  $\theta_r$ . Consequently,  $G$  contains  $k$  vertex-disjoint models of  $\theta_r$ , as required.  $\square$

A very similar proof can be used to show the following lemma, using Proposition 6.

**Lemma 4.** *There is a function  $h_r(k) = O(k^3r \text{polylog } kr)$  such that, for every positive integers  $k$  and  $r$  and every graph  $G$ , if  $\text{tw}(G) \geq h_r(k)$ , then  $\text{pack}_{\theta_r}^V(G) \geq k$ .*

### 3.3. From vertices to edges

In this section, we show how an estimation of a vertex-Erdős–Pósa gap can be derived from the bound on the treewidth obtained in Section 3.2. The proof of the two following lemmas are inspired from the proof of [10, Lemma 2].

**Lemma 5** (Adapted from Lemma 2 of [10]). *Let  $k \geq 3, r$  be two positive integers and  $G$  a graph such that  $\text{pack}_{\theta_r}^V(G) = k$ . Then  $G$  has a  $(\text{tw}(G) + 1)$ -separation triple  $(V_1, S, V_2)$  such that  $\frac{1}{3}k \leq \text{pack}_{\theta_r}^V(G[V_1]) \leq \frac{2}{3}k$ .*

**Proof.** Let  $(T, \mathcal{V})$  be an optimal nice tree decomposition of  $G$ . For all  $t \in V(T)$ , let  $H_t$  be the subset of  $V(G)$  equal to  $(\bigcup_{t' \in \text{desc}_T(t)} V_{t'}) \setminus V_t$ , that is, informally, the subset of vertices that are in bags below  $V_t$  but not in  $V_t$ . We also define the function  $p: V(T) \rightarrow \mathbb{N}$  as:  $\forall t \in V(T), p(t) = \text{pack}_{\theta_r}^V(G[H_t])$ , which counts the number of vertex-disjoint models of  $\theta_r$  in the subgraph of  $G$  induced by  $H_t$ .

**Remark 2.** The function  $p$  is nondecreasing along every path from a vertex of  $T$  to the root of  $T$ , because if a vertex  $t' \in V(T)$  is a child of a vertex  $t \in V(T)$ , then  $H_{t'} \subseteq H_t$ , and thus  $\text{pack}_H^V(G[H_{t'}]) \leq \text{pack}_H^V(G[H_t])$ .

**Remark 3.** As  $T$  is a nice decomposition of  $G$ , its vertices can be of four different types. We make remarks about the value taken by  $p$  depending on the type of the vertices:

**Base node  $t$ :**  $p(t) = 0$ , because since  $t$  has no descendant,  $H_t = \emptyset$ ;

**Introduce node  $t$  with child  $t'$ :** as the unique element of  $V_t \setminus V_{t'}$  cannot appear in the bags of  $\text{desc}_T(t')$  (by definition of a tree decomposition),  $H_t = H_{t'}$  and then  $p(t) = p(t')$ ;

**Forget node  $t$  with child  $t'$ :**  $H_t$  contains one vertex more than  $H_{t'}$  therefore  $p(t) - p(t') \in \{0, 1\}$ ;

**Join node  $t$  with children  $t_1$  and  $t_2$ :**  $H_t = H_{t_1} \cup H_{t_2}$ , but  $H_{t_1}$  and  $H_{t_2}$  are disjoint and there is no edge between the vertices of  $H_{t_1}$  and of  $H_{t_2}$  in  $G[H_t]$  (otherwise the set  $V_{t_1} = V_{t_2}$  would contain an endpoint of this edge, which also belongs to  $H_{t_1}$  or  $H_{t_2}$ , and this is contradictory). Thus there is no  $\theta_r$ -model in  $G[H_t]$  that uses (simultaneously) vertices of  $H_{t_1}$  and of  $H_{t_2}$ , and therefore  $p(t) = p(t_1) + p(t_2)$ .

Let  $t \in V(T)$  be a node such that  $p(t) > \frac{2}{3}k$  and such that for every child  $t'$  of  $t, p(t') \leq \frac{2}{3}k$ . Let us make some claims about  $t$ .

**Claim 1.** *such a  $t$  exists.*

**Proof of Claim 1.** The value of  $p$  on the root  $r$  of  $T$  is  $k$  (because  $G[H_r] = G$ ) and according to the previous remark, the value of  $p$  on base nodes is 0. As  $p$  is nondecreasing on a path from a base node to the root (see Remark 2), such a vertex  $t$  exists.  $\diamond$

**Claim 2.**  *$t$  is unique.*

**Proof of Claim 2.** To show that  $t$  is unique, we assume by contradiction that there is another  $t' \in V(T)$  with  $t' \neq t$  and  $p(t') > \frac{2}{3}k$ , and such that for every child  $t''$  of  $t'$ ,  $p(t'') \leq \frac{2}{3}k$ . Three cases can occur:

- (i)  $t'$  is a descendant of  $t$ . However,  $p$  is nondecreasing along any path from a vertex to the root (Remark 2) and  $p(t') \geq \frac{2}{3}k$ , whereas the value of  $p$  for each child of  $t$  is at most  $\frac{2}{3}k$ : this is a contradiction.
- (ii)  $t$  is a descendant of  $t'$ . The same argument applies (symmetric situation).
- (iii)  $t$  and  $t'$  are not in the above situations. Let  $v$  be the least common ancestor of  $t$  and  $t'$ . As  $p$  is nondecreasing along any path from a vertex to the root, the child  $v_t$  (respectively  $v_{t'}$ ) of  $v$  of which  $t$  (respectively  $t'$ ) is descendant should be such that  $p(v_t) > \frac{2}{3}k$  (respectively  $p(v_{t'}) > \frac{2}{3}k$ ). By definition of  $v$ , we have  $v_t \neq v_{t'}$ . As  $v$  is a join node,  $p(v) = p(v_t) + p(v_{t'}) > \frac{4}{3}k$ , which is impossible.  $\diamond$

**Claim 3.**  $t$  is either a forget node or a join node.

**Proof of Claim 3.** By definition of  $t$ , the value  $p(t)$  is different from the value(s) taken by  $p$  over the child(ren) of  $t$ . This can only occur in the cases of a join node or a forget node.  $\diamond$

We now present a  $(\mathbf{tw}(G) + 1)$ -separation triple  $(V_1, S, V_2)$  of  $G$  with the required properties.

**Case 1:**  $t$  is a forget node with  $t'$  as child.

Let  $S = V_{t'}$ ,  $V_1 = H_{t'}$ , and  $V_2 = V(G) \setminus (V_1 \cup S)$ .

**Case 2:**  $t$  is a join node with  $t_1, t_2$  as children.

As  $\frac{2}{3}k < p(t) = p(t_1) + p(t_2)$  (Remark 3), there is  $i \in \{1, 2\}$  such that  $p(t_i) \geq \frac{k}{3}$ . Let  $S = V_{t_i}$ ,  $V_1 = H_{t_i}$ , and  $V_2 = V(G) \setminus (V_1 \cup S)$ .

In both cases, we have:

- (i)  $\frac{1}{3}k \leq \mathbf{pack}_{\theta_r}^v(G[V_1]) \leq \frac{2}{3}k$  by definition of  $V_1$  and  $t$ ;
- (ii)  $(V_1, S, V_2)$  is a partition of  $V(G)$ ;
- (iii) there is no edge between a vertex in  $V_1$  and a vertex of  $V_2$  (intuitively,  $S$  separates  $V_1$  and  $V_2$ ); and
- (iv)  $|S| \leq \mathbf{tw}(G) + 1$ , because  $S$  is a bag of an optimal tree decomposition of  $G$ .

In the case where  $t$  is a forget node, the inequality  $\frac{1}{3}k \leq \mathbf{pack}_{\theta_r}^v(G[V_1])$  of (i) holds because  $p(t') \geq p(t) - 1 > \frac{2}{3}k - 1 \geq \frac{k}{3}$  (cf. Remark 3). To see why (iii) is true, assume by contradiction that there are two vertices  $u \in V_1$  and  $v \in V_2$  such that  $\{u, v\} \in E(G)$ . Let  $s_0 \in V(T)$  be the child of  $t$  such that  $S = V_{s_0}$  (cf. the two different cases above). By definition of  $V_1$  there is a vertex  $s_1 \in V(T)$  of  $T$  in  $\text{desc}_T(s_0)$  whose bag  $V_{s_1}$  contains  $u$ . By definition of  $V_2$ , the vertex  $v$  does not belong to the bag  $V_{s_0}$  nor to a bag of a descendant of  $s_0$ . Let  $s_2$  be a vertex of  $T$  containing  $u$  and which is, according to the previous remark, not the bag of a descendant of  $s_0$  nor  $s_0$ .

As  $(T, \mathcal{V})$  is a tree decomposition of  $G$  and  $\{u, v\} \in E(G)$ , we have the following:

- there is a vertex  $s \in V(T)$  whose bag contains both  $u$  and  $v$ ;
- the subgraph of  $T$  induced by vertices whose bags contain  $u$  (respectively  $v$ ) is connected.

Consequently there is a path in  $T$  from  $s_1$  to  $s$  (respectively from  $s_2$  to  $s$ ) each bag of which contains  $u$  (respectively  $v$ ). As  $s$  is on the (only) path of  $T$  linking  $s_1$  to  $s_2$ , one of  $u, v$  belongs to the bag  $V_s$ . But this contradicts the fact that  $(V_1, S, V_2)$  is a partition of  $V(G)$ .

We conclude that  $(V_1, S, V_2)$  is a  $(\mathbf{tw}(G) + 1)$ -separation triple of  $G$  with the required properties.  $\square$

A function  $h: \mathbb{R} \rightarrow \mathbb{R}$  is said to be *superadditive* if for every  $x$  and every  $y$  in its domain,  $f(x) + f(y) \leq f(x + y)$ .

**Lemma 6** (Adapted from Lemma 5.4 in [2]). Let  $h_r$  be a superadditive function such that for every graph  $G$  and every positive integers  $r$  and  $k$ , if  $\mathbf{tw}(G) \geq h_r(k)$  then  $G \geq_{\mathbf{tw}}(k + 1) \cdot \theta_r$ . For every graph  $G$  and every positive integer  $k$ , if  $\mathbf{pack}_{\theta_r}^v(G) = k$  then we have

$$\mathbf{cover}_{\theta_r}^v(G) \leq 3 \cdot h_r(k) \log(k + 1).$$

**Proof.** We proceed by induction on  $k$ .

Base case  $k = 0$ : Clear.

Induction step  $k > 0$ : We assume that the lemma holds for every positive integer  $k' < k$ . Let  $G$  be a graph such that  $\mathbf{pack}_{\theta_r}^v(G) = k$ . First, remark that  $\mathbf{tw}(G) < h_r(k)$ , otherwise by definition of  $h_r$  we would have  $\mathbf{pack}_{\theta_r}^v(G) > k$ . Thus, by Lemma 5  $G$  contains a  $h_r(k)$ -separation triple  $(V_1, S, V_2)$  such that  $k/3 \leq \mathbf{pack}_{\theta_r}^v(G[V_1]) \leq 2k/3$ . This implies that  $k_1, k_2 \leq \lfloor 2k/3 \rfloor$ , where  $k_i = \mathbf{pack}_{\theta_r}^v(G[V_i])$  for every  $i \in \{1, 2\}$ . Also we have  $k_1 + k_2 \leq k$  as  $G_1$  and  $G_2$  are two vertex-disjoint subgraphs of  $G$ .



The triple  $(V_1, S, V_2)$  is a partition of  $V(G)$ , so the following holds:

$$\begin{aligned} \text{cover}_{\theta_r}^v(G) &\leq \text{cover}_{\theta_r}^v(G[V_1]) + \text{cover}_{\theta_r}^v(G[V_2]) + |S| \\ &\leq \text{cover}_{\theta_r}^v(G[V_1]) + \text{cover}_{\theta_r}^v(G[V_2]) + h_r(k) \\ &\leq 3 \cdot h_r(k_1) \log(k_1 + 1) + 3 \cdot h_r(k_2) \log(k_2 + 1) + h_r(k) \quad (\text{induction hyp.}). \end{aligned}$$

If  $k = 1$ , then  $k_1 = k_2 = 0$  and we have  $\text{cover}_{\theta_r}^v(G) \leq h_r(k) \leq 3 \cdot h_r(k) \log(k + 1)$ . We may now assume  $k \geq 2$ . Observe that in this case, as  $k_i \leq \lfloor \frac{2}{3}k \rfloor$ , we get  $k_i + 1 \leq \frac{3}{4}(k + 1)$  for every  $i \in \{1, 2\}$ .

$$\begin{aligned} \text{cover}_{\theta_r}^v(G) &\leq 3 \cdot (h_r(k_1) + h_r(k_2)) \log\left(\frac{3(k + 1)}{4}\right) + h_r(k) \\ &\leq 3 \cdot h_r(k) \log\left(\frac{3(k + 1)}{4}\right) + h_r(k) \quad (\text{superadditivity of } h_r) \\ &\leq 3 \cdot h_r(k) \log(k + 1) - 3 \cdot \log(4/3)h_r(k) + h_r(k) \\ &\leq 3 \cdot h_r(k) \log(k + 1). \end{aligned}$$

This concludes the proof.  $\square$

**Corollary 1.** Let  $f_r$  be the vertex-Erdős–Pósa gap of  $\theta_r$ . Then we have

- $f_r(k) = O(kr^2 \text{ polylog } kr)$ ;
- $f_r(k) = O(k^3r \text{ polylog } kr)$ .

These estimations follow from [Lemmas 3, 4 and 6](#).

The following lemma shows how to translate a vertex-Erdős–Pósa gap into an edge-Erdős–Pósa gap in the case of  $\theta_r$ . The main idea of the proof is that if the considered graph has small maximum degree, a small edge-hitting set can be constructed from a small vertex-hitting set. On the other hand, a big maximum degree forces a large packing of  $\theta_r$ -models.

**Lemma 7.** If  $f_r$  is the vertex-Erdős–Pósa gap of  $\theta_r$ , then the edge-Erdős–Pósa gap of  $\theta_r$  is less than  $2kr \cdot f_r(k)$ .

**Proof.** Let  $G$  be a graph, let  $r \geq 2$  be an integer and let  $f_r$  is the vertex-Erdős–Pósa gap of  $\theta_r$ .

We want to prove that if  $G$  contains less than  $k$  edge-disjoint models of  $\theta_r$ , then it has a  $\theta_r$ -edge-hitting set of size less than  $2kr \cdot f_r(k)$ .

According to [Remark 1](#), we can assume that  $G$  is biconnected. If it is not the case, we consider its biconnected components separately (if it has no biconnected component then the lemma is trivial).

First, remark  $\Delta(G) < 2kr$ , otherwise by [Lemma 2](#)  $G$  would contain at least  $k$  edge-disjoint  $\theta_r$ -models.

Notice that if  $G$  does not contain  $k$  edge-disjoint  $\theta_r$ -models, it does not contain  $k$  vertex-disjoint  $\theta_r$ -models either. Consequently, there is a set  $X \subseteq V(G)$  meeting every  $\theta_r$  model of  $G$  and such that  $|X| \leq f_r(k)$ . Let us consider the set  $Y \subseteq E(G)$  of edges incident to vertices of  $X$ , i.e.  $Y = \{\{u, v\} \in E(G), u \in X\}$ . Remark that as  $\Delta(G) < 2kr$ , we have  $|Y| \leq 2kr \cdot f_r(k)$ . Now, assume that there is a  $\theta_r$ -model in  $G$  not having edges in  $Y$ . None of its vertices is in  $X$ , which is contradictory. So  $Y$  is a  $\theta_r$ -edge hitting set of the required size. This concludes the proof.  $\square$

**Corollary 2.** An edge-gap of  $O(k^3r^3)$  for  $\theta_r$  can be derived from [Proposition 2](#).

**Proof of Theorem 1.** It follows from the application of [Lemma 7](#) to the estimations of the vertex-Erdős–Pósa gap of  $\theta_r$  given in [Corollary 1](#).

#### 4. Further research

The main question, initiated in this paper, is whether for every planar graph  $J$ , the class  $\mathcal{M}(J)$  satisfies this edge variant of the Erdős–Pósa property. As for the vertex version, it is easy to see that the planarity of  $J$  is necessary. For instance, if  $J = K_5$ , consider as graph  $G$  an  $n$ -vertex toroidal wall, which is a 3-regular graph embeddable in the torus that contains  $K_5$  as a minor. One can check that  $G$  does not contain two edge-disjoint models of  $K_5$ , but  $\Omega(\sqrt{n})$  edges of  $G$  are needed in order to hit all its  $K_5$ -models.

Moreover, a second question is: when this property holds, does it hold with a polynomial gap for all graphs? Also, finding lower bounds on this gap for specific graphs is another interesting and complementary question. Let us mention that, as it is the case for the vertex version (see [\[5,8\]](#)), for any non-acyclic planar graph  $J$  for which the edge variant of the Erdős–Pósa property holds for  $\mathcal{M}(J)$ , we have that  $f_{\mathcal{M}(J)}(k) = \Omega(k \log k)$ . Indeed, let  $G$  be an  $n$ -vertex cubic graph with treewidth  $\Omega(n)$  and girth  $\Omega(\log n)$  (such graphs are well-known to exist). Since  $J$  is planar, the treewidth of any graph excluding  $J$  as a minor is bounded by a constant [\[18\]](#), hence any set of edges of  $G$  meeting all models of  $J$  has size  $\Omega(n)$  (as the removal of an edge may decrease the treewidth by at most two). On the other hand, since  $J$  contains a cycle and the girth of  $G$  is  $\Omega(\log n)$ , any

model of  $J$  in  $G$  contains  $\Omega(\log n)$  edges (assuming that  $J$  does not have isolated vertices), and therefore  $G$  contains  $O(n/\log n)$  edge-disjoint models of  $J$  (here we have used that the degree of  $G$  is bounded), easily implying that  $f_{\mathcal{M}(J)}(k) = \Omega(k \log k)$ . In particular, it holds that  $f_{\mathcal{M}(\theta_r)}(k) = \Omega(k \log k)$  for any  $r \geq 2$ , so a first avenue for further work in this direction is to optimize the gap function  $f_{\mathcal{M}(\theta_r)}(k)$  given in [Theorem 1](#).

Finally, when the graphs  $G$  (in which the packings or coverings are taken) are restricted to classes of bounded degree, the proof of [Lemma 7](#) can easily be adapted to prove that the bound of the vertex version also holds for the edge version.

## References

- [1] E. Birmel e, J. Bondy, B. Reed, Brambles, prisms and grids, in: A. Bondy, J. Fonlupt, J.-L. Fouquet, J.-C. Fournier, J.L. Ram ırez Alfons ın (Eds.), *Graph Theory in Paris*, in: Trends in Mathematics, Birkh user, Basel, 2007, pp. 37–44.
- [2] C. Chekuri, J. Chuzhoy, Large-treewidth graph decompositions and applications, in: Proceedings of the Forty-fifth Annual ACM Symposium on Theory of Computing, STOC'13, ACM, New York, NY, USA, 2013, pp. 291–300.
- [3] C. Chekuri, J. Chuzhoy, Polynomial bounds for the grid-minor theorem, in: Proceedings of the 46th Annual ACM Symposium on Theory of Computing, STOC'14, ACM, New York, NY, USA, 2014, pp. 60–69.
- [4] R. Diestel, *Graph Theory*, third ed., in: Graduate Texts in Mathematics, vol. 173, Springer-Verlag, Heidelberg, 2005.
- [5] P. Erd os, L. P osa, On independent circuits contained in a graph, *Canad. J. Math.* 17 (1965) 347–352.
- [6] S. Fiorini, T. Huyhn, G. Joret, personal communication, 2013.
- [7] S. Fiorini, G. Joret, I. Sau, Optimal Erd os–P osa property for pumpkins, Manuscript, 2013.
- [8] S. Fiorini, G. Joret, D.R. Wood, Excluded forest minors and the Erd os–P osa property, *Combin. Probab. Comput.* 22 (5) (2013) 700–721.
- [9] F.V. Fomin, D. Lokshtanov, N. Misra, G. Philip, S. Saurabh, Quadratic upper bounds on the Erd os–P osa property for a generalization of packing and covering cycles, *J. Graph Theory* 74 (4) (2013) 417–424.
- [10] F.V. Fomin, S. Saurabh, D.M. Thilikos, Strengthening Erd os–P osa property for minor-closed graph classes, *J. Graph Theory* 66 (3) (2011) 235–240.
- [11] G. Joret, C. Paul, I. Sau, S. Saurabh, S. Thomass e, Hitting and harvesting pumpkins, *SIAM J. Discrete Math.* 28 (3) (2014) 1363–1390.
- [12] K.-I. Kawarabayashi, Y. Kobayashi, Edge-disjoint odd cycles in 4-edge-connected graphs, in: 29th International Symposium on Theoretical Aspects of Computer Science, STACS, 2012, pp. 206–217.
- [13] T. Kloks, *Treewidth. Computations and Approximations*, in: LNCS, vol. 842, Springer, 1994.
- [14] A. Leaf, P. Seymour, Treewidth and planar minors, Manuscript, 2012.
- [15] J.-F. Raymond, D. Thilikos, Polynomial gap extensions of the Erd os–P osa theorem, in: J. Ne et ıl, M. Pellegrini (Eds.), *The Seventh European Conference on Combinatorics, Graph Theory and Applications*, in: CRM Series, vol. 16, Scuola Normale Superiore, 2013, pp. 13–18.
- [16] J.-F. Raymond, D.M. Thilikos, Low Polynomial Exclusion of Planar Graph Patterns, ArXiv e-prints, May 2013.
- [17] B.A. Reed, D.R. Wood, Polynomial treewidth forces a large grid-like-minor, *European J. Combin.* 33 (3) (2012) 374–379.
- [18] N. Robertson, P.D. Seymour, Graph Minors. V. Excluding a planar graph, *J. Combin. Theory Ser. B* 41 (2) (1986) 92–114.