



HAL
open science

Antistrong digraphs

Jørgen Bang-Jensen, Stéphane Bessy, Bill Jackson, Matthias Kriesell

► **To cite this version:**

Jørgen Bang-Jensen, Stéphane Bessy, Bill Jackson, Matthias Kriesell. Antistrong digraphs. Journal of Combinatorial Theory, Series B, 2017, 122, pp.68-90. 10.1016/j.jctb.2016.05.004 . lirmm-01348862

HAL Id: lirmm-01348862

<https://hal-lirmm.ccsd.cnrs.fr/lirmm-01348862v1>

Submitted on 26 Jul 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Antistrong digraphs ^{*}

Jørgen Bang-Jensen[†] Stéphane Bessy[‡] Bill Jackson[§] Matthias Kriesell[¶]

April 2, 2015

Abstract

An antidirected trail in a digraph is a trail (a walk with no arc repeated) in which the arcs alternate between forward and backward arcs. An antidirected path is an antidirected trail where no vertex is repeated. We show that it is NP-complete to decide whether two vertices x, y in a digraph are connected by an antidirected path, while one can decide in linear time whether they are connected by an antidirected trail. A digraph D is antistrong if it contains an antidirected (x, y) -trail starting and ending with a forward arc for every choice of $x, y \in V(D)$. We show that antistrong connectivity can be decided in linear time. We discuss relations between antistrong connectivity and other properties of a digraph and show that the arc-minimal antistrong spanning subgraphs of a digraph are the bases of a matroid on its arc-set. We show that one can determine in polynomial time the minimum number of new arcs whose addition to D makes the resulting digraph the arc-disjoint union of k antistrong digraphs. In particular, we determine the minimum number of new arcs which need to be added to a digraph to make it antistrong. We use results from matroid theory to characterize graphs which have an antistrong orientation and give a polynomial time algorithm for constructing such an orientation when it exists. This immediately gives analogous results for graphs which have a connected bipartite 2-detachment. Finally, we study arc-decompositions of antistrong digraphs and pose several problems and conjectures.

Keywords: antidirected path, bipartite representation, matroid, detachment, anticonnected digraph

1 Introduction

We refer the reader to [1] for notation and terminology not explicitly defined in this paper. An **antidirected path** in a digraph D is a path in which the arcs alternate between forward and backward arcs. The digraph D is said to be **anticonnected** if it contains an antidirected path between x and y for every pair of distinct vertices x, y of D . Anticonnected digraphs were studied in [4], where several properties such as antihamiltonian connectivity have been considered. We will show in Theorem 2.2 below that it is NP-complete to decide whether a given digraph contains an antidirected path between given vertices.

Our main purpose is to introduce a related connectivity property based on the concept of a **forward antidirected trail**, i. e. a walk with no arc repeated which begins and ends with a forward arc and in which the arcs alternate between forward and backward arcs. A digraph D is **antistrong** if it has at least three vertices and contains a forward antidirected (x, y) -trail for every pair of distinct vertices x, y of D . We say that D is **k -arc-antistrong** if it has at least three vertices and contains k arc-disjoint forward antidirected (x, y) -trails for all distinct $x, y \in V(D)$.

^{*}This work was initiated while Bang-Jensen and Jackson attended the program “Graphs, hypergraphs and computing” at Institute Mittag-Leffler, spring 2014. Support from IML is gratefully acknowledged

[†]Department of Mathematics and Computer Science, University of Southern Denmark, Odense DK-5230, Denmark (email:bj@imada.sdu.dk). The research of Bang-Jensen was also supported by the Danish research council under grant number 1323-00178B

[‡]LIRMM, Université Montpellier, 161 rue Ada 34392 Montpellier Cedex 5 FRANCE (email:stephane.bessy@lirmm.fr)

[§]School of Mathematical Sciences, Queen Mary University of London, Mile End Road London E1 4NS, England (email:b.jackson@qmul.ac.uk)

[¶]Institut für Mathematik, Technische Universität Ilmenau, Weimarer Strasse 25, D-98693 Ilmenau, Germany

1 The paper is organized as follows. First we show that, from an algorithmic point of view, an-
2 ticonnectivity is not an easy concept to work with, since deciding whether a digraph contains an
3 anticonnected path between a given pair of vertices is NP-complete. Then we move to the main topic
4 of the paper, antistrong connectivity, and show that this relaxed version of anticonnectivity is easy to
5 check algorithmically. In fact, we show in Section 3 that there is a close relation between antistrong
6 connectivity of a digraph D and its so called bipartite representation $B(D)$, namely D is antistrong
7 if and only if $B(D)$ is connected. This allows us in Section 4 to find the minimum number of new
8 arcs we need to add to a digraph which is not antistrong so that the resulting digraph is antistrong.
9 Furthermore, using the bipartite representation we show in Section 5 that the arc-minimal antistrong
10 spanning subdigraphs of a digraph D form the bases of a matroid on the arc-set of D . More generally,
11 we show that the subsets of A which contain no closed antidirected trails are the independent sets
12 of a matroid on A . In Section 6 we study the problem of deciding whether a given undirected graph
13 has an antistrong orientation. We show how to reduce this problem to a matroid problem and give a
14 characterization of those graphs that have an antistrong orientation. Our proof leads to a polynomial
15 time algorithm which either finds an antistrong orientation of the given input graph G or produces a
16 certificate which shows that G has no such orientation. In Section 7 we show that being orientable as
17 an antistrong digraph can be expressed in terms of connected 2-detachments of graphs (every vertex v
18 is replaced by two copies v', v'' and every original edge uv becomes an edge between precisely one of the
19 4 possible pairs $u'v', u'v'', u''v', u''v''$) with the extra requirement that the 2-detachment is bipartite
20 and contains no edge of the form $u'v'$ or $u''v''$. This immediately leads to a characterization of graphs
21 having such a 2-detachment. Finally, in Section 8 we show that one can decide in polynomial time
22 whether a given digraph D has a spanning antistrong subdigraph D' so that $D - A(D')$ is connected in
23 the underlying sense (while it is NP-hard to decide whether a given digraph contains a non-separating
24 strong spanning subdigraph).
25 We conclude the paper with some remarks and open problems.

26 2 Anticonnectivity

27 It was shown in [4] that every connected graph G has an anticonnected orientation. This can be seen
28 by considering a breath first search tree rooted at some vertex r . Let $\{r\} = L_0, L_1, L_2, \dots, L_k$ be the
29 distance classes of G . Orient all edges between r and L_1 from r to these vertices, orient all edges
30 between $L_1 \cup L_3$ and L_2 from L_2 to $L_1 \cup L_3$, orient all edges from L_4 to $L_3 \cup L_5$ etc. Finally, orient
31 all the remaining, not yet oriented edges arbitrarily.

32 We will show that it is NP-complete to decide if a digraph has an antidirected path between two
33 given vertices. We need the following result which is not new, as it follows from a result in [8] on the
34 vertex analogue, but we include a new and short proof for completeness.

35 **Theorem 2.1** *It is NP-complete to decide for a given graph $G = (V, E)$, two specified vertices $x, y \in V$
36 and pairs of distinct edges $\mathcal{P} = \{(e_1, f_1), (e_2, f_2), \dots, (e_p, f_p)\}$, all from E , whether G has an (x, y) -
37 path which avoids at least one edge from each pair in \mathcal{P} .*

38 **Proof:** We first slightly modify a very useful polynomial reduction, used in many papers such
39 as [3], from 3-SAT to a simple path problem and then show how to extend this to a reduction from
40 3-SAT to the problem above. For simplicity our proof uses multigraphs but it is easy to change to
41 graphs.

42 Let $W[u, v, p, q]$ be the graph (the variable gadget) with vertices $\{u, v, y_1, y_2, \dots, y_p, z_1, z_2, \dots, z_q\}$
43 and the edges of the two (u, v) -paths $uy_1y_2 \dots y_pv, uz_1z_2 \dots z_qv$.

44 Let \mathcal{F} be an instance of 3-SAT with variables x_1, x_2, \dots, x_n and clauses C_1, C_2, \dots, C_m . The ordering
45 of the clauses C_1, C_2, \dots, C_m induces an ordering of the occurrences of a variable x and its negation
46 \bar{x} in these. With each variable x_i we associate a copy of $W[u_i, v_i, p_i + 1, q_i + 1]$ where x_i occurs p_i
47 times and \bar{x}_i occurs q_i times in the clauses of \mathcal{F} . Identify end vertices of these graphs by setting
48 $v_i = u_{i+1}$ for $i = 1, 2, \dots, n - 1$. Let $s = u_1$ and $t = v_n$ and denote by G' the resulting graph. In
49 G' we respectively denote by $y_{i,j}$ and $z_{i,j}$ the vertices y_j and z_j in the copy of W associated with the
50 variable x_i .

51 Next, for each clause C_i we associate this with 3 edges from G' as follows: assume C_i contains variables
52 x_j, x_k, x_l (negated or not). If x_j is not negated in C_i and this is the r th copy of x_j (in the order of the

1 clauses that use x_j), then we associate C_i with the edge $y_{j,r}y_{j,r+1}$ and if C_i contains $\overline{x_j}$ and this is
2 the k th occurrence of $\overline{x_j}$, then we associate C_i with the edge $z_{j,k}z_{j,k+1}$. We make similar associations
3 for the other two literals of C_i . Thus for each clause C_i we now have a set E_i of three distinct edges
4 $e_{i,1}, e_{i,2}, e_{i,3}$ from G' and $E_i \cap E_j = \emptyset$ for $i \neq j$.
5 Now it is easy to check that G' has an (s, t) -path which avoids at least one edge from each of the sets
6 E_1, E_2, \dots, E_m if and only if \mathcal{F} is satisfiable. Indeed, the (s, t) -path goes through the ‘ z -vertices’ of
7 the copy of W associated with x_i if and only if x_i is set to TRUE to satisfy \mathcal{F} .

8 Let us go back to the original problem. Let H be the multigraph consisting of vertices c_0, c_1, \dots, c_m
9 and three edges (denoted $f_{i,1}, f_{i,2}, f_{i,3}$) from c_{i-1} to c_i for $i \in \{1, \dots, m\}$. Let G denote the multigraph
10 we obtain from G' and H by identifying t and c_0 . Let $x = s$ and $y = c_m$. Finally, form three disjoint
11 pairs of arcs $(e_{i,1}, f_{i,1}), (e_{i,2}, f_{i,2}), (e_{i,3}, f_{i,3})$ between E_i and $\{f_{i,1}, f_{i,2}, f_{i,3}\}$ for every $i \in \{1 \dots m\}$.
12 By the observations above it is easy to check that G has an (x, y) -path which avoids at least one arc
13 from each of the forbidden pairs if and only if \mathcal{F} is satisfiable. \diamond

15 **Theorem 2.2** *It is NP-complete to decide whether a given digraph contains an antirected path*
16 *between given vertices x, y .*

17 **Proof:** The following proof is due to Anders Yeo (private communication, April 2014). Let
18 $G = (V, E)$ be a graph with two specified vertices $x, y \in V$ and pairs of distinct edges $\mathcal{P} =$
19 $\{(e_1, f_1), (e_2, f_2), \dots, (e_p, f_p)\}$, all from E . We will show how to construct a digraph D_G with specified
20 vertices s, t such that D_G contains an antirected (s, t) -path if and only if G has an (x, y) -path which
21 avoids at least one edge from each pair in \mathcal{P} . Since the construction can be done in polynomial time
22 this and Theorem 2.1 will imply the result.

23 Let k be the maximum number of pairs in \mathcal{P} involving the same edge from E . Let P be an
24 antirected path of length $2k + 2$ which starts with a forward arc (and hence ends with a backward
25 arc). Now construct D_G as follows: start from G and first replace every edge uv with a private copy
26 P_{uv} of P (no internal vertices are common to two such paths). Then for each pair $(e_i, f_i) \in \mathcal{P}$ we
27 identify one sink of P_{e_i} with one source of P_{f_i} so that the resulting vertex has in- and out-degree 2.
28 By the choice of the length of P we can identify in pairs, i. e. no three vertices will be identified. Note
29 that all the original vertices of G will be sources in D_G . The remaining (new vertices) will be called
30 *internal vertices*.

31 Finally let $s = x$ and $t = y$. We claim that D_G has an antirected (s, t) -path if and only
32 if G has an (x, y) -path which uses at most one edge from each of the pairs in \mathcal{P} . Suppose first
33 that $xx_1x_2 \dots x_{r-1}x_r y$ is a path in G which uses at most one edge from each of the pairs in \mathcal{P} .
34 Then $P_{xx_1}P_{x_1x_2} \dots P_{x_{r-1}x_r}P_{x_r y}$ is an antirected (s, t) -path in D_G (no vertex is repeated since the
35 identifications above were only done for paths corresponding to pairs in \mathcal{P}). Conversely, suppose
36 D_G contains an antirected (s, t) -path Q . By the way we identified vertex pairs according to \mathcal{P} , the
37 internal vertices have in- and out-degree at most 2, and if an internal vertex is on two paths P_{e_i}, P_{f_i}
38 then it has both its in-neighbours on P_{e_i} and both its out-neighbours on P_{f_i} . This implies that Q will
39 either completely traverse a path P_{e_i} or not touch any internal vertex of that path. Hence it cannot
40 traverse both P_{e_i} and P_{f_i} if $(e_i, f_i) \in \mathcal{P}$, and it follows that if we delete all internal vertices of Q and
41 add back the edges of G corresponding to each of the traversed paths, we obtain an (x, y) -path in G
42 that uses at most one edge from each pair in \mathcal{P} . \diamond

44 3 Properties of antistrong digraphs

45 For every digraph D we can associate an undirected bipartite graph which contains all the information
46 we need to study antistrong connectivity. The **bipartite representation** [1, Page 19] of a digraph
47 $D = (V, A)$ is the bipartite graph $B(D) = (V' \cup V'', E)$, where $V' = \{v' | v \in V\}$, $V'' = \{v'' | v \in V\}$
48 and $E = \{v'w'' | vw \in A\}$.

49 **Proposition 3.1** *Let $D = (V, A)$ be a digraph with $|V| \geq 3$. The following are equivalent.*

1 (a) D is antistrong

2 (b) $B(D)$ is connected.

3 (c) For every choice of distinct vertices x, y , the digraph D contains both an antidirected (x, y) -trail
4 $T_{x,y}$ of even length starting on a forward arc and an antidirected (x, y) -trail $\bar{T}_{x,y}$ of even length
5 starting on a backward arc.

6 **Proof:** Suppose (a) holds. Then, following the edges corresponding to the arcs of a forward
7 antidirected (x, y) -trail, $B(D)$ contains an (x', y'') -path for every pair of distinct vertices $x, y \in V$.
8 Now, if x and y are distinct vertices of D , we choose a third vertex z in D ($z \neq x$ and $z \neq y$),
9 and the union of an (x', z'') -path and a (z'', y') -path contains an (x', y') -path in $B(D)$. Similarly we
10 obtain an (x'', y'') -path in $B(D)$ for every pair of distinct vertices $x, y \in V$. Finally, for any $x \in V$,
11 an (x', x'') -path in $B(D)$ can be found in the union of an (x', y') -path and a (y', x'') -path, where y
12 is a vertex of D distinct from x . Hence (a) \Rightarrow (b) holds. Conversely, (b) \Rightarrow (a) holds, since any
13 (x', y'') -path in $B(D)$ corresponds to a forward antidirected (x, y) -path in D which starts and ends
14 with a forward arc.

15 Now to prove (b) \Rightarrow (c), it suffices to remark that $T_{x,y}$ and $\bar{T}_{x,y}$ correspond to an (x', y') -path and
16 an (x'', y'') -path in $B(D)$, respectively. Finally, to see that (c) \Rightarrow (b) holds, it suffices to show that
17 if (c) holds, then $B(D)$ contains an (x', y'') -path for all $x, y \in V$ (possibly equal). This follows by
18 considering a neighbour z'' of x' and a (z'', y'') -path in $B(D)$. \diamond

19
20 Proposition 3.1 implies the next two results.

21 **Corollary 3.2** *One can check in linear time whether a digraph is antistrong.*

22 **Corollary 3.3** *No bipartite digraph is antistrong.*

23 Recall that a digraph is **k-strong** if it has at least $k+1$ vertices and it remains strong after deletion
24 of any set of at most $k-1$ vertices. The digraph obtained from three disjoint independent sets X, Y, Z
25 each of size k by adding all arcs from X to Y , from Y to Z , and from Z to X is k -strong. However,
26 $B(D)$ has three connected components. This shows that no condition on the strong connectivity will
27 guarantee that a digraph is antistrong.

28
29 Recall that D is k -arc-antistrong if it contains k arc-disjoint forward antidirected (x, y) -trails for
30 every ordered pair of distinct vertices x, y . We can check in time $O(mk)$ whether a digraph has k
31 arc-disjoint forward antidirected (x, y) -trails for given vertices x, y , because they correspond to edge-
32 disjoint (x', y'') -paths in $B(D)$ whose existence can be checked by using flows, see e.g. [1, Section 5.5].
33 So we can check in polynomial time if a digraph is k -arc-antistrong.

34 **Theorem 3.4** *If D is $2k$ -arc-antistrong, then it contains k arc-disjoint antistrong spanning subdi-*
35 *graphs.*

36 **Proof:** Since D is $2k$ -arc-antistrong, $B(D)$ is $2k$ -edge-connected. We can now use Nash-Williams' theorem
37 (see [1, Theorem 9.4.2] for instance) to deduce that $B(D)$ has k edge-disjoint spanning trees.
38 Proposition 3.1 now gives the required set of k arc-disjoint antistrong spanning subdigraphs of D . \diamond

39
40 **Theorem 3.5** *There exists a polynomial time algorithm which for a given digraph D and a natural*
41 *number k either returns k arc-disjoint spanning antistrong subdigraphs of D or correctly answers that*
42 *no such set exists.*

43 **Proof:** This follows from the fact that such subdigraphs exist if and only if $B(D)$ has k edge-
44 disjoint spanning trees, and the existence of such trees can be checked via Edmonds' algorithm for
45 matroid partition [5]. \diamond

46
47 The corresponding problem for containing two arc-disjoint *strong* spanning subdigraphs is NP-
48 complete (see e.g. [1, Theorem 13.10.1]).

Theorem 3.6 *It is NP-complete to decide whether a digraph D contains two spanning strong subdigraphs D_1, D_2 which are arc-disjoint.*

4 Antistrong connectivity augmentation

Note that every complete digraph on at least 3 vertices is antistrong. Hence it is natural to ask for the minimum number of new arcs one has to add to a digraph in order to make it antistrong.

Theorem 4.1 *There exists a polynomial time algorithm for finding, for a given digraph $D = (V, A)$ on at least 3 vertices, a minimum cardinality set of new arcs F such that the digraph $D' = (V, A \cup F)$ is antistrong.*

Proof: Let D be a digraph on $n \geq 3$ vertices which is not antistrong. By Proposition 3.1, its bipartite representation $B(D)$ is not connected. First observe that in the bipartite representation each new arc added to D will correspond to an arc from a vertex u' of V' to a vertex $v'' \in V''$ such that $u \neq v$ back in V . So we are looking for the minimum number of new edges of type $u'v''$ with $u \neq v$ whose addition to $B(D)$ makes it connected while preserving the bipartition V', V'' . Note that, as long as $n \geq 3$, in which case $B(D)$ has at least 6 vertices, we can always obtain a connected graph by adding edges that are legal according to the definition above. So the number of edges we need is exactly the number of connected components of $B(D)$ minus one.¹

To find an optimal augmentation we add all missing edges between V' and V'' to $B(D)$, except for those of the form $v'v''$ and give the new edges cost 1, while all original edges get cost 0. Now find a minimum weight spanning tree in the resulting weighted complete bipartite graph. The edges of cost 1 correspond to an optimal augmenting set back in D . \diamond

The complexity of the analogous question for k -arc-antistrong connectivity is open.

Problem 4.2 *Given a digraph D and a natural number k , can we find in polynomial time a minimum cardinality set of new arcs whose addition to D results in a digraph D' which is k -arc-antistrong?*

Problem 4.2 is easily seen to be equivalent to the following problem on edge-connectivity augmentation of bipartite graphs.

Problem 4.3 *Given a natural number k and a bipartite graph $B = (X, Y, E)$ with $|X| = |Y| = p$ which admits a perfect matching M in its bipartite complement, find a minimum cardinality set of new edges F such that $F \cap M = \emptyset$ and $B + F$ is k -edge-connected and bipartite with the same bipartition as B .*

Theorem 4.1 can be extended to find the minimum number of new arcs whose addition to D gives a digraph with k arc-disjoint antistrong spanning subdigraphs D_1, \dots, D_k , provided that $V(D)$ is large enough to allow the existence of k such subdigraphs. Note that since each D_i needs at least $2n - 1$ arcs and we do not allow parallel arcs, we need n to be large enough, in particular we must have $n \geq 2k + 1$.

Theorem 4.4 *There exists a polynomial time algorithm for determining, for a given digraph D on at least 3 vertices, whether one can add some edges to D such that the resulting digraph is simple (no parallel arcs) and has k arc-disjoint antistrong spanning subdigraphs. In the case when such a set exists, the algorithm will return a minimum cardinality set of arcs A' such that $D' = (V, A \cup A')$ contains k arc-disjoint antistrong spanning subdigraphs.*

Proof: This follows from the fact that the minimum set of new arcs is exactly the minimum number of new edges, not of the form $v'v''$ that we have to add to $B(D)$ such that the resulting bipartite graph is simple and has k edge-disjoint spanning trees. This number can be found using matroid

¹This number is also equal to $(2n - 1) - r(A)$ where r is the rank function of the matroid $M(D)$ which we define in Section 5.

1 techniques as follows. Add all missing edges from V' to V'' and give those of the form $v'v''$ very
2 large cost (larger than $2nk$) and the other new edges cost 1. Now, if the resulting complete bipartite
3 digraph $K_{n,n}$ has k -edge-disjoint spanning trees of total cost less than $2nk$, then the set of new edges
4 added will form a minimum augmenting set and otherwise no solution exists. Recall from matroid
5 theory that k edge-disjoint spanning trees in $K_{n,n}$ correspond to k edge-disjoint bases in the cycle
6 matroid $M(K_{n,n})$ of $K_{n,n}$ which again corresponds to an independent set of size $k(2n - 1)$ in the
7 union $M = \bigvee_{i=1}^k M(K_{n,n})$. This means that we can solve the problem by finding a minimum cost
8 base B of M and then either return the arcs which correspond to edges of cost 1 in B or decide that
9 no solution exists when the cost of B is more than $2kn$. We leave the details to the reader. \diamond

11 5 A matroid for antistrong connectivity

12 Having seen the equivalence between antistrong connectivity of digraph D on n vertices and connec-
13 tivity of its bipartite representation $B(D)$ (see Proposition 3.1), and recalling from matroid theory
14 that $B(D)$ is connected if and only if the cycle matroid $M(B(D))$ has rank $|V(B(D))| - 1$, it is natural
15 to ask how antistrong connectivity can be expressed as a matroid property on D itself.

16 For $F \subseteq A$, we denote by $h(F)$ and $t(F)$ the numbers of vertices that are heads, respectively tails,
17 of one or more arcs in F .

Recall that the independent sets of the cycle matroid $M(G)$ of a graph $G = (V, E)$ are those
subsets $I \subseteq E$ for which we have $|I'| \leq \nu(I') - 1$ for all $\emptyset \neq I' \subseteq I$, where $\nu(I')$ is the number of
end vertices of the edges in I' . Inspired by this we define set I of arcs in a digraph $D = (V, A)$ to be
independent if

$$|I'| \leq h(I') + t(I') - 1 \quad \text{for all } \emptyset \neq I' \subseteq I, \quad (1)$$

18 A set $S \subseteq A$ is **dependent** if it is not independent.

19 **Proposition 5.1** *Let $D = (V, A)$ be a digraph. A subset $I \subseteq A$ is independent if and only if the*
20 *corresponding edge set I in $B(D)$ forms a forest. Every inclusion-minimal dependent set $S \subseteq A$*
21 *corresponds to a cycle in $B(D)$ and conversely.*

22 **Proof:** Suppose $I \subseteq A$ is independent and consider the corresponding edge set \tilde{I} in $B(D)$. If \tilde{I} is
23 not a forest, then some subset $\tilde{I}' \subseteq \tilde{I}$ will be a cycle C in $B(D)$ with p vertices in each of V', V'' for
24 some $p \geq 2$. The set \tilde{I}' corresponds to a set $I' \subseteq I$ with $h(I') + t(I') - 1 = p + p - 1 < 2p = |I'|$,
25 contradicting that I is independent. The other direction follows from the fact that every forest F in
26 $B(D)$ spans at least $|E(F)| + 1$ vertices in $B(D)$ and every subset of a forest is again a forest. The
27 last claim follows from the fact that every minimal set of edges which does not form a forest in $B(D)$
28 forms a cycle in $B(D)$. \diamond

30 The previous proposition implies that a set of arcs of a digraph is dependent if and only if it
31 contains a closed trail of even length consisting of alternating forward and backward arcs. We will
32 refer to such a trail as a **closed antidirected trail**, or **CAT** for short.

33 **Theorem 5.2** *Let $D = (V, A)$ be a digraph and \mathcal{I} be the family of all independent sets of arcs in D .*
34 *Then $M(D) = (A, \mathcal{I})$ is a graphic matroid with rank equal to the size of a largest collection of arcs*
35 *containing no closed alternating trail.*

36 **Proof:** It follows immediately from Proposition 5.1 that a set I belongs to \mathcal{I} if and only if the
37 corresponding edge set \tilde{I} is independent in the cycle matroid on $B(D)$. \diamond

39 **Theorem 5.3** *A digraph D is antistrong if and only if $M(B(D))$ has rank $2|V| - 1$.*

40 **Proof:** The rank of $M(B(D))$ equals the size of a largest acyclic set of edges in $B(D)$. This has size
41 $2|V| - 1$ precisely when $B(D)$ has a spanning tree H . Back in D , the arcs corresponding to $E(H)$
42 contain antidirected forward trails between any pair of distinct vertices. \diamond

6 Antistrong orientations of graphs

Recall that, by Robbins' theorem (see e.g. [1, Theorem 1.6.1]) a graph G has a strongly connected orientation if and only if G is 2-edge-connected. For antistrong orientations we have the following consequence of Proposition 3.1 which implies that there is no lower bound to the (edge-) connectivity which guarantees an antistrong orientation of a graph.

Proposition 6.1 *No bipartite graph can be oriented as an antistrong digraph.*

The purpose of this section is to characterize graphs which can be oriented as antistrong digraphs.

Theorem 6.2 *Suppose $G = (V, E)$ and $|E| = 2|V| - 1$. Then G has an antistrong orientation if and only if*

$$|E(H)| \leq 2|V(H)| - 1 \text{ for all nonempty subgraphs } H \text{ of } G, \text{ and} \quad (2)$$

$$|E(H)| \leq 2|V(H)| - 2 \text{ for all nonempty bipartite subgraphs } H \text{ of } G. \quad (3)$$

We derive Theorem 6.2 from the following characterization of graphs which can be oriented as digraphs with no closed antidirected trail (CAT).

Theorem 6.3 *A graph $G = (V, E)$ has an orientation with no CAT if and only if G satisfies (2) and (3). In particular no n vertex graph with at least $2n$ edges and no n vertex bipartite graph with at least $2n - 1$ edges admits a CAT-free orientation.*

It is not hard to see that Theorem 6.3 implies Theorem 6.2. Assume that Theorem 6.3 holds and consider a graph $G = (V, E)$ with $|E| = 2|V| - 1$. Suppose that G has an antistrong orientation D . Then $B(D)$ is connected by Proposition 3.1. As $B(D)$ has $2|V| - 1 = |V(B(D))| - 1$ edges it is a tree. So D is a CAT-free orientation of G and, by Theorem 6.3, conditions (2) and (3) hold for G . Conversely, if (2) and (3) hold for G , then G has a CAT-free orientation by Theorem 6.3, and we can deduce as above that this orientation is also an antistrong orientation of G .

We next show that (2) and (3) are necessary conditions for a CAT-free orientation. For the necessity of (2), suppose that some nonempty subgraph H has $|E(H)| \geq 2|V(H)|$ and that D is any orientation of G . Then $B(D)$ has at least $2|V(H)|$ edges between $V(H)'$ and $V(H)''$, implying that it contains a cycle. Hence D is not CAT-free. The necessity of (3) can be seen as follows. Suppose H is a bipartite subgraph on $2|V(H)| - 1$ edges and let \vec{H} be an arbitrary orientation of H . Since no bipartite graph has an antistrong orientation it follows that $B(\vec{H})$ is not connected, and, as it has $2|V(H)| - 1 = |V(B(\vec{H}))| - 1$ edges, it contains a cycle. This corresponds to a CAT in \vec{H} .

Most of the remainder of this section is devoted to a proof of sufficiency in Theorem 6.3. We first show that, for an arbitrary graph $G' = (V', E')$, the edge sets of all subgraphs G of G' which satisfy (2) and (3) are the independent sets of a matroid on E' . We then show that this matroid is the matroid union of the cycle matroid and the 'even bicircular matroid' of G' (defined below). This allows us to partition the edge-set of a graph G which satisfies (2) and (3) into a forest and an 'odd pseudoforest'. We then use this partition to define a CAT-free orientation of G . We first recall some results from matroid theory. We refer a reader unfamiliar with submodular functions and matroids to [7].

Suppose E is a set and $f : 2^E \rightarrow \mathbb{Z}$ is a submodular, nondecreasing set function which is nonnegative on $2^E \setminus \{\emptyset\}$. Edmonds [6], see [7, Theorem 13.4.2], showed that f induces a matroid M_f on E in which $S \subseteq E$ is independent if $|S'| \leq f(S')$ for all $\emptyset \neq S' \subseteq S$. The rank of a subset $S \subseteq E$ in M_f is given by the min-max formula

$$r_f(S) = \min_{\mathcal{P}} \left\{ \left| S \setminus \bigcup_{T \in \mathcal{P}} T \right| + \sum_{T \in \mathcal{P}} f(T) \right\}, \quad (4)$$

where the minimum is taken over all subpartitions \mathcal{P} of S (where a **subpartition** of S is a collection of pairwise disjoint nonempty subsets of S). Note that the matroid $M(D)$ defined in the previous section is induced on the arc-set of the digraph D by the set function $h + t - 1$.

1 Given a graph $G = (V, E)$ and $S \subseteq E$ let $G[S]$ be the **subgraph induced** by S i.e. the subgraph
2 of G with edge-set S and vertex-set all vertices incident to S . Let $\nu, \beta : 2^E \rightarrow \mathbb{Z}$ by putting $\nu(S)$ equal
3 to the number of vertices incident to S , and $\beta(S)$ equal to the number of bipartite components of
4 $G[S]$. It is well known that ν is submodular, nondecreasing, and nonnegative on 2^E and that $M_{\nu-1}(G)$
5 is the cycle matroid of G . The function $\nu - \beta$ is also known to be submodular, nondecreasing, and
6 nonnegative on 2^E and hence induces a matroid $M_{\nu-\beta}(G)$ on E which we call the **even bicircular**
7 **matroid** of G , see for example [11]. The independent sets of $M_{\nu-\beta}(G)$ are the edge sets of the **odd**
8 **pseudoforests** of G , i. e. subgraphs in which each connected component contains at most one cycle,
9 and if such a cycle exists then it is odd.

10 The above mentioned properties of ν and $\nu - \beta$ imply that $2\nu - 1 - \beta$ is submodular, nondecreasing,
11 and nonnegative on $2^E \setminus \{\emptyset\}$. We will show that the independent sets in $M_{2\nu-1-\beta}(G)$ are the edge
12 sets of the subgraphs which satisfy (2) and (3).

13 **Lemma 6.4** *Let $G = (V, E)$ be a graph and $\mathcal{I} = \{I \subseteq E : G[I] \text{ satisfies (2) and (3)}\}$. Then \mathcal{I} is the*
14 *family of independent sets of the matroid $M_{2\nu-1-\beta}(G)$. In addition, the rank of a subset $S \subseteq E$ in*
15 *this matroid is $r_{2\nu-1-\beta}(S) = \min_{\mathcal{P}} \{|S \setminus \bigcup_{T \in \mathcal{P}} T| + \sum_{T \in \mathcal{P}} (2\nu(T) - 1 - \beta(T))\}$ where the minimum*
16 *is taken over all subpartitions \mathcal{P} of S .*

17 **Proof.** We first suppose that some $S \subseteq E$ is not independent in $M_{2\nu-1-\beta}(G)$. Then we may choose
18 a nonempty $S' \subseteq S$ with $|S'| > 2\nu(S') - 1 - \beta(S')$, and subject to this condition, such that $|S'|$ is as
19 small as possible. The minimality of S' implies that $H = G[S']$ is connected. So $\beta(S') = 1$ if and only
20 if H is bipartite (and 0 otherwise) and we may now deduce that that $H \subseteq G[S]$ fails to satisfy (2) or
21 (3).

22 We next suppose that $G[S]$ fails to satisfy (2) or (3) for some $S \subseteq E$. Then there exists a nonempty
23 subgraph H of $G[S]$ such that either $|E(H)| > 2|V(H)| - 1$, or H is bipartite and $|E(H)| > 2|V(H)| - 2$.
24 Then $S' = E(H)$ satisfies $|S'| > 2\nu(S') - 1 - \beta(S')$ so S is not independent in $M_{2\nu-1-\beta}(G)$.

25 The expression for the rank function of $M_{2\nu-1-\beta}(G)$ follows immediately from (4). \diamond

26
27 The **matroid union** of two matroids $M_1 = (E, \mathcal{I}_1)$ and $M_2 = (E, \mathcal{I}_2)$ on the same ground
28 set E is the matroid $M_1 \vee M_2 = (E, \mathcal{I})$ where $\mathcal{I} = \{I_1 \cup I_2 : I_1 \in \mathcal{I}_1 \text{ and } I_2 \in \mathcal{I}_2\}$. Suppose
29 $f_1, f_2 : E \rightarrow \mathbb{Z}$ are submodular, nondecreasing, and nonnegative on $2^E \setminus \{\emptyset\}$. Then $f_1 + f_2$ will also be
30 submodular, nondecreasing, and nonnegative on $2^E \setminus \{\emptyset\}$ and hence will induce the matroid $M_{f_1+f_2}$.
31 Every independent set in $M_{f_1} \vee M_{f_2}$ is independent in $M_{f_1+f_2}$, but the converse does not hold in
32 general. Katoh and Tanigawa [9, Lemma 2.2] have shown that the equality $M_{f_1+f_2} = M_{f_1} \vee M_{f_2}$ does
33 hold whenever the minimum in formula (4) for the ranks $r_{f_1}(S)$ and $r_{f_2}(S)$ is attained for the same
34 subpartition of S , for all $S \subseteq E$. This allows us to deduce

35 **Lemma 6.5** *For any graph $G = (V, E)$, we have $M_{2\nu-1-\beta}(G) = M_{\nu-1}(G) \vee M_{\nu-\beta}(G)$.*

36 **Proof.** This follows from the above mentioned result of Katoh and Tanigawa, and the facts that
37 $r_{\nu-1}(S) = \sum_{T \in \mathcal{P}} (\nu(T) - 1)$ and $r_{\nu-\beta}(S) = \sum_{T \in \mathcal{P}} (\nu(T) - \beta(T))$ where \mathcal{P} is the partition of S given
38 by the connected components of $G[S]$ (since $r_{\nu-1}(S)$ and $r_{\nu-\beta}(S)$ are equal to the number of edges
39 in a maximum forest and a maximum odd pseudoforest, respectively, in $G[S]$). \diamond

40

41 Lemma 6.4 and Lemma 6.5 immediately give the following.

42 **Lemma 6.6** *Let $G = (V, E)$ be a graph. Then G satisfies (2) and (3) if and only if E can be*
43 *partitioned into a forest and an odd pseudoforest.*

44 We provide an alternative graph theoretic proof of this lemma in the Appendix.

45

46 We next show that every graph whose edge set can be partitioned into a spanning tree and an
47 odd pseudoforest has a CAT-free orientation.




Figure 1: A CAT-free orientation of the union of a spanning tree T and a spanning pseudoforest P ; T governs the bipartition X, Y (white/grey), its edges are drawn outside (or on) the disk spanned by the vertices. The edges of P are embedded in the interior of that disk, the root vertex is the encircled topmost one, the precious edge is the dashed one.

1 **Theorem 6.7** Let G be the edge-disjoint union of a spanning tree T and an odd pseudoforest P .²
 2 Then G has a CAT-free orientation. In addition, such an orientation can be constructed in linear
 3 time given T and P .

4 **Proof:** Let X, Y be the unique (up to renaming the two sets) bipartition of T and orient all edges
 5 of T from Y to X . If P has no edges we are done since there are no cycles in G . Let P_1, \dots, P_k be
 6 the connected components of P . We shall show that we can orient the edges of P_1, \dots, P_k in such a
 7 way that none of the resulting arcs of these (now oriented) pseudoforests $\vec{P}_1, \dots, \vec{P}_k$ can belong to a
 8 closed antidirected trail. Clearly this will imply the lemma. For each P_i we choose a root vertex r_i of
 9 P_i as follows. If P_i is a tree then we choose r_i to be an arbitrary vertex of P_i . If P_i contains an odd
 10 cycle C_i then we choose r_i to be a vertex of C_i such that r_i has at least one neighbour $s_i \in C_i$ which
 11 belongs to the same set in the bipartition (X, Y) as r_i (this is possible since C_i is odd). We will refer
 12 to the edge $r_i s_i$ as a **precious edge** of P_i . Put $T_i = P_i - r_i s_i$ if P_i contains a cycle and otherwise
 13 put $T_i = P_i$.

14 We orient the edges of T_i as follows. Every edge of P_i with one end in X and the other in Y is
 15 oriented from X to Y . Every edge uv of T_i with $u, v \in X$ is oriented towards r_i in T_i (so if v is closer
 16 to r_i than u in T_i we orient the edge from u to v and otherwise we orient it from v to u , see Figure 1).
 17 Every edge pq of T_i with $p, q \in Y$ is oriented away from r_i in T_i . Finally, if P_i contains a precious
 18 edge $r_i s_i$, then we orient $r_i s_i$ from r_i to s_i if $r_i, s_i \in X$, and from s_i to r_i if $r_i, s_i \in Y$. Let D denote
 19 the resulting orientation of G . The digraph D can be constructed in linear time if we traverse each
 20 tree T_i by a breath first search rooted at r_i .

21 We use induction on $|E(P)|$ to show that D is CAT-free. As noted above, this is true for the base
 22 case when $E(P) = \emptyset$. Suppose that $E(P) \neq \emptyset$ and choose an edge uv in some P_i according to the
 23 following criteria. If P_i is not a cycle then choose v to be a vertex of degree one in P_i distinct from
 24 r_i and u to be the neighbour of v in P_i . If P_i is an odd cycle then choose $v = r_i$ and $u = s_i$. We will
 25 show that uv belongs to no CAT in D . By symmetry, we may suppose that $v \in X$.

²Note that G may have parallel edges, but no more than two copies of any edge, in which case one copy is in T and the other in P

We first consider the case when v is a vertex of degree one in P_i . Below $d^+(v), d^-(v)$ denote the out-degree, respectively, the in-degree of the vertex v . We have two possible subcases:

- $u \in Y$. Since $v \in X$, we oriented uv from v to u . All the other edges incident to v belong to T and were oriented towards v . Then $d^+(v) = 1$ and the arc vu cannot be part of a CAT.
- $u \in X$. Since $v \in X$, we oriented uv from v to u (as u is closer to r_i than v in T_i). As previously we have $d^+(v) = 1$ and the arc vu cannot be part of a CAT.

Since $D - uv$ is CAT-free by induction, D is also CAT-free.

We next consider the case when P_i is an odd circuit, $v = r_i$ and $u = s_i$. Let t_i be the neighbour of r_i in P_i distinct from s_i . We again have two possible subcases:

- $t_i \in X$. Since $r_i \in X$, we oriented the edge $t_i r_i$ from t_i to r_i . Then $d^+(r_i) = 1$, and the arc $r_i s_i$ cannot be part of a CAT.
- $t_i \in Y$. Let q_i be the neighbour of s_i in P_i which is distinct from r_i . The choice of r_i implies that $q_i \in Y$, and hence that $s_i q_i$ is oriented from s_i to q_i . Then $d^+(s_i) = 1$, and the arc $r_i s_i$ cannot be part of a CAT.

Since $D - r_i s_i$ is CAT-free by induction, D is also CAT-free. ◇

Proof of Theorem 6.3 (sufficiency): Let $G = (V, E)$ be a graph satisfying (2) and (3). By Lemma 6.6, E can be partitioned into a forest F and an odd pseudoforest P . By adding a suitable set of edges to G , we may assume that $|E| = 2|V| - 1$. (This follows by considering the matroid $M_{2\nu-1-\beta}(2K_n)$ on the edge set of the graph $2K_n$ with vertex set V in which all pairs of vertices are joined by two parallel edges. It is easy to check that $2K_n$ has an edge-disjoint forest and odd pseudoforest with a total of $2|V| - 1$ edges. Thus the rank of $M_{2\nu-1-\beta}(2K_n)$ is $2|V| - 1$. Since E is an independent set in $M_{2\nu-1-\beta}(2K_n)$, it can be extended to an independent set with $2|V| - 1$ edges.) The fact that $|E| = 2|V| - 1$ implies that F is a spanning tree of G . We can now apply Theorem 6.7 to deduce that G has a CAT-free orientation. ◇

We have seen that Theorem 6.3 implies Theorem 6.2, and hence that a graph $G = (V, E)$ has an antistrong orientation if and only if the rank of $M_{2\nu-1-\beta}(G)$ is equal to $2|V| - 1$. We can now apply the rank formula (4) to characterize graphs which admit an antistrong orientation.

Theorem 6.8 *A graph $G = (V, E)$ has an antistrong orientation if and only if*

$$e(\mathcal{Q}) \geq |\mathcal{Q}| - 1 + b(\mathcal{Q}) \tag{5}$$

for all partitions \mathcal{Q} of V , where $e(\mathcal{Q})$ denotes the number of edges of G between the different parts of \mathcal{Q} and $b(\mathcal{Q})$ the number of parts of \mathcal{Q} which induce bipartite subgraphs of G .

Proof: Suppose that G has no antistrong orientation. Then the rank of $M_{2\nu-1-\beta}(G)$ is less than $2|V| - 1$ so there exists a subpartition \mathcal{P} of E such that

$$\alpha(\mathcal{P}) := \left| E \setminus \bigcup_{T \in \mathcal{P}} T \right| + \sum_{T \in \mathcal{P}} (2\nu(T) - 1 - \beta(T)) < 2|V| - 1. \tag{6}$$

We may assume that \mathcal{P} has been chosen such that:

- (i) $\alpha(\mathcal{P})$ is as small as possible;
- (ii) subject to (i), $|\mathcal{P}|$ is as small as possible;
- (iii) subject to (i) and (ii), $|\bigcup_{T \in \mathcal{P}} T|$ is as large as possible.

1 Let $\mathcal{P} = \{E_1, E_2, \dots, E_t\}$ and let $H_i = (V_i, E_i)$ be the subgraph of G induced by E_i for all $1 \leq i \leq t$.
 2 We will show that H_i is a (vertex-)induced connected subgraph of G and that $V_i \cap V_j = \emptyset$ for all $i \neq j$.

First, suppose that H_i is disconnected for some $1 \leq i \leq t$. Then we have $H_i = H'_i \cup H''_i$ for some subgraphs $H'_i = (V'_i, E'_i)$ and $H''_i = (V''_i, E''_i)$ with $V'_i \cap V''_i = \emptyset$. Let $\mathcal{P}' = (\mathcal{P} \setminus \{E_i\}) \cup \{E'_i, E''_i\}$. We have

$$2\nu(E_i) - 1 - \beta(E_i) > 2\nu(E'_i) - 1 - \beta(E'_i) + 2\nu(E''_i) - 1 - \beta(E''_i)$$

3 since, $\nu(E_i) = \nu(E'_i) + \nu(E''_i)$ and $\beta(E_i) = \beta(E'_i) + \beta(E''_i)$. This implies that $\alpha(\mathcal{P}') < \alpha(\mathcal{P})$ and
 4 contradicts (i). Hence H_i is connected and $\beta(E_i) \in \{0, 1\}$ for all $1 \leq i \leq t$.

Next, suppose that $V_i \cap V_j \neq \emptyset$ for some $1 \leq i < j \leq t$. Let $\mathcal{P}' = (\mathcal{P} \setminus \{E_i, E_j\}) \cup \{E_i \cup E_j\}$. We have

$$2\nu(E_i) - 1 - \beta(E_i) + 2\nu(E_j) - 1 - \beta(E_j) \geq 2\nu(E_i \cup E_j) - 1 - \beta(E_i \cup E_j)$$

5 since, if $|V_i \cap V_j| = 1$, then $\nu(E_i) + \nu(E_j) = \nu(E_i \cup E_j) + 1$ and $\beta(E_i) + \beta(E_j) \leq \beta(E_i \cup E_j) + 1$,
 6 and, if $|V_i \cap V_j| \geq 2$, then $\nu(E_i) + \nu(E_j) \geq \nu(E_i \cup E_j) + 2$ and $\beta(E_i) + \beta(E_j) \leq 2$. This implies that
 7 $\alpha(\mathcal{P}') \leq \alpha(\mathcal{P})$. Since $|\mathcal{P}'| < |\mathcal{P}|$ this contradicts (i) or (ii). Hence $V_i \cap V_j = \emptyset$ for all $i \neq j$.

8 Finally, suppose that $H_i \neq G[V_i]$. Then some $e \in E \setminus \bigcup_{T \in \mathcal{P}} T$ has both end vertices in E_i .
 9 Let $E'_i = E_i + e$ and $\mathcal{P}' = \mathcal{P} - E_i + E'_i$. This implies that $\alpha(\mathcal{P}') \leq \alpha(\mathcal{P})$. Since $|\mathcal{P}'| = |\mathcal{P}|$ and
 10 $|\bigcup_{T \in \mathcal{P}'} T| > |\bigcup_{T \in \mathcal{P}} T|$, this contradicts (i) or (iii). Hence $H_i = G[V_i]$.

11 Let \mathcal{Q} be the partition of V obtained from $\{V_1, V_2, \dots, V_t\}$ by adding the remaining vertices of G
 12 as singletons. Then $|E \setminus \bigcup_{T \in \mathcal{P}} T| = e(\mathcal{Q})$ and $\sum_{T \in \mathcal{P}} (2\nu(T) - 1 - \beta(T)) = 2|V| - |\mathcal{Q}| - b(\mathcal{Q})$. We
 13 can now use (6) to deduce that $e(\mathcal{Q}) < |\mathcal{Q}| - 1 + b(\mathcal{Q})$.

Suppose, on the other hand, that $e(\mathcal{Q}) < |\mathcal{Q}| - 1 + b(\mathcal{Q})$ for some partition $\mathcal{Q} = \{V_1, V_2, \dots, V_s\}$ of
 V . Let $G[V_i] = (V_i, E_i)$ for $1 \leq i \leq s$ and $\mathcal{P} = \{E_i : E_i \neq \emptyset, 1 \leq i \leq s\}$. Then $|E \setminus \bigcup_{T \in \mathcal{P}} T| = e(\mathcal{Q})$
 and

$$\sum_{T \in \mathcal{P}} (2\nu(T) - 1 - \beta(T)) = 2|V| - |\mathcal{Q}| - b(\mathcal{Q}) - \sum_{T \in \mathcal{P}} (\beta(T) - \beta^*(T)) \leq 2|V| - |\mathcal{Q}| - b(\mathcal{Q}),$$

where $\beta^*(T) = \min\{\beta(T), 1\}$. A straightforward calculation now gives

$$\left| E \setminus \bigcup_{T \in \mathcal{P}} T \right| + \sum_{T \in \mathcal{P}} (2\nu(T) - 1 - \beta(T)) < 2|V| - 1$$

14 and hence G has no antistrong orientation. ◇

15

16

17 **Corollary 6.9** *Every 4-edge-connected nonbipartite graph has an antistrong orientation.*

18 **Proof:** Suppose $G = (V, E)$ is 4-edge-connected and not bipartite and let \mathcal{Q} be a partition of V . If
 19 $\mathcal{Q} = \{V\}$ then $e(\mathcal{Q}) = 0 = |\mathcal{Q}| - 1 + b(\mathcal{Q})$ since G is not bipartite, and if $\mathcal{Q} \neq \{V\}$ then $e(\mathcal{Q}) \geq 2|\mathcal{Q}| \geq$
 20 $|\mathcal{Q}| - 1 + b(\mathcal{Q})$ since G is 4-edge-connected. Hence G has an antistrong orientation by Theorem 6.8. ◇

21

22 **Corollary 6.10** *Every nonbipartite graph with three edge-disjoint spanning trees has an antistrong*
 23 *orientation.*

24 **Proof:** We give two proofs of this corollary.

25 Suppose $G = (V, E)$ is a nonbipartite graph with three edge-disjoint spanning trees and let \mathcal{Q}
 26 be a partition of V . If $\mathcal{Q} = \{V\}$ then $e(\mathcal{Q}) = 0 = |\mathcal{Q}| - 1 + b(\mathcal{Q})$ since G is not bipartite, and if
 27 $\mathcal{Q} \neq \{V\}$ then $e(\mathcal{Q}) \geq 3(|\mathcal{Q}| - 1)$ since G has three edge-disjoint spanning trees. Since $|\mathcal{Q}| \geq 2$,
 28 $2(|\mathcal{Q}| - 1) \geq |\mathcal{Q}| \geq b(\mathcal{Q})$ and $e(\mathcal{Q}) \geq |\mathcal{Q}| - 1 + b(\mathcal{Q})$. Hence G has an antistrong orientation by
 29 Theorem 6.8.

30 We could also remark that if T_1, T_2 and T_3 denote three edge-disjoint spanning trees of G , then
 31 there exists $e \in G$ such that $T_1 + e$ is not bipartite. Then depending if $e \in T_2$ or not, $\{T_1 + e, T_3\}$ or

1 $\{T_1 + e, T_2\}$ is an edge-disjoint pair of a spanning odd pseudo-tree and a spanning tree of G . Let H
2 denote this subgraph of G . Then using Theorem 6.7, H has a CAT-free orientation which is also an
3 antistrong orientation of H since $|E(H)| = 2|V(H)| - 1$. So G has also an antistrong orientation. \diamond

5 Corollary 6.10 is tight in the sense that there exist graphs with many edge-disjoint trees, two
6 spanning and the others missing just three vertices, which have no antistrong orientation. Consider
7 the graph G obtained by identifying one vertex of a complete bipartite graph $K_{k,k}$ and a complete
8 graph K_4 . Then G has no antistrong orientation. Indeed, consider the partition \mathcal{Q} of $V(G)$ into
9 four parts: the copy of $K_{k,k}$, and one part for each remaining vertex of K_4 . We have $e(\mathcal{Q}) = 6 <$
10 $|\mathcal{Q}| - 1 + b(\mathcal{Q}) = 4 - 1 + 4$ and then G has no antistrong orientation by Theorem 6.8.

11 Since $M_{2\nu-1-\beta}(G) = M_{\nu-1}(G) \vee M_{\nu-\beta}(G)$, we can use Edmonds' matroid partition algorithm [5]
12 to determine the rank of $M_{2\nu-1-\beta}(G)$ in polynomial time, and hence determine whether G has an
13 antistrong orientation. Moreover, when such an orientation exists, we can use Edmonds' algorithm
14 to construct an edge-disjoint spanning tree and pseudoforest with a total of $2|V| - 1$ edges, and then
15 use the construction from the proof of Theorem 6.7 to obtain the desired antistrong orientation in
16 polynomial time. This gives

17 **Corollary 6.11** *There exists a polynomial time algorithm which finds, for a given input graph G ,*
18 *either an antistrong orientation D of G , or a certificate, in terms of a subpartition \mathcal{P} which violates*
19 *(5), that G has no such orientation.*

20 7 Connected bipartite 2-detachments of graphs

21 We now show a connection between antistrong orientations of a graph G and so-called detachments
22 of G . We need only the special case of 2-detachments (see e.g. [10] for results on detachments).

23 A **2-detachment** of a graph $G = (V, E)$ is any graph $H = (V' \cup V'', E')$ which can be obtain
24 from G by replacing every vertex $v \in V$ with two new vertices v', v'' and then for each original edge
25 uv adding precisely one of the four edges $u'v', u'v'', u''v', u''v''$ to E' .

26 **Lemma 7.1** *A graph $G = (V, E)$ has an antistrong orientation if and only if G has a 2-detachment*
27 *$H = (V' \cup V'', E')$ which is connected and bipartite with bipartition V', V'' (we call such a 2-detachment*
28 *good).*

29 **Proof:** Suppose G has a good 2-detachment $H = (V' \cup V'', E')$. Then there are no edges of the form
30 $u'v'$ and no edges of the form $u''v''$. Hence the orientation D that we get by orienting the edges of
31 the form $u'v''$ from u to v will be an antistrong orientation of G by Proposition 3.1. Conversely, if D
32 is an antistrong orientation of G , then $B(D)$ is a good 2-detachment of G . \diamond

34 We can now use Theorem 6.8 and the subsequent remark to deduce the following.

Theorem 7.2 *A graph $G = (V, E)$ has a good 2-detachment if and only if*

$$e(\mathcal{Q}) \geq |\mathcal{Q}| - 1 + b(\mathcal{Q}) \tag{7}$$

35 *for all partitions \mathcal{Q} of V . Furthermore, there exists a polynomial time algorithm which returns such a*
36 *2-detachment when it exists and otherwise returns a certificate, in terms of a partition violating (7),*
37 *that no such detachment exists.*

38 8 Non-separating antistrong spanning subdigraphs

39 While there are polynomial time algorithms for checking the existence of two edge-disjoint spanning
40 trees [5], or two arc-disjoint branchings (spanning out-trees) in a digraph (see e.g. [1, Corollary 9.3.2]),
41 checking whether we can delete a strong spanning subdigraph and still have a connected digraph is
42 difficult. Let $UG(D)$ denote the underlying undirected graph of a digraph D .

1 **Theorem 8.1** [3] *It is NP-complete to decide whether a given digraph D contains a spanning strong*
 2 *subdigraph H such that $UG(D - A(H))$ is connected.*

3 If we replace “strong” by “antistrong” above, the problem becomes solvable in polynomial time.

4 **Theorem 8.2** *We can decide in polynomial time for a given digraph $D = (V, A)$ on n vertices whether*
 5 *D contains a spanning antistrong subdigraph $H = (V, A')$ such that $UG(D - A')$ is connected.*

6 **Proof:** We may assume that D is antistrong, since this can be checked in linear time by verifying
 7 that $B(D)$ is connected. Let $M_1 = (A, \mathcal{I})$ be the cycle matroid of the underlying graph $UG(D)$ of
 8 D and let $M_2 = M(D) = (A, \mathcal{I}(D))$ be the matroid from Section 5 whose bases are the antistrong sets
 9 consisting of $2n - 1$ arcs. Let $M = M_1 \vee M_2$ be the union of the matroids M_1, M_2 , that is, a set X of
 10 arcs is independent in M if and only we can partition X into X_1, X_2 such that X_i is independent in
 11 M_i . For each of the matroids M_1, M_2 we can check in polynomial time whether a given subset X of
 12 arcs is independent in M_1 and M_2 (for M_1 we need to check that there is no cycle in $UG(D)[X]$ and
 13 for M_2 we need to check that there is no cycle in the subgraph of $B(D)[E_X]$ induced by the edges E_X
 14 corresponding to X in $B(D)$). Thus it follows from Edmonds’ algorithm for matroid partitioning [5]
 15 that we can find a base of M in polynomial time using the independence oracles of M_1, M_2 . The
 16 desired digraph H exists if and only if the size of a base in M is $(2n - 1) + (n - 1) = 3n - 2$. \diamond

17
 18 A similar proof gives the following.

19 **Theorem 8.3** *We can decide in polynomial time whether a digraph D contains $k + \ell$ arc-disjoint*
 20 *spanning subdigraphs $D_1, \dots, D_{k+\ell}$ such that D_1, \dots, D_k are antistrong and $UG(D_{k+1}), \dots, UG(D_{k+\ell})$*
 21 *are connected.*

22 9 Remarks and open problems

23 We saw in Theorem 3.6 that it is NP-complete to decide whether a given digraph contains two arc-
 24 disjoint spanning strong subdigraphs. We would be interested to know what happens if we modify
 25 the problem as follows.

26 **Question 9.1** *Can we decide in polynomial time whether D contains arc-disjoint spanning subdi-*
 27 *graphs D_1, D_2 such that D_1 is antistrong and D_2 is strongly connected?*

28 Inspired by Theorem 8.2 it is natural to ask the following intermediate question.

29 **Question 9.2** *Can we decide in polynomial time whether D contains arc-disjoint spanning subdi-*
 30 *graphs D_1, D_2 such that D_1 is antistrong and $UG(D_2)$ is 2-edge-connected?*

31 The following conjecture was raised in [2].

32 **Conjecture 9.3** [2] *There exists a natural number k such that every k -arc-strong digraph has arc-*
 33 *disjoint strong spanning subdigraphs D_1, D_2 .*

34 Perhaps the following special case may be easier to study.

35 **Conjecture 9.4** *There exists a natural number k such that every digraph D which is both k -arc-strong*
 36 *and k -arc-antistrong has arc-disjoint strong spanning subdigraphs D_1, D_2 .*

*

37 **Acknowledgement.** Bang-Jensen and Jackson wish to thank Jan van den Heuvel for stimulating
 38 discussions about antistrong connectivity. They also thank the Mittag-Leffler Institute for providing
 39 an excellent working environment.

References

- [1] J. Bang-Jensen and G. Gutin, Digraphs: Theory, Algorithms and Applications, 2nd edition, Springer Verlag, London 2009.
- [2] J. Bang-Jensen and A. Yeo, Decomposing k -arc-strong tournaments into strong spanning subdigraphs, *Combinatorica* **24** (2004), 331–349.
- [3] J. Bang-Jensen and A. Yeo, Arc-disjoint spanning sub(di)graphs in digraphs, *Theoret. Comp. Sci.* **438** (2012), 48–54.
- [4] G. Chartrand, H. Gavlas, M. Schultz, and C. E. Wall, Anticonnected digraphs, *Utilitas Math.* **51** (1997), 41–54.
- [5] J. Edmonds, Minimum partition of a matroid into independent sets, *J. Res. Nat. Bur. Standards, Sec. B.* **69** (1965), 67–72.
- [6] J. Edmonds, Submodular functions, matroids, and certain polyhedra, in: *Combinatorial Structures and their Applications*, eds. R. Guy, H. Hanani, N. Sauer, and J. Schönheim, Gordon and Breach, New York 1970, 69–87.
- [7] A. Frank, *Connections in combinatorial optimization*, Oxford Lecture series in mathematics and its applications **38** (2011).
- [8] H. N. Gabow, S. N. Maheshwari, and L. J. Osterweil, On two problems in the generation of program test paths, *IEEE Trans. Software Eng. SE* **2** (1976), 227–231.
- [9] N. Katoh and S. Tanigawa, A rooted-forest partition with uniform vertex demand, *J. Comb. Optim.* **24** (2012), 67–98.
- [10] C. St. J. A. Nash-Williams, Connected detachments of graphs and generalized Euler trails, *J. London Math. Soc.* **31** (2) (1985), 17–29.
- [11] S. Tanigawa, Matroids of gain graphs in applied discrete geometry, *Trans. Amer. Math. Soc.*, to appear, see arXiv:1207.3601.
- [12] W. Whiteley, Some matroids from discrete geometry, in: *Matroid Theory*, eds. J.E Bonin et al. *Contemporary Mathematics* **197**, Amer. Math. Soc., Providence (1996), 171–311.

1 A Appendix: a graph theoretical proof of Lemma 6.6

2 In this appendix, we give a graph theoretical proof of Lemma 6.6, recalled below.

3 **Lemma 6.6** *Let $G = (V, E)$ be a graph. Then G satisfies*

$$|E(H)| \leq 2|V(H)| - 1 \text{ for all nonempty subgraphs } H \text{ of } G, \text{ and} \quad (2)$$

$$|E(H)| \leq 2|V(H)| - 2 \text{ for all nonempty bipartite subgraphs } H \text{ of } G \quad (3)$$

4 *if and only if E can be partitioned into a forest and an odd pseudoforest.*

5
6 **Proof:** Recall that a **pseudoforest** is a graph in which each connected component contains at
7 most one cycle, and it is called **odd** if it does not contain even cycles. A theorem due to Whiteley [12]
8 (see also [7] p.367 for a short proof based on Edmonds' branching theorem) asserts that a graph
9 satisfies condition (2) if and only if its edge set can be partitioned into a forest and a pseudoforest.
10 So let us denote by a **2-decomposition** of a graph $G = (V, E)$ a pair (G_b, G_r) of spanning subgraphs
11 $G_b = (V, E_b)$ and $G_r = (V, E_r)$ such that $\{E_b, E_r\}$ is a partition of E and G_b is a forest of G and G_r
12 is a pseudoforest of G . We will call any sub-structure — edge, component, subgraph etc. — of G_r or
13 of G_b **red** or **black**, respectively, and for a subgraph H of G we denote by H_r and H_b the subgraph
14 of H induced by its red or black edges, respectively.

15 Without loss of generality we may assume that G is connected, and that G_b is a spanning tree of G
16 (otherwise we could move edges from G_r to G_b to make G_b connected). The **canonical bipartition** of
17 a 2-decomposition (G_b, G_r) of a connected graph G is the unique bipartition given by any 2-colouring
18 of G_b . Moreover, an edge of G_r which does not cross this bipartition is called (as previously) a
19 **precious edge** in (G_b, G_r) . A 2-decomposition of a 2-decomposable graph is **nice** if every red cycle
20 contains at least one precious edge.

21 First we establish the next claim.

22 **Claim 1** *A connected graph which has a 2-decomposition admits a nice 2-decomposition if and only*
23 *if (3) holds.*

24 **Proof:** First observe that for any subgraph H of G with at least one black edge, we have $|E(H)| =$
25 $|E(H_b)| + |E(H_r)| \leq (|V(H_b)| - 1) + |V(H_r)| = 2|V(H)| - 1$. For any red subgraph H with at least
26 one edge, we get $|E(H)| = |E(H_r)| \leq |V(H_r)| \leq 2|V(H)| - 2$. In particular (2) holds for every
27 2-decomposable graph.

28 The necessity is quite clear. Indeed, consider a nice 2-decomposition (G_b, G_r) of G and assume
29 that (3) does not hold. Thus there exists H a bipartite subgraph of G with $|E(H)| = 2|V(H)| - 1$. So
30 equality holds in the previous computation and we have $|E(H_b)| = |V(H_b)| - 1$ and $|E(H_r)| = |V(H_r)|$.
31 In particular H_b is a spanning tree of H and H_r contains at least one cycle C . As (G_b, G_r) is nice,
32 C contains a precious edge xy . As H_b is connected, there exists a black path P from x to y and P
33 has even length because x and y belong to the same part of the canonical bipartition of (G_b, G_r) . So
34 $P \cup xy$ forms an odd cycle of H , a contradiction.

35 Now let us prove the sufficiency. Let (G_b, G_r) be a 2-decomposition of G . A red component R
36 of the decomposition is **bad** if it is not a tree and its (hence unique) cycle does not contain any
37 precious edges. If we remove from a bad component R all its precious edges, we obtain several
38 connected components, one of which contains the cycle of R . We call this component the **core** of R
39 and denote it by $c(R)$. For convenience we use $c(R)$ below to denote both a vertex set and the red
40 subgraph induced by these vertices. Note that $G[c(R)]$ is bipartite as $c(R)$ contains no precious edge.
41 A **sequence** of the decomposition (G_b, G_r) is a list $\mathcal{R} = (c(R_1), \dots, c(R_i))$ of the cores of its bad red
42 components in decreasing order of cardinality.

43 Among all the 2-decompositions of G , we choose one whose sequences $\mathcal{R} = (c(R_1), \dots, c(R_i))$
44 satisfy

- 45 (a) i is minimal, and
- 46 (b) subject to (a), $|c(R_i)|$ is minimal.

47 We will prove that this 2-decomposition (G_b, G_r) is nice, that is, $\mathcal{R} = \emptyset$. Assume that it is not the
48 case and consider $\{X_1, \dots, X_p\}$ the black components of $G[c(R_i)]$ (that is, the connected components

1 of $G_b[c(R_i)]$). If $p = 1$, then $G[c(R_i)]$ is connected in black, and as $G_r[c(R_i)]$ is unicyclic, the bipartite
2 graph $G[c(R_i)]$ violates (3), a contradiction. So we must have $p \geq 2$. Now denote by W_1, \dots, W_q
3 the connected components of $G_b \setminus c(R_i)$ and construct a graph T' on $\{X_1, \dots, X_p, W_1, \dots, W_q\}$ by
4 connecting two vertices of T' if there exists an edge in G_b between the corresponding components. In
5 other words, we contract the (connected) vertex sets $X_1, \dots, X_p, W_1, \dots, W_q$ in G_b to single vertices.
6 So T' is a tree. Finally we consider T the minimal subtree of T' containing $\{X_1, \dots, X_p\}$. By definition
7 the leaves of T are in $\{X_1, \dots, X_p\}$ and as $p \geq 2$, T has at least two such leaves. So we consider a leaf
8 X_k of T which does not contain entirely the red cycle of R_i (this could occur even without violating
9 (3) if X_k is not connected in red for instance). We denote by $W_{k'}$ the only neighbour of X_k in T .
10 Now, we specify two edges, one black and one red in order to ‘change their color’ and obtain a con-
11 tradiction. First denote by uv the unique black edge between X_k and $W_{k'}$. We suppose that $u \in X_k$
12 and $v \in W_{k'}$ (so $v \notin c(R_i)$). Now we look at a 1-orientation of $c(R_i)$ (this is an orientation of $c(R_i)$)
13 in which every vertex has out-degree at most 1) and consider a maximal oriented red path leaving u
14 with all its vertices in X_k . As X_k does not contain entirely the red cycle of R_i , this path ends at a ver-
15 tex $x \in X_k$ which has a unique red out-neighbour $y \in X_{k''}$ for some $k'' \neq k$. We select this red edge xy .

16
17 Notice that the unique black path from x to y contains the edge uv . Indeed the unique path from
18 X_k to $X_{k''}$ in T corresponds to the unique black path P from X_k to $X_{k''}$ in G . As X_k is a leaf of
19 T , the first edge of P is uv and as X_k and $X_{k''}$ are respectively connected in black, the unique black
20 path from x to y contains P and so it contains the edge uv . This implies that $(G_b + xy) - uv$ is also
21 a spanning tree of G . Moreover, its bipartition is the same as the bipartition of G_b . Indeed as xy is
22 an edge lying inside the core of R_i , it is not precious and $G_b + xy$ is still bipartite and has the same
23 bipartition as G_b . Removing uv does not affect the bipartition (because $(G_b + xy) - uv$ is connected).

24 To conclude, we focus on the red part of the new 2-decomposition $((G_b + xy) - uv, (G_r + uv) - xy)$.
25 By construction, the component X of $G_r - xy$ containing u (and also x) is a red tree. Remark that
26 $X = R_i$ if and only if xy is an edge of the cycle of R_i . If v does not belong to R_i , then by adding
27 uv we attached X to another red component in $(G_r + uv) - xy$. As X contains at least one vertex,
28 namely u , $|c(R_i)|$ has decreased, a contradiction to (b) in the choice of (G_b, G_r) (or to (a) if $X = R_i$).
29 If v belongs to R_i but v does not belong to X (in this case we have $X \neq R_i$), then $v \in R_i \setminus c(R_i)$
30 and by adding uv we attached X to a vertex of $R_i \setminus c(R_i)$. Once again, $|c(R_i)|$ has decreased, a
31 contradiction to (b) in the choice of (G_b, G_r) . Finally if v belongs to X then adding uv produces a
32 new red unicyclic component. However as the red path in G_r from v to u starts in $R_i \setminus c(R_i)$ and
33 ends in $c(R_i)$, it contains a precious edge. So that newly created red unicyclic component is not bad,
34 and $|c(R_i)|$ has decreased. Hence, again, we either contradict (b), or (a) if $X = R_i$. contradicting (a). \diamond

35
36 Now to finish the proof, we will show how to go from a nice 2-decomposition of a connected graph
37 to a decomposition into a spanning tree and an odd pseudoforest (i.e. a pseudoforest in which every
38 cycle has odd length). Let $G = (V, E)$ be a connected graph which admits a nice 2-decomposition and
39 consider a nice 2-decomposition (G_b, G_r) of G with a minimum number of even red cycles. We will
40 show by contradiction that this decomposition has no even red cycle. Assume it is not the case and
41 denote by C_1, \dots, C_l the even red cycles of G_r . In each of these, select a precious edge $e_i = x_i y_i$ and
42 let $X = \{x_1, y_1, x_2, y_2, \dots, x_l, y_l\}$. Exchanging two edges between G_b and G_r will modify the structure
43 of G_b , and some previously selected precious edges could become not precious any more. To avoid
44 this we will find a vertex u with the following property

45
46 \mathcal{P} : There exists a component B of $G_b \setminus u$ such that one of the following hold:

- 47 • $B \cap X$ contains only one element and this is not in the same component of G_r as u (Case A).
- 48 • $(B \cup \{u\}) \cap X$ contains exactly two elements and they are the end vertices of some e_i (Case B).

49 First assume we have found such a vertex u and let us see how to conclude, depending of which
50 the two cases A or B we are in.

51
52 *Case A.* Denote by x_i the only element of $B \cap X$ and by B_i the red component of G_r containing x_i .
53 As u does not belong to B_i , we can find an edge vw along the black path in G_b from u to x_i such
54 that w and x_i are in the same component of $G_b - vw$, w belongs to B_i and v does not belong to B_i .

1 So we exchange the colors of vw and x_iy_i . The graph $B_i - x_iy_i$ is a tree and when we add vw to G_r
2 we connect this tree to another component of G_r . The component of $G_b - vw$ containing v is a tree
3 containing all the vertices of X except x_i . So the precious edges e_j with $j \neq i$ are still precious edges,
4 and this is also the case in $(G_b - vw) + x_iy_i$ which is a spanning tree of G . So, we reduce the number
5 of even red cycle of the nice 2-decomposition (G_b, G_r) , a contradiction.

6
7 *Case B.* Denote by x_i and y_i the two elements of $(B \cup \{u\}) \cap X$ and also by B_i the red component
8 of G_r containing the precious edge x_iy_i . If the black path P in G_b between x_i and y_i is not totally
9 contained in B_i then we can select a vertex u' not belonging to B_i along this path and end up in the
10 previous case with u replaced by u' . So P is totally contained in B_i . Then, as $P + x_iy_i$ is an odd cycle
11 (because x_iy_i is precious), we can find along P two consecutive vertices vw which are in the same part
12 of the bipartition induced by the bipartite graph $G_r[B_i]$. So we exchange x_iy_i and vw . As previously
13 $G_b + x_iy_i - vw$ is a spanning tree of G such that all the edges e_j with $j \neq i$ are still precious and vw
14 is also precious. The graph $G_r - x_iy_i + vw$ is now a pseudoforest, and we have reduced the number
15 of even red cycles of the nice 2-decomposition (G_b, G_r) , a contradiction.

16 Finally, let us see how to find a vertex u in G which has property \mathcal{P} . Consider T' the minimal
17 subtree of G_b containing all the vertices of the set X . In particular all the leaves of T' are elements
18 of X . Then build the tree T from T' by replacing iteratively each vertex of degree 2 in T' and not
19 belonging to X by an edge linking its two neighbours in T' . The vertices of T are now vertices of X
20 or have degree at least three in T . Assume first that a leaf f of T has its neighbour f' in X . Denote
21 by B the component of $G_b \setminus f'$ containing f . By construction f is the unique element of $B \cap X$. We
22 select $u = f'$. If f and f' are in different components of G_r then we are in *Case A*, otherwise we are
23 in *Case B*.

24 So we can assume that all the leaves of T are neighbours of vertices of T which are not in X and have
25 degree at least three in T . Consider now a leaf f' of the tree obtained from T by removing its leaves.
26 Denote by L the set of neighbours of f' in T which are leaves of T . If $|L| = 2$ and L consists of the
27 end vertices of some e_i then we choose $u = f'$ and are in *Case B*. Otherwise, let B_i be the component
28 of G_r containing f' and consider a vertex f of L not belonging to B_i (this exists as $|B_i \cap X| = 2$).
29 Then we choose $u = f'$ and B to be the component of $G_b \setminus f'$ containing f and we are in *Case A*. \diamond
30