



**HAL**  
open science

## Aperiodic Tilings and Entropy

Bruno Durand, Guilhem Gamard, Anaël Grandjean

► **To cite this version:**

Bruno Durand, Guilhem Gamard, Anaël Grandjean. Aperiodic Tilings and Entropy. DLT: Developments in Language Theory, Aug 2014, Ekaterinburg, Russia. pp.166-177, 10.1007/978-3-319-09698-8\_15 . lirmm-01480693

**HAL Id: lirmm-01480693**

**<https://hal-lirmm.ccsd.cnrs.fr/lirmm-01480693v1>**

Submitted on 1 Mar 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Aperiodic tilings and entropy

Bruno DURAND, Guilhem GAMARD, Anaël GRANDJEAN

August 27, 2014

- 1 Introduction
- 2 J. Kari & K. Culik's tileset
- 3 Aperiodicity
- 4 Positive entropy
- 5 Conclusion

# Wang tiles


## Wang tiles

- Finite set of colors
- Alphabet = colored squares
- Adjacent borders have matching colors

$$c = \{\text{red, yellow, blue}\}$$

$$\Sigma = \left\{ \begin{array}{c} \text{blue} \\ \text{red} \end{array} \text{ square}, \begin{array}{c} \text{red} \\ \text{blue} \end{array} \text{ square}, \text{yellow square} \right\}$$

$$\Sigma' = \left\{ \begin{array}{c} \text{blue} \\ \text{red} \end{array} \text{ square} \right\}$$



# Aperiodic tilesets

## Definition

A set of tiles is **aperiodic** when:

- it can cover the plane;
- it cannot cover the plane periodically.

Cover such that adjacent borders have matching colors

# The history of small aperiodic tilesets

## Self-similar

1964	R. Berger	> 20,000 tiles
1966	D. Knuth	96 tiles
1971	R. Robinson	52 tiles
1974	R. Penrose	32 tiles
1986	R. Ammann, B. Grünbaum, G. Shephard	16 tiles

## Not self-similar

1996	J. Kari	14 tiles
1996	K. Culik	13 tiles

# Our result

## Theorem

*The Kari-Culik tileset has positive entropy.*

Intuitively:

- Description of a  $n \times n$ -square takes  $\Omega(n^2)$  bits
- It contains dense “random” bits

Note: entropy zero was conjectured.

1 Introduction

2 J. Kari & K. Culik's tileset

3 Aperiodicity

4 Positive entropy

5 Conclusion




## A function with aperiodic orbits

Consider this function:



$$f: \left[\frac{1}{3}; 2\right] \rightarrow \left[\frac{1}{3}; 2\right]$$
$$x \mapsto \begin{cases} 2x & \text{if } x \leq 1 \\ x/3 & \text{if } x \geq 1 \end{cases}$$

- Its orbits, i.e. sequences  $u_x = (f^{on}(x))_{n \in \mathbb{N}}$ , are aperiodic
- Its orbits are also dense in  $\left[\frac{1}{3}; 2\right]$

# The general idea





## Multiplications done by transducers





- $(2, 0)$  stands for “read 2, write 0”

## Multiplications done by transducers




- $(2, 0)$  stands for “read 2, write 0”

## From transducers to tile sets




- States of  $M_{1/3}$ ,  $M_2$ : disjoint colors
- One line = one run


# An aperiodic set of tiles



# An aperiodic set of tiles



# An aperiodic set of tiles





- 1 Introduction
- 2 J. Kari & K. Culik's tileset
- 3 Aperiodicity**
- 4 Positive entropy
- 5 Conclusion

## Lines and averages

### Definition

The *average* of a sequence  $(u_n)$  is:

$$\text{avg}(u) = \lim_{n \rightarrow \infty} \frac{1}{2n+1} \sum_{k=-n}^n u_k$$

Tilings: average of a line = average of northern sides

# Aperiodicity

Theorem (J. Kari and K. Culik, 1996)

*The Kari-Culik tileset is aperiodic.*

## Sketch of proof.

- Suppose there is a periodic tiling. Then each line has an average. The averages are periodic: contradiction.
- To tile the plane, start from  $\dots 11111111 \dots$  and run the transducers forever.





- 1 Introduction
- 2 J. Kari & K. Culik's tileset
- 3 Aperiodicity
- 4 Positive entropy**
- 5 Conclusion


# Each line has an average

## Lemma

*In any tiling, each line has an average.*

## Sketch of proof.

Consider a line without an average.



Density  $\implies \exists n$  s.t.  $f^{on}(x) < 1 < f^{on}(y)$



# Entropy

$C(n)$  = the number of  $n \times n$ -squares found in any tiling

## Definition





We call the **entropy** the following quantity:

$$E = \lim_{n \rightarrow \infty} \frac{\log C(n)}{n^2}$$

- Classical definition in dynamical systems
- With 13 tiles, if  $C(n) \sim 13^{\epsilon n^2}$ , then  $E = \epsilon$

## Substitutive pairs

are pairs of distinct patterns with the same borders.









# Substitutive pairs generate entropy

## Lemma

*If a substitutive square is found in any  $n \times n$ -square of any tiling, then the entropy of the tiles is positive.*



 = substitutive square




# Substitutive pairs appear often (1/2)

## Lemma

*Whenever a pattern  $0111^\alpha 0$  occurs on a line of tiles, there is a substitutive square intersecting this pattern.*

## Sketch of proof.

Case analysis. □



Middle case


## Substitutive pairs appear often (1/2)

### Lemma

*Whenever a pattern  $0111^\alpha 0$  occurs on a line of tiles, there is a substitutive square intersecting this pattern.*

### Sketch of proof.

Case analysis. □



Leftmost case


# Substitutive pairs appear often (1/2)

## Lemma

*Whenever a pattern  $0111^\alpha 0$  occurs on a line of tiles, there is a substitutive square intersecting this pattern.*

## Sketch of proof.

Case analysis. □



Rightmost case

## Substitutive pairs appear often (2/2)

### Lemma

*In any line with an average  $\in ]\frac{3}{4}; \frac{9}{10}[$ , a pattern of the form  $0111^\alpha 0$  appears regularly.*

### Sketch of proof.

If there are no “0” regularly, then the average is 1.

If there are no “111” regularly, then the average is  $\leq \frac{3}{4}$ . □

- All orbits regularly meet the interval  $] \frac{3}{4}; \frac{9}{10}[$
- Hence substitutive squares appear often enough

- 1 Introduction
- 2 J. Kari & K. Culik's tileset
- 3 Aperiodicity
- 4 Positive entropy
- 5 Conclusion**

# Many thanks for your attention!

## Our result

- The entropy of the Kari-Culik tileset is positive
- The Kari-Culik-tilings are not all self-similar

## Open problems

- Characterize the language of words which can appear on K.C.'s lines?
- Is there a tileset working the same way, but with 0 entropy?
- Is there a sub-shift of finite type  $A$ , with positive entropy, such that any subshift of finite type  $\subset A$  also has positive entropy?