



HAL
open science

Structure and Enumeration of K_4 -minor-free links and link diagrams

Juanjo Rué, Dimitrios M. Thilikos, Vasiliki Velona

► **To cite this version:**

Juanjo Rué, Dimitrios M. Thilikos, Vasiliki Velona. Structure and Enumeration of K_4 -minor-free links and link diagrams. *Electronic Notes in Discrete Mathematics*, 2018, 68, pp.119-124. 10.1016/j.endm.2018.06.021 . lirmm-01890505

HAL Id: lirmm-01890505

<https://hal-lirmm.ccsd.cnrs.fr/lirmm-01890505>

Submitted on 8 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Structure and Enumeration of K_4 -minor-free links and link diagrams

Juanjo Rué^{1,2}

Department of Mathematics, Universitat Politècnica de Catalunya, Barcelona, Spain

Dimitrios M. Thilikos^{1,2}

ALGCo project-team, LIRMM, Université de Montpellier, CNRS, Montpellier, France
Department of Mathematics, National and Kapodistrian University of Athens, Athens, Greece

Vasiliki Velona^{1,2}

Department of Economics, Universitat Pompeu Fabra, Barcelona, Spain
Department of Mathematics, Universitat Politècnica de Catalunya, Barcelona, Spain

Abstract

We study the class \mathcal{L} of link types that admit a K_4 -minor-free diagram, i.e., they can be projected on the plane so that the resulting graph does not contain any subdivision of K_4 . We prove that \mathcal{L} is the closure of a subclass of torus links under the operation of connected sum. Using this structural result, we enumerate \mathcal{L} and subclasses of it, with respect to the minimal number of crossings or edges in a projection of $L \in \mathcal{L}$. Further, we enumerate (both exactly and asymptotically) all connected K_4 -minor-free link diagrams, all minimal connected K_4 -minor-free link diagrams, and all K_4 -minor-free diagrams of the unknot.

Keywords: series-parallel graphs, links, knots, generating functions, asymptotic enumeration, map enumeration

1 Introduction

The exhaustive generation of knots and links is a well-established problem in low dimensional geometry (see [7, Ch.5], for instance). However, there are very few enumerative results in the literature and they are relatively recent, (see [10] and [6]). Moreover, there seems to be no known results connecting graph theoretic classes with link classes.

Let us start with some formal definitions. A *knot* K is a smooth embedding of the 1-dimensional sphere \mathbb{S}^1 in \mathbb{R}^3 . A *link* is a finite disjoint union of knots $L = K_1 \cup \dots \cup K_n$. Two links L_1 and L_2 are equivalent if there is a continuous and injective function $h : \mathbb{R}^3 \times [0, 1] \rightarrow \mathbb{R}^3$, such that $h(L_1, 0) = L_1$ and $h(L_1, 1) = L_2$. A link equivalent to a set of non intertwined circles is called a *trivial link*. If it is a knot, we also call it the *unknot*.

Consider a link L and a sphere \mathbb{S}^2 embedded in such a way that it meets the link transversely in exactly two points P_1 and P_2 . Then we can discern two different links L_1, L_2 , after connecting P_1 and P_2 . The first corresponds to the part of L in the interior of the sphere and the second to the part in the exterior. We then call L a *connected sum* with *factors* L_1, L_2 , denoted $L_1 \# L_2$. A link that does not have non-trivial factors is called *prime*, otherwise *composite*. A link is *split* if there is a sphere embedded in the link complement that separates the link. Each of the components is called a *disjoint component* of the link and, conversely, the link is their *disjoint sum*.

Let $L \in \mathbb{R}^3$ be a link and let $\pi : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a projection map. If for all $x \in L$, $|\pi^{-1}(x)| = 2$ or 1 , and all the double points are finite and transverse, then the projection is said to be *regular*. For all knots defined here, there exists a regular projection [2, Ch. 3]. This allows to work with *link-diagrams*, i.e. a triple (V, E, σ) , where (V, E) is a 4-regular plane graph and $\sigma : V(G) \rightarrow \binom{E(G)}{2}$, where, for every $v \in V(G)$, $\sigma(v)$ is a set of two opposite edges of the embedding of G and encodes which pair is overcrossing. Notice that a link-diagram may also have vertexless edges. We call these edges *trivial*. The *crossing number* of a link L is the minimum number of crossings that can have a diagram of it. Such a diagram is called *minimal*. We also say that a diagram (V, E, σ) is K_4 -minor-free if the graph (V, E) does not contain

¹The first and the third author have partially been supported by the MINECO projects MTM2014-54745-P, MTM2017-82166-P and the María de Maetzu research grant MDM-2014-0445 (BGSMath). The second author has been supported by the project “ESIGMA” (ANR-17-CE40-0028). The third author has been supported under an FPI grant from the MINECO research project MTM2015-67304-PI.

²Emails: juan.jose.rue@upc.edu, sedthilk@thilikos.info, vasiliki.velona@upf.edu

any subgraph that is a subdivision of a K_4 .

A *torus link* is a link that can be embedded on the torus. They are denoted by $T(p, q)$, $p, q \in \mathbb{Z}$, where p and q are the number of times that the link crosses the meridian and the longitude cycle, respectively. Let \mathcal{T}_2 be the closure of torus links of type $T(2, q)$, $q \in \mathbb{Z} - \{0\}$, under the connected sum operation. Let \mathcal{L} be the class of links that have a K_4 -minor-free link-diagram and $\mathbf{mcl}(\mathcal{T}_2)$ be the closure of \mathcal{T}_2 under disjoint sum.

2 Structure of K_4 -links and enumeration

Our first result gives the type of links that admit a K_4 -minor-free diagram.

Theorem 2.1 *The links that admit a K_4 -minor-free diagram are exactly the ones in $\mathbf{mcl}(\mathcal{T}_2)$, i.e. $\mathcal{L} = \mathbf{mcl}(\mathcal{T}_2)$.*

Based on the above, we derive a precise description of links that admit a K_4 -minor-free diagram. Let $\bar{\mathcal{L}}$ be the class of non-split links in \mathcal{L} and $\hat{\mathcal{L}}$ the class of links in \mathcal{L} with no trivial disjoint components, with size defined as their crossing number. We will obtain the asymptotic growth of \mathcal{L} with respect to the number of edges in a minimal diagram (not the crossing number, so as to account also for trivial disjoint components). Let $L(z)$, $\bar{L}(z)$, and $\hat{L}(z)$ be the respective generating functions. Finally, let \mathcal{T} be the class of non plane, unrooted trees, where the vertices are labelled with multisets of the set $\{1\} \cup \{\pm 3, \pm 5, \dots\}$ and the edges are labelled with an element of the set $\{2, \pm 4, \pm 6, \dots\}$. These labels will encode crossing numbers, hence a label i corresponds to size $|i|$. The size of $T \in \mathcal{T}$ is the sum of its labels. See Figure 1 for an example of such a tree.




Figure 1. An object of \mathcal{T} .

Proposition 2.2 $\bar{\mathcal{L}} \cong \mathcal{T}$ and $\text{MSet}_{\geq 1}(\mathcal{T} \setminus T_1) \cong \hat{\mathcal{L}}$, where MSet is the multiset operator with at least one element and $T_1 \in \mathcal{T}$ is the tree with one vertex and label 1.

Using Proposition 2.2 and the framework of the *Symbolic Method* (see [4]), we derive combinatorial expressions for \mathcal{L} that translate to generating functions. Let $\bar{\mathcal{K}}$ and \mathcal{K} be the classes of prime knots and composite knots plus the trivial knot in \mathcal{L} , respectively. For prime knots in \mathcal{L} , i.e. prime torus knots $T(2, 2i + 1)$, $i \in \mathbb{Z} \setminus \{0, -1\}$, it holds that $\bar{K}(z) = 2 \sum_{i \geq 1} z^{2i+1} = \frac{2z^3}{1-z^2}$, counting according to their crossing number. For \mathcal{K} , it is enough to consider non-empty multisets of prime torus knots, therefore $\mathcal{K} = \text{MSet}_{\geq 1}(\bar{\mathcal{K}})$, which translates to the expression

$$K(z) = \exp \left(\sum_{k \geq 1} \frac{1}{k} \bar{K}(z^k) \right) - 1 = \exp \left(\sum_{k \geq 1} \frac{1}{k} \frac{2z^{3k}}{1-z^{2k}} \right) - 1.$$

Let \mathcal{G} be the class of unrooted and unlabelled non-plane trees. Notice that one cannot replace immediately z for \mathcal{K} in the ogf $G(z)$, since the vertices are not distinguishable. Hence, to continue, one needs to use cycle indices. Let \mathcal{G}^\bullet be the class of rooted and unlabelled non-plane trees. For the cycle index of \mathcal{G}^\bullet , it is known (see [1]) that

$$\mathcal{Z}_{\mathcal{G}^\bullet}(s_1, s_2, \dots) = s_1 \exp \left(\sum_{k \geq 1} \frac{1}{k} \mathcal{Z}_{\mathcal{G}^\bullet}(s_k, s_{2k}, \dots) \right).$$

Let \mathcal{E} be the combinatorial class of the edge labels of \mathcal{T} , hence $E(z) = z^2 + \frac{2z^4}{1-z^2}$, and let $f(z) = E(z)K(z)$. We can now obtain the ordinary generating function of $\mathcal{F} = \mathcal{T}^\bullet \circ (\mathcal{E} \times \mathcal{L}_c)$. By Polya's Theorem, the latter satisfies the equation $F(z) = \mathcal{Z}_{\mathcal{G}^\bullet}(f(z), f(z^2), \dots)$. Then, it holds that

$$\bar{L}(z) = \frac{F(z)}{E(z)} + \frac{E(z)}{2} \left(-\frac{F(z)^2}{E(z)^2} + \frac{F(z^2)}{E(z^2)} \right),$$

by the Dissymmetry Theorem (see [1]). The first terms of $\bar{L}(z)$ are as follows:

$$\bar{L} = 1 + z^2 + 2z^3 + 3z^4 + 4z^5 + 9z^6 + 12z^7 + 26z^8 + 40z^9 + \dots$$

We can then get asymptotic estimates by means of complex analytic tools:

Theorem 2.3 $\bar{\mathcal{L}}$ has asymptotic growth of the form:

$$[z^n] \bar{L}(z) \sim \frac{cn^{-3/2}}{\Gamma(-1/2)} \rho^{-n}, \quad \rho \approx 0.44074, \quad c \approx 1.45557.$$

Additionally, $\hat{\mathcal{L}}$ has the same type of asymptotic growth with the same ρ and $c \approx 3.61691$.

The generating function $L(z)$ is equal to $\hat{L}(z^2) \frac{1}{1-z}$, since a link diagram of n vertices has $2n$ non-trivial edges and a number of trivial edges. We thus obtain the following corollary.

Corollary 2.4 \mathcal{L} has asymptotic growth of the form:

$$[z^n]L(z) \sim \frac{cn^{-3/2}}{\Gamma(-1/2)}\rho^{-n},$$

where $\rho \approx 0.44074$ and $c \approx 8.97779$ or $c \approx 3.95687$, depending on whether n is even or odd, respectively.

Finally, we also get asymptotic estimates for the coefficients of $K(z)$. The following result is a consequence of Meinardus' Theorem (see [4, VIII.23]).

Theorem 2.5 \mathcal{K} has asymptotic growth of the form:

$$[z^n]K(z) \sim cn^{-7/4} \exp(c'n^{1/2}),$$

where $c = \frac{e^{2 \ln(2)\zeta(0)}(\Gamma(2)\zeta(2))^{5/4}}{2\sqrt{\pi}} \approx 0.26275$, $c' = 2\sqrt{\Gamma(2)\zeta(2)} \approx 2.56509$, and $\zeta(z)$ is the Riemann zeta function.

3 Enumeration of families of link diagrams

Enumerating connected K_4 -free link-diagrams is equivalent to enumerating 4-regular unrooted planar maps which are K_4 -minor-free, that we call $\bar{\mathcal{M}}_1$. We first give a construction for the rooted ones, \mathcal{M}_1 , with respect to edges, adapting the construction for rooted 4-regular maps in [8]. With this we obtain a functional system of equations that can be analyzed by using the analytic machinery developed by Drmota in [3]. Finally, by adapting an argument by Richmond and Wormald in [9], we are able to show that we can unroot the maps under study, showing that the maps in our class with non-trivial automorphisms are exponentially few. This study gives the following result:

Theorem 3.1 The class of connected K_4 -free link diagrams $\bar{\mathcal{M}}_1$ satisfies that

$$[z^n]\bar{\mathcal{M}}_1(z) \sim \frac{1}{2n} \frac{cn^{-3/2}}{\Gamma(-1/2)}\rho^{-n}2^n, \quad \rho \approx 0.31184, \quad c \approx 1.52265$$

Refining the combinatoric analysis in the previous case (with more auxiliary classes) and using again an adaptation of the argument of Richmond and Wormald in [9], we are able to asymptotically enumerate also the class of minimal link diagrams $\bar{\mathcal{M}}_2$, as well as the class of K_4 -free link diagrams of the unknot, $\bar{\mathcal{M}}_3$:

Theorem 3.2 The class of connected K_4 -free minimal link diagrams $\bar{\mathcal{M}}_2$, and

the class of K_4 -free link diagrams of the unknot, \bar{M}_3 satisfy:

$$[z^n]\bar{M}_2(z) \sim \frac{1}{2n} \frac{c_2 n^{-3/2}}{\Gamma(-1/2)} \rho_2^{-n}, \quad \rho_2 \approx 0.41456, \quad c_2 \approx 0.81415,$$

$$[z^n]\bar{M}_3(z) \sim \frac{1}{2n} \frac{c_3 n^{-3/2}}{\Gamma(-1/2)} \rho_3^{-n}, \quad \rho_3 \approx 0.23188, \quad c_3 \approx 2.19020.$$

Acknowledgement. All authors like to thank Lefteris Kirousis that contributed to the inception of this research.

References

- [1] F. Bergeron, G. Labelle, and P. Leroux. *Combinatorial species and tree-like structures*, volume 67. Cambridge University Press, 1998.
- [2] P. R. Cromwell. *Knots and links*. Cambridge University Press, 2004.
- [3] M. Drmota. *Systems of functional equations. Random Structures and Algorithms*, 10, 103-124, 1997.
- [4] P. Flajolet and R. Sedgewick. *Analytic combinatorics*. Cambridge University press, 2009.
- [5] A. Kawauchi. *A survey of knot theory*. Birkhäuser, 2012.
- [6] S. Kunz-Jacques and G. Schaeffer. The asymptotic number of prime alternating links. In *FPSAC'2001*, pages 10–p. Arizona State University, 2001.
- [7] W. Menasco and M. Thistlethwaite. *Handbook of knot theory*. Elsevier, 2005.
- [8] M. Noy, C. Requilé, and J. Rué. Enumeration of labeled 4-regular planar graphs. *Submitted*, available on-line at [arXiv: 1709.04678](https://arxiv.org/abs/1709.04678).
- [9] L. B. Richmond and N. C. Wormald. Almost all maps are asymmetric. *J. of Combinatorial Theory, Series B*, 63(1):1–7, 1995.
- [10] C. Sundberg and M. Thistlethwaite. The rate of growth of the number of prime alternating links and tangles. *Pacific J. of mathematics*, 182(2):329–358, 1998.