



HAL
open science

On the König-Egerváry theorem for k -paths

Stéphane Bessy, Pascal Ochem, Dieter Rautenbach

► **To cite this version:**

Stéphane Bessy, Pascal Ochem, Dieter Rautenbach. On the König-Egerváry theorem for k -paths. Journal of Graph Theory, 2019, 91 (1), pp.73-87. 10.1002/jgt.22421 . lirmm-02078255

HAL Id: lirmm-02078255

<https://hal-lirmm.ccsd.cnrs.fr/lirmm-02078255>

Submitted on 25 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the König-Egerváry Theorem for k -Paths

Stéphane Bessy¹, Pascal Ochem¹, and Dieter Rautenbach²

¹ Laboratoire d'Informatique, de Robotique et de Microélectronique de Montpellier, Montpellier, France, stephane.bessy@lirmm.fr, pascal.ochem@lirmm.fr

² Institute of Optimization and Operations Research, Ulm University, Ulm, Germany, dieter.rautenbach@uni-ulm.de

Abstract

The famous König-Egerváry theorem is equivalent to the statement that the matching number equals the vertex cover number for every induced subgraph of some graph if and only if that graph is bipartite. Inspired by this result, we consider the set \mathcal{G}_k of all graphs such that, for every induced subgraph, the maximum number of disjoint paths of order k equals the minimum order of a set of vertices intersecting all paths of order k . For $k \in \{3, 4\}$, we give complete structural descriptions of the graphs in \mathcal{G}_k . Furthermore, for odd k , we give a complete structural description of the graphs in \mathcal{G}_k that contain no cycle of order less than k . For these graph classes, our results yield efficient recognition algorithms as well as efficient algorithms that determine maximum sets of disjoint paths of order k and minimum sets of vertices intersecting all paths of order k .

Keywords: König-Egerváry theorem; matching; vertex cover; k -path vertex cover; bipartite graph

MSC2010: 05C69

1 Introduction

The famous Kőnig-Egerváry theorem [4,8] states that the matching number $\nu(G)$ of a bipartite graph G equals its vertex cover number $\tau(G)$. Since a graph is bipartite if and only if it contains no odd cycle C_{2k+1} as an induced subgraph, and $\nu(C_{2k+1}) = k < k + 1 = \tau(C_{2k+1})$, the Kőnig-Egerváry theorem is equivalent to the statement that $\nu(H) = \tau(H)$ for every induced subgraph H of some graph G if and only if G is bipartite. Considering a matching as a packing of paths of order 2, and a vertex cover as a set of vertices intersecting every path of order 2, it is natural to ask for generalizations of the Kőnig-Egerváry theorem for longer paths, and to consider the corresponding graph classes generalizing the bipartite graphs.

In the present paper we study such generalizations.

We consider finite, simple, and undirected graphs as well as finite and undirected multi-graphs that may contain loops and parallel edges. Let k be a positive integer, and let G be a graph. A k -path and a k -cycle in G is a not necessarily induced path and cycle of order k in G , respectively. A set of disjoint k -paths in G is a k -matching in G , and a set of vertices of G intersecting every k -path in G is a k -vertex cover in G . The k -matching number $\nu_k(G)$ of G is the maximum cardinality of a k -matching in G , and the k -vertex cover number $\tau_k(G)$ of G is the minimum cardinality of a k -vertex cover in G .

Clearly,

$$\nu_k(G) \leq \tau_k(G).$$

Let \mathcal{G}_k be the set of all graphs G such that $\nu_k(H) = \tau_k(H)$ for every induced subgraph H of G . As noted above, the Kőnig-Egerváry theorem is equivalent to the statement that \mathcal{G}_2 is the set of all bipartite graphs. Since $\nu_1(G) = \tau_1(G) = n(G)$ for every graph G of order $n(G)$, the set \mathcal{G}_1 contains all graphs.

For $k \in \{3, 4\}$, we give complete structural descriptions of the graphs in \mathcal{G}_k . Furthermore, for odd k , we give a complete structural description of the graphs in \mathcal{G}_k that contain no cycle of order less than k .

Among the two parameters $\nu_k(G)$ and $\tau_k(G)$, only the latter seems to have received considerable attention in the literature [2,3,9]. Note that a set X of vertices of a graph G is a 3-vertex cover if and only if its complement $V(G) \setminus X$ is a so-called *dissociation set* [1,13], that is, a set of vertices inducing a subgraph of maximum degree at most 1. Probably motivated by this connection, the 3-vertex cover number has been studied in detail [6,7,10–12]. For every k at least 3, the hardness of the k -vertex cover number has been shown in [3]. It follows from known results (cf. [GT12] in [5]) that, for every integer k at least 3, it is NP-complete to decide for a given graph G whose order $n(G)$ is a multiple of k , whether $\nu_k(G) = \frac{n(G)}{k}$, that is, whether G has a *perfect* k -matching.

For a positive integer k and a graph G , let \mathcal{P}_k be the set of all k -paths of G . The parameters $\nu_k(G)$ and $\tau_k(G)$ are the optimum values of the following integer linear programs.

$$\nu_k(G) \left\{ \begin{array}{ll} \max & \sum_{P \in \mathcal{P}_k} x_P \\ \text{s.t.} & \sum_{P \in \mathcal{P}_k: u \in V(P)} x_P \leq 1 \quad \forall u \in V(G) \\ & x_P \in \{0, 1\} \quad \forall P \in \mathcal{P}_k \end{array} \right.$$

$$\tau_k(G) \left\{ \begin{array}{ll} \min & \sum_{u \in V(G)} y_u \\ \text{s.t.} & \sum_{u \in V(P)} y_u \geq 1 \quad \forall P \in \mathcal{P}_k \\ & y_u \in \{0, 1\} \quad \forall u \in V(G) \end{array} \right.$$

Relaxing “ $\in \{0, 1\}$ ” in both programs to “ ≥ 0 ” yields a pair of dual linear programs, whose optimal values we denote by $\nu_k^*(G)$ and $\tau_k^*(G)$, respectively. Since $\nu_k(G) = \nu_k^*(G) = \tau_k^*(G) = \tau_k(G)$ for a given graph G in \mathcal{G}_k , linear programming allows to determine $\nu_k(G)$ and $\tau_k(G)$ for G in polynomial time. Furthermore, since \mathcal{G}_k is closed under taking induced subgraphs, iteratively considering the removal of individual vertices, one can use linear programming to determine in polynomial time an induced subgraph G' of G of minimum order with $\nu_k(G) = \nu_k(G') = \tau_k(G') = \tau_k(G)$. Note that a maximum k -matching in G' covers all vertices of G' , and is also a maximum k -matching in G , and that a minimum k -vertex cover in G' is also a minimum k -vertex cover in G . Now, within G' , one can use linear programming to iteratively identify in polynomial time k -paths as well as vertices whose removal reduces the k -matching number as well as k -vertex cover by exactly 1, respectively. Clearly, the identified k -paths form a maximum k -matching in G , and the identified vertices form a minimum k -vertex cover in G .

We discuss some generic examples of graphs in \mathcal{G}_k , namely,

- forests,
- k -subdivisions of multigraphs, and,
- $k/2$ -subdivisions of bipartite multigraphs for even k .

Trivially, every graph of order less than k belongs to \mathcal{G}_k , which implies that the local structure of the graphs in \mathcal{G}_k is not simple.

The fact that all forests belong to \mathcal{G}_k follows by an inductive argument using the following lemma. In fact, the lemma yields a simple polynomial time reduction algorithm that determines a maximum k -matching as well as a minimum k -vertex cover in a given forest. An efficient algorithm computing a minimum k -vertex cover in a given forest was presented in [3].

Lemma 1 *Let k be a positive integer. If the graph G is the union of a tree T and a graph G' such that T and G' share exactly one vertex x , the tree T contains a k -path, but the forest $T - x$ contains no k -path, then $\nu_k(G) = \nu_k(G' - x) + 1$ and $\tau_k(G) = \tau_k(G' - x) + 1$.*

Proof: Every k -path in T contains x . Hence, if \mathcal{P} is a k -matching in G , then at most one path in \mathcal{P} intersects $V(T)$. Removing any such path yields a k -matching in $G' - x$, which implies $\nu_k(G) \leq \nu_k(G' - x) + 1$. Conversely, if \mathcal{P}' is a k -matching in $G' - x$, then adding a k -path contained in T , yields a k -matching in G , which implies $\nu_k(G) \geq \nu_k(G' - x) + 1$.

If X is a k -vertex cover in G , then X intersects $V(T)$, and $X \setminus V(T)$ is a k -vertex cover in $G' - x$, which implies $\tau_k(G) \geq \tau_k(G' - x) + 1$. Conversely, adding x to any k -vertex cover in $G' - x$ yields a k -vertex cover in G , which implies $\tau_k(G) \leq \tau_k(G' - x) + 1$. \square

The following lemma captures some natural cycle conditions for the graphs in \mathcal{G}_k .

For an integer n , let $[n]$ be the set of positive integers at most n .

Lemma 2 *Let k and p be positive integers.*

- (i) *Every cycle of order at least k in every graph in \mathcal{G}_k has order 0 modulo k .*
- (ii) *A set X of vertices of the cycle $C_{pk} : u_1 u_2 \dots u_{pk} u_1$ of order pk is a minimum k -vertex cover in C_{pk} if and only if $X = \{u_{i+(j-1)k} : j \in [p]\}$ for some $i \in [k]$.*
- (iii) *If G is in \mathcal{G}_3 , C is a cycle in G , and u and v are distinct vertices of C that have neighbors outside of $V(C)$, then $\text{dist}_C(u, v) \equiv 0 \pmod{3}$.*
- (iv) *If G is in \mathcal{G}_4 , C is a cycle of length at least 4 in G , and u and v are distinct vertices of C that have neighbors outside of $V(C)$, then $\text{dist}_C(u, v) \equiv 0 \pmod{2}$.*

Proof: If the graph G arises by adding some edges to the cycle C_n of order n , where n is at least k , then $\nu_k(G) = \lfloor \frac{n}{k} \rfloor \leq \lceil \frac{n}{k} \rceil = \tau_k(C_n) \leq \tau_k(G)$, which implies (i). The value of $p = \tau_k(C_{pk})$ and the fact that every k -vertex cover in C_{pk} has to contain at least one of any k consecutive vertices of C_{pk} implies (ii).

If G , C , u , and v are as in (iii), u' is a neighbor of u outside of $V(C)$, v' is a neighbor of v outside of $V(C)$, and G' is the subgraph of G induced by $V(C) \cup \{u', v'\}$, then $\nu_3(G') = \lfloor \frac{n(C) + |\{u', v'\}|}{3} \rfloor = \frac{n(C)}{3}$. Since $G \in \mathcal{G}_3$, we obtain $\tau_3(G') = \frac{n(C)}{3} = \tau_3(C)$, which implies that every minimum 3-vertex cover in G' is a minimum 3-vertex cover in C , and, hence, as described in (ii). Since u and v must both belong to every minimum 3-vertex cover in G' , their distance on C must be a multiple of 3.

Now, if G , C , u , and v are as in (iv), and u' , v' , and G' are as above, then $\nu_4(G') = \lfloor \frac{n(C) + |\{u', v'\}|}{4} \rfloor = \frac{n(C)}{4}$. Again every minimum 4-vertex cover in G' is a minimum 4-vertex cover in C , and, hence, as described in (ii). Since every minimum 4-vertex cover in G' contains either u or both vertices at distance 2 from u within C , and the same holds for v , the distance of u and v on C must be even. \square

Lemma 2 (i) and (iii) suggest that subdividing every edge of a multigraph $k - 1$ times yields a natural candidate for a graph in \mathcal{G}_k . For a positive integer k , let the k -subdivision $Sub_k(H)$ of a multigraph H arise by subdividing every edge of H exactly $k - 1$ times, that is,

- every edge between distinct vertices u and v is replaced by a $(k + 1)$ -path between u and v whose internal vertices have degree 2, and
- every loop incident with some vertex u is replaced by a k -cycle containing u and $k - 1$ further vertices of degree 2.

Note that the k -subdivision of a forest is a forest. Together with Lemma 1, the following lemma implies that $Sub_k(H)$ belongs to \mathcal{G}_k for every multigraph H .

Lemma 3 *Let k be a positive integer. If the graph G contains an induced subgraph B such that*

- $B = Sub_k(H)$ for some connected multigraph H that contains a cycle, and
- every component K of $G - V(H)$ that contains a vertex from $V(B) \setminus V(H)$ satisfies $\nu_k(K) = 0$,

then $\nu_k(G) = \nu_k(G - V(H)) + n(H)$, and $\tau_k(G) = \tau_k(G - V(H)) + n(H)$.

Proof: Since H is connected and contains a cycle, it contains an edge e incident with some vertex r such that $H - e$ contains a spanning tree T of H . Rooting T in r , assigning e to r , and assigning to every other vertex of H , the edge to its parent within T , yields an injective function $f : V(H) \rightarrow E(H)$ such that u is incident with $f(u)$ for every vertex u of H .

Let \mathcal{P}_f be k -matching of order $n(H)$ in B that contains, for every vertex u of H , the k -path formed within B by u and the subdivided edge $f(u)$. Recall that the components of $G - V(H)$ that contain a vertex from $V(B) \setminus V(H)$ contain no k -paths. Therefore, adding \mathcal{P}_f to any k -matching in $G - V(H)$ yields $\nu_k(G) \geq \nu_k(G - V(H)) + n(H)$. Conversely, if \mathcal{P} is a k -matching in G , then, since every k -path in G that intersects $V(B)$ contains a vertex of H , the set \mathcal{P} contains at most $n(H)$ paths intersecting $V(B)$. Removing all such paths from \mathcal{P} yields a k -matching in $G - V(H)$, which implies $\nu_k(G) \leq \nu_k(G - V(H)) + n(H)$.

If X is a k -vertex cover in $G - V(H)$, then $X \cup V(H)$ is a k -vertex cover in G , which implies $\tau_k(G) \leq \tau_k(G - V(H)) + n(H)$. Now, let X be a k -vertex cover in G . Clearly, $X' = X \cap V(B)$ is a k -vertex cover in B . If some vertex u of H does not belong to X' , then X' must intersect all

subdivided edges of H incident with u , in particular, X' contains a vertex from the subdivided edge $f(u)$. Since f is injective, this easily implies that X' contains at least $n(H)$ vertices. Since $X \setminus X'$ is a k -vertex cover in $G - V(H)$, we obtain $\tau_k(G) \geq \tau_k(G - V(H)) + n(H)$. \square

For even values of k , Lemma 2 (i) and (iv) suggest yet another construction based on subdivisions of bipartite multigraphs. The following lemma captures the essence of this construction.

Lemma 4 *If k is a positive even integer, and $G = \text{Sub}_{k/2}(H)$ for some bipartite connected multigraph H that contains a cycle, then $\nu_k(G) = \tau_k(G)$.*

Proof: In view of the Kőnig-Egerváry theorem, and, since H is bipartite, it suffices to show that $\nu_k(G) \geq \nu(H)$ and $\tau_k(G) \leq \tau(H)$.

let M be a matching in H . Contracting the edges in M yields a connected multigraph that contains a cycle, and arguing similarly as in the proof of Lemma 3, we obtain the existence of an injective function $f : M \rightarrow E(H) \setminus M$ such that the edges e and $f(e)$ are adjacent for every edge e in M . Now, for every edge e in M , the $(k/2 + 1)$ -path corresponding to the subdivided edge e and the $(k/2 - 1)$ -path corresponding to the interior of the subdivided edge $f(e)$ form a k -path in G . Since M is a matching and f is injective, all these k -paths are disjoint, which implies $\nu(H) \leq \nu_k(G)$.

If X is a vertex cover in H , then every component of $G - X$ is a $(k/2 - 1)$ -subdivision of some star. Hence, $G - X$ contains no k -path, which implies $\tau(H) \geq \tau_k(G)$. \square

2 The graphs in \mathcal{G}_3 and \mathcal{G}_4

In this section we characterize the graphs in \mathcal{G}_k for $k \in \{3, 4\}$ by describing their blocks and conditions imposed on their cutvertices. As it turns out, the three generic examples of graphs in \mathcal{G}_k discussed in the introduction are the main building blocks of the considered graphs.

Recall that a cutvertex of a graph G is a vertex x of G for which $G - x$ has more components than G , and that a block of G is a maximal connected subgraph B of G such that B itself has no cutvertex. An endblock of G is a block of G that contains at most one cutvertex of G . A block is trivial if it is either K_1 or K_2 .

Let \mathcal{H}_3 be the set of all graphs G such that every non-trivial block B of G satisfies the following condition.

- (i) $B = \text{Sub}_3(H)$ for some multigraph H , and every cutvertex of G that belongs to B is a vertex of H .

Theorem 5 $\mathcal{G}_3 = \mathcal{H}_3$.

Proof: In order to show that $\mathcal{G}_3 \subseteq \mathcal{H}_3$, it suffices to show that $G \in \mathcal{H}_3$ for every connected graph G in \mathcal{G}_3 . If G is a tree, then all blocks of G are trivial, and, hence, $G \in \mathcal{H}_3$. If G is a cycle, then Lemma 2(i) implies that $n(G)$ is a multiple of 3, and, hence, $G = \text{Sub}_3(C_{n(G)/3}) \in \mathcal{H}_3$. Now, we may assume that G is neither a tree nor a cycle. Let B be a non-trivial block of G . By Lemma 2(i), the order of every cycle in B is a multiple of 3. Suppose that B contains a path $P : u_0 \dots u_\ell$ such that u_0 and u_ℓ have degree at least 3 in G , and $u_1, \dots, u_{\ell-1}$ have degree 2 in G . Since $B - u_1$ is connected, the path P is contained in a cycle C such that u_0 and u_ℓ both have neighbors outside of $V(C)$. By Lemma 2(iii), the length ℓ of P is a multiple of 3, in particular, no two vertices of B of degree at least 3 in G are adjacent. Let H be the multigraph that arises by replacing every path or cycle $u_0 u_1 u_2 u_3 \dots u_{3p-3} u_{3p-2} u_{3p-1} u_{3p}$ of length $3p$ such that u_0 and u_{3p} have degree at least 3 in G , and u_1, \dots, u_{3p-1} have degree 2 in G , by the path or cycle $u_0 u_3 \dots u_{3p-3} u_{3p}$ of length p . Clearly, $B = \text{Sub}_3(H)$, and every cutvertex of G that belongs to B is a vertex of H , that is, $G \in \mathcal{H}_3$. Altogether, we obtain $\mathcal{G}_3 \subseteq \mathcal{H}_3$.

It follows easily from its definition that \mathcal{H}_3 is a hereditary class of graphs, that is, it is closed under taking induced subgraphs. Therefore, in order to show the reverse inclusion $\mathcal{H}_3 \subseteq \mathcal{G}_3$, it suffices to show that $\nu_3(G) = \tau_3(G)$ for every connected graph G in \mathcal{H}_3 , which we do by induction on the order of G . If G is a tree, then Lemma 1 implies $\nu_3(G) = \tau_3(G)$. If G is a cycle, then the order of G is a multiple of 3, and, hence, $\nu_3(G) = \tau_3(G)$. Now, we may assume that G is neither a tree nor a cycle. Let B be a non-trivial block of G . Let $B = \text{Sub}_3(H)$ for some multigraph H such that every cutvertex of G that belongs to B is a vertex of H . By Lemma 3 applied to B , we obtain $\nu_3(G) = \nu_3(G - V(H)) + n(H)$ and $\tau_3(G) = \tau_3(G - V(H)) + n(H)$. Since \mathcal{H}_3 is hereditary, we obtain, by induction, $\nu_3(G - V(H)) = \tau_3(G - V(H))$, which implies $\nu_3(G) = \tau_3(G)$ and completes the proof. \square

For some positive integer p , let the graph $T(p)$ arise by adding an edge between the two vertices in a partite set of order 2 of the complete bipartite graph $K_{2,p}$. Note that $T(1)$ is a triangle, and that $T(2)$ arises by removing one edge from K_4 .

Let \mathcal{H}_4 be the set of all graphs G such that every non-trivial block B of G satisfies the following condition.

- (i) Either $B = \text{Sub}_2(H)$ for some bipartite multigraph H , and every cutvertex of G that belongs to B is a vertex of H ,
- (ii) or $B = K_4$ is an endblock,
- (iii) or $B = T(2)$ is an endblock, and, if B contains a cutvertex x of G , then x has degree 2 in B ,
- (iv) or $B = T(p)$ for some positive integer p , at most two cutvertices of G belong to B , every cutvertex of G that belongs to B has degree $p+1$ in B , and, if B contains two cutvertices of G , then there is one cutvertex x of G in B such that every vertex in $N_G(x) \setminus V(B)$ has degree 1 in G .

See Figure 1 for an illustration of (iv).




Figure 1: $T(p)$ as a non-endblock of a graph in \mathcal{G}_4 .

Theorem 6 $\mathcal{G}_4 = \mathcal{H}_4$.

Proof: As before, in order to show that $\mathcal{G}_4 \subseteq \mathcal{H}_4$, we show that $G \in \mathcal{H}_4$ for every connected graph $G \in \mathcal{G}_4$. If G is a tree, then clearly $G \in \mathcal{H}_4$. If G is a cycle, then Lemma 2(i) implies that $n(G)$ is either 3 or a multiple of 4, and, hence, $G \in \mathcal{H}_4$. Now, we may assume that G is neither a tree nor a cycle. Let B be a non-trivial block of G .

The three graphs G_1 , G_2 , and G_3 in Figure 2 are forbidden subgraphs for the graphs in \mathcal{G}_4 . In fact, each of these graphs contains a 4-path but has order less than 8, which implies that

adding edges yields graphs with 4-matching number 1. Conversely, their 4-vertex cover number is 2, and adding edges can only increase this value.




Figure 2: Three forbidden subgraphs for the graphs in \mathcal{G}_4 .

First, we assume that B contains two adjacent vertices x and y with exactly p common neighbors z_1, \dots, z_p , where $p \geq 2$. Let $Z = \{z_1, \dots, z_p\}$ and $U = \{x, y\} \cup Z$. If x has a neighbor x' in B outside of U , then, since B has no cutvertex, a path in $B - x$ between x' and $U \setminus \{x\}$ together with a suitable path within $B[U]$ yields a cycle of order at least 4 whose order is not a multiple of 4, contradicting Lemma 2(i). Hence, x , and, by symmetry, y do not have neighbors in B outside of U . A similar argument also implies that z_1, \dots, z_p do not have neighbors in B outside of U , which implies that $V(B) = U$.

If Z is not independent, and $p \geq 3$, then B contains a cycle of order 5, contradicting Lemma 2(i). Hence, if Z is not independent, then $p = 2$, which implies that B is K_4 . Since G does not contain G_1 as a subgraph, we obtain that B is an endblock, that is, B is as in (ii) in the definition of \mathcal{H}_4 . Hence, we may assume that Z is independent. If some vertex in Z is a cutvertex of G , then, since G does not contain G_1 or G_3 as a subgraph, we obtain that $p = 2$, and that B is an endblock, that is, B is as in (iii) in the definition of \mathcal{H}_4 . Hence, we may assume that no vertex in Z is a cutvertex of G , which implies that at most two cutvertices of G belong to B , and that every cutvertex of G that belongs to B has degree $p + 1$ in B . Furthermore, if B contains two cutvertices of G , then, since G does not contain G_2 as a subgraph, there is one cutvertex x of G in B such that every vertex in $N_G(x) \setminus V(B)$ has degree 1 in G , that is, B is as in (iv) in the definition of \mathcal{H}_4 .

Next, we assume that B contains a triangle with vertices x, y , and z , but that no two adjacent vertices in B have more than one common neighbor. Arguing as above, we obtain $V(B) = \{x, y, z\}$, and, since G does not contain G_1 or G_2 as a subgraph, it follows that B is as in (iv) in the definition of \mathcal{H}_4 . Hence, we may assume that B contains no triangle.

Suppose that B contains a path $P : u_0 \dots u_\ell$ such that u_0 and u_ℓ have degree at least 3 in G , and $u_1, \dots, u_{\ell-1}$ have degree 2 in G . Since $B - u_1$ is connected, the path P is contained in a cycle C such that u_0 and u_ℓ both have neighbors outside of $V(C)$. By Lemma 2(iv), the length ℓ of P is even, in particular, no two vertices of B of degree at least 3 in G are adjacent. Let H be the multigraph that arises by replacing every path or cycle $u_0 u_1 u_2 \dots u_{2p-2} u_{2p-1} u_{2p}$ of length $2p$ such that u_0 and u_{2p} have degree at least 3 in G , and u_1, \dots, u_{2p-1} have degree 2 in G , by the path or cycle $u_0 u_2 \dots u_{2p-2} u_{2p}$ of length p . Clearly, $B = \text{Sub}_2(H)$, and every cutvertex of G that belongs to B is a vertex of H , that is, B is as in (i) in the definition of \mathcal{H}_4 . Altogether, it follows that $G \in \mathcal{H}_4$, which implies $\mathcal{G}_4 \subseteq \mathcal{H}_4$.

Again, it follows easily from its definition that \mathcal{H}_4 is a hereditary class of graphs. Hence, in order to show the reverse inclusion $\mathcal{H}_4 \subseteq \mathcal{G}_4$, it suffices to show that $\nu_4(G) = \tau_4(G)$ for every connected graph G in \mathcal{H}_4 , which we do by induction on the sum of the order and the size of G . As in the proof of Theorem 5, we may assume that G is neither a tree nor a cycle. If G contains a block B as in (ii) or (iii) in the definition of \mathcal{H}_4 , then it is easy to see that

$\nu_4(G) = \nu_4(G - V(B)) + 1$ and $\tau_4(G) = \tau_4(G - V(B)) + 1$. If G contains a block B as in (iv) in the definition of \mathcal{H}_4 , then we consider a graph G' obtained from G by removing an edge of B that is incident with every cutvertex in B . This graph G' is in \mathcal{H}_4 , has less edges than G , and satisfies $\nu_4(G) = \nu_4(G')$ and $\tau_4(G) = \tau_4(G')$. In all these cases, we obtain $\nu_4(G) = \tau_4(G)$ by induction. Hence, we may assume that G contains no such block.

Let B be a non-trivial block of G . Let X be the set of cutvertices of G that belong to B . For $x \in X$, let G_x be the component of $G - (V(B) \setminus \{x\})$ that contains x . We may assume that B is chosen in such a way that there is a vertex x^* in X such that G_x is a tree for every vertex x in $X \setminus \{x^*\}$. If some tree G_x with x in $X \setminus \{x^*\}$ contains a 4-path, then Lemma 1 implies the existence of an induced subgraph G' of G with $\nu_4(G) = \nu_4(G') + 1$ and $\tau_4(G) = \tau_4(G') + 1$, and $\nu_4(G) = \tau_4(G)$ follows by induction. Hence, for every vertex x in $X \setminus \{x^*\}$, the tree G_x is a star. Let X' be the set of vertices x in $X \setminus \{x^*\}$, for which G_x is not a star with center vertex x , that is, G_x contains a 3-path P_x starting in x . Let B' be the union of B and the paths P_x for x in X' . If $B = \text{Sub}_2(H)$, where H is as in (i) in the definition of \mathcal{H}_4 , then $B' = \text{Sub}_2(H')$ for the multigraph H' that arises from H by attaching a vertex of degree 1 to every vertex in X' . Clearly, H' is bipartite, connected, and contains a cycle.

First, suppose that x^* belongs to some minimum vertex cover in H' . By the Kőnig-Egerváry Theorem, this implies that every maximum matching in H' contains an edge incident with x^* . Let M be a maximum matching in H' . Similarly as in the proofs of Lemma 3 and Lemma 4, we obtain the existence of an injective function $f : M \rightarrow E(H') \setminus M$ such that the edges e and $f(e)$ are adjacent for every edge e in M . Adding the $\nu(H')$ disjoint 4-paths in B' corresponding to M , each formed using a subdivided edge e in M and the interior of the subdivided edge $f(e)$, to a maximum 4-matching in $G_{x^*} - x^*$ implies $\nu_4(G) \geq \nu_4(G_{x^*} - x^*) + \nu(H')$. Adding to a minimum 4-vertex cover in $G_{x^*} - x^*$ a minimum vertex cover in H' that contains x^* but none of the vertices of degree 1 in $V(H') \setminus V(H)$, yields a 4-vertex cover in G , which implies $\tau_4(G) \leq \tau_4(G_{x^*} - x^*) + \tau(H')$. Now, by induction and the Kőnig-Egerváry Theorem for H' , we obtain $\nu_4(G) \geq \nu_4(G_{x^*} - x^*) + \nu(H') = \tau_4(G_{x^*} - x^*) + \tau(H') \geq \tau_4(G) \geq \nu_4(G)$, that is, $\nu_4(G) = \tau_4(G)$.

Now, we may assume that x^* belongs to no minimum vertex cover in H' , which implies that every minimum vertex cover in H' contains all neighbors of x^* in H' . Furthermore, by the Kőnig-Egerváry Theorem, this implies that some maximum matching M in H' contains no edge incident with x^* . Similarly as in the proof of Lemma 4, we obtain the existence of an injective function $f : \{x^*\} \cup M \rightarrow E(H') \setminus M$ such that x^* and $f(x^*)$ are incident, and e and $f(e)$ are adjacent for every $e \in M$. Let G' arise from G_{x^*} by attaching a vertex of degree 1 to x^* , corresponding to the internal vertex of the subdivided version of $f(x^*)$. Arguing similarly as above, we obtain $\nu_4(G) \geq \nu_4(G') + \nu(H')$ and $\tau_4(G) \leq \tau_4(G') + \tau(H')$, and $\nu_4(G) = \tau_4(G)$ follows by induction and the Kőnig-Egerváry Theorem for H' , which completes the proof. \square

3 Graphs without short cycles in \mathcal{G}_k for odd k

For general k , an explicit characterization of \mathcal{G}_k , similar to the ones that we obtained for \mathcal{G}_3 and \mathcal{G}_4 in the previous section, might not be possible. For instance, every graph of order less than k without a cutvertex is a block of some graph in \mathcal{G}_k , and already in the characterization of \mathcal{G}_4 , we encountered sporadic blocks that required special attention. Nevertheless, if we consider an odd k as well as the graphs in \mathcal{G}_k that do not contain short cycles, then the sporadic blocks should disappear.

Let k be a positive odd integer. Let \mathcal{G}'_k be the set of all graphs in \mathcal{G}_k that contain no cycle of order less than k . Note that \mathcal{G}'_3 actually coincides with \mathcal{G}_3 . Let \mathcal{H}'_k be the set of all graphs G such that every non-trivial block B of G satisfies the following condition.

- (i) $B = \text{Sub}_k(H)$ for some multigraph H , and every component K of $G - V(H)$ that contains a vertex from $V(B) \setminus V(H)$ is a tree without a k -path.

As before our goal is to show that \mathcal{G}'_k and \mathcal{H}'_k coincide. The following lemma deals with some rather simple graphs in \mathcal{G}'_k for which it is surprisingly difficult to show that they belong to \mathcal{H}'_k .

Lemma 7 *Let k be a positive odd integer, and let p be a positive integer. If the graph G in \mathcal{G}_k arises from the cycle $C_{pk} : u_1 u_2 \dots u_{pk} u_1$ of order pk by attaching, for every i in $[pk]$, a path P_i of order p_i to the vertex u_i , where $0 \leq p_i < (k-1)/2$, then $G \in \mathcal{H}'_k$.*

Proof: It suffices to show that $\nu_k(G) = p$. Indeed, if $\nu_k(G) = p$, then $\nu_k(G) = \tau_k(G) = \tau_k(C_{pk}) = p$, and Lemma 2(ii) implies the existence of a minimum k -vertex cover X of G with $X = \{u_{i+(j-1)k} : j \in [p]\}$ for some $i \in [k]$. It follows that the unique cycle C_{pk} in G , which is the only non-trivial block of G , is the k -subdivision of the cycle $u_i u_{i+k} u_{i+2k} \dots u_{i+(p-1)k} u_i$ of order p with vertex set X , and that every component of $G - X$ is a tree without a k -path, that is, $G \in \mathcal{H}'_k$. Hence, for a contradiction, we assume that $\nu_k(G) > p$.

Since removing an endvertex from G can reduce the k -matching number by at most 1, we may assume, by considering a suitable induced subgraph of G , that $\nu_k(G) = p + 1$, and that $\nu_k(G - x) = p$ for every endvertex x of G . For i in $[pk]$, let P_i be the path $u_i^1 \dots u_i^{p_i}$, where, for $p_i \geq 1$, the vertex u_i^1 is a neighbor of u_i . Note that the order of G is $pk + p_1 + \dots + p_{pk}$, and that the endvertices of G are the vertices $u_i^{p_i}$ for those i in $[pk]$ with $p_i \geq 1$. Let \mathcal{P} be a maximum k -matching in G . A path P in \mathcal{P} that is not completely contained in C_{pk} is called *special*. By the choice of G , for every special path P in \mathcal{P} , there are two distinct indices i and j in $[pk]$ with $\max\{p_i, p_j\} \geq 1$ such that P is the path

$$\underbrace{u_i^{p_i} \dots u_i^1}_{P_i} \underbrace{u_i u_{i+1} \dots u_{j-1} u_j}_{\subseteq C_{pk}} \underbrace{u_j^1 \dots u_j^{p_j}}_{P_j}, \quad (1)$$

where we identify indices modulo pk for the subpath $u_i u_{i+1} \dots u_{j-1} u_j$ of P that is contained in C_{pk} . If $p_i \geq 1$, then P is said to have the *left leg* P_i . If $p_j \geq 1$, then P is said to have the *right leg* P_j . Since G contains at most $\nu_k(C_{pk}) = p$ disjoint non-special paths, and every special path contains at most $2 \max\{p_1, \dots, p_{pk}\} < k - 1$ vertices that do not belong to C_{pk} , the set \mathcal{P} contains at least two special paths. By the choice of G , for every i in $[pk]$ with $p_i \geq 1$, the path P_i is either the left leg or the right leg of some path in \mathcal{P} .

Let i in $[pk]$ be such that P_i is the left leg of some path P in \mathcal{P} as in (1). By the choice of G , the graph $G_i = G - u_i^{p_i}$ satisfies $\nu_k(G_i) = p$. Similarly as above, this implies the existence of a minimum k -vertex cover X_i in G_i with $X_i = \{u_{r+(s-1)k} : s \in [p]\}$ for some $r \in [k]$. We will show that $r = i - p_i$, which implies that X_i is uniquely determined. Since X_i has order p , and intersects all p paths in $\mathcal{P} \setminus \{P\}$, it contains no vertex of P , and, hence, no vertex from $u_i u_{i+1} \dots u_{j-1} u_j$. Since $G_i - X_i$ contains no k -path, this implies that $r \in \{i - p_i, i - p_i + 1, \dots, i - 1\}$. Now, if r is not $i - p_i$, then $r \in \{i - p_i + 1, \dots, i - 1\}$, the set X_i contains no vertex from $u_i u_{i+1} \dots u_{i+k-p_i+1}$, and $u_i^{p_i-1} \dots u_i^1 u_i u_{i+1} \dots u_{i+k-p_i+1}$ is a k -path in $G_i - X_i$, which is a contradiction. Hence, $r = i - p_i$ as claimed. Symmetrically, if P_i is the right leg of some path in \mathcal{P} , then G_i has a unique minimum k -vertex cover X_i with $X_i = \{u_{i+p_i+(s-1)k} : s \in [p]\}$.

We consider some cases.

Case 1 *No path in \mathcal{P} has a right leg.*

In this case, every special path in \mathcal{P} contains at most $\max\{p_1, \dots, p_{pk}\} < (k-1)/2$ vertices that do not belong to C_{pk} , which implies that \mathcal{P} contains at least three special paths. By symmetry, we may assume that the indices r , s , and t in $[pk]$ are chosen in such a way that

- $r < s < t$,

- $p_s \leq p_t$,
- P_r , P_s , and P_t are left legs of three special paths in \mathcal{P} , and
- no other special path in \mathcal{P} intersects the subpath $u_r \dots u_s \dots u_t$ of C_{pk} .

By the choice of G , in this case it follows that every vertex of C_{pk} belongs to some path in \mathcal{P} . Therefore, the final condition in the choice of r , s , and t implies that

$$s \equiv (r + k - p_r) \pmod{k} \quad \text{and} \quad t \equiv (s + k - p_s) \pmod{k}.$$

Since X_r contains the vertex u_{r-p_r} , this implies that $u_s \in X_r$, and that X_r contains no vertex from $u_{t-k+p_s+1}u_{t-k+p_s+2} \dots u_t$. See Figure 3 for an illustration.




Figure 3: The situation in Case 1, where vertices in X_r are indicated by the square boxes, and the paths in \mathcal{P} are shown in bold.

Nevertheless, since $p_s \leq p_t$, the graph $G_r - X_r$ contains the path $u_{t-k+p_s+1}u_{t-k+p_s+2} \dots u_t u_t^1 \dots u_t^{p_t}$ of order $k - 1 - p_s + 1 + p_t \geq k$, which is a contradiction.

Case 2 *Some special path in \mathcal{P} has a right leg, and some special path in \mathcal{P} has a left leg.*

By symmetry, we may assume that the indices s and t in $[pk]$ are such that

- $s < t$,
- $p_s \leq p_t$,
- P_s is the right leg of a special path in \mathcal{P} , and P_t is the left leg of a special path in \mathcal{P} , and
- no other special path in \mathcal{P} intersects the subpath $u_s \dots u_t$ of C_{pk} .

We may assume that the non-special paths in \mathcal{P} that intersect $u_s \dots u_t$ are chosen in such a way that their removal from $u_s \dots u_t$ leaves a path of the form $u_s \dots u_{s+s'}$ for some $s' \geq 0$. Since $\nu_k(G_s) = p$, we have $s' \leq p_s - 1$. If $s' \leq p_s - 2$, then X_s contains no vertex from $u_{t-k+p_s-s'}u_{t-k+p_s-s'+1} \dots u_t$, and $G_s - X_s$ contains the path $u_{t-k+p_s-s'}u_{t-k+p_s-s'+1} \dots u_t u_t^1 \dots u_t^{p_t}$ of order $k - p_s + s' + 1 + p_t > k$, which is a contradiction. See Figure 4 for an illustration. Hence, we obtain $s' = p_s - 1$, which implies that $u_t \in X_s$.




Figure 4: Illustration of the proof that $s' = p_s - 1$.

If the path P' in \mathcal{P} whose right leg is P_s also has a left leg, say P_r for some $r < s$, then X_r contains x_{r-p_r} , and, hence, also u_{s+p_s+1} as well as u_{t+1} but no vertex from $u_{t-k+2}u_{t-k+3} \dots u_t$. See Figure 5 for an illustration.




Figure 5: Illustration of the proof that P' has no left leg.

Now, $G_r - X_r$ contains the path $u_{t-k+2}u_{t-k+3} \dots u_t u_t^1 \dots u_t^{p_t}$ of order $k - 2 + 1 + p_t \geq k$, which is a contradiction. Hence, P' has no left leg, and equals $u_{s-k+p_s+1}u_{s-k+p_s+2} \dots u_s u_s^1 \dots u_s^{p_s}$.

Let $r < s$ be maximum such that some special path P'' in \mathcal{P} contains u_r . By the choice of G , and, since P' has no left leg, we obtain that $r \equiv (s - k + p_s) \pmod k$.

First, suppose that $p_r = 0$, that is, P'' has no right leg. Since P'' is special, it has a left leg, say P_q for some $q < r$. Here things work as previously; X_q contains u_{q-p_q} , and, hence, also u_{r+1} , u_{s-k+p_s+1} , u_{s+p_s+1} , and u_{t+1} but no vertex from $u_{t-k+2}u_{t-k+3} \dots u_t$. Now, $G_q - X_q$ contains the path $u_{t-k+2}u_{t-k+3} \dots u_t u_t^1 \dots u_t^{p_t}$ of order $k - 2 + 1 + p_t \geq k$, which is a contradiction. Hence, $p_r \geq 1$, that is, the path P'' has P_r as its right leg. If $p_r \geq p_t$, then X_t contains u_{t-p_t} , and, hence, also $u_{s-p_t+p_s}$ as well as u_{r+k-p_t} but no vertex from $u_r u_{r+1} \dots u_{r+k-p_t-1}$. See Figure 6 for an illustration.




Figure 6: Illustration of the proof that $p_r \not\leq p_t$.

Now, $G_t - X_t$ contains the path $u_r^{p_r} \dots u_r^1 u_r u_{r+1} \dots u_{r+k-p_t-1}$ of order $p_r + 1 + k - p_t - 1 \geq k$, which is a contradiction. Conversely, if $p_r < p_t$, then X_r contains u_{r+p_r} , and, hence, also u_{t-k+p_r} but no vertex from $u_{t-k+p_r+1}u_{t-k+p_r+2} \dots u_t$. Now, $G_r - X_r$ contains the path $u_{t-k+p_r+1}u_{t-k+p_r+2} \dots u_t u_t^1 \dots u_t^{p_t}$ of order $k - p_r - 1 + 1 + p_t \geq k$, which is a contradiction. This completes the proof. \square

We proceed to the main result in this section, which actually contains Theorem 5 as a special case. In view of its simplicity, we kept the separate proof of Theorem 5.

Theorem 8 $\mathcal{G}'_k = \mathcal{H}'_k$ for every positive odd integer k .

Proof: As before, in order to show that $\mathcal{G}'_k \subseteq \mathcal{H}'_k$, we show that $G \in \mathcal{H}'_k$ for every connected graph $G \in \mathcal{G}'_k$. By Lemma 2(i), the order of every cycle in G is a multiple of k . We may again assume that G is neither a tree nor a cycle. Let B be a non-trivial block of G .

First, we assume that B is not just a cycle, that is, it contains vertices that are of degree at least 3 in B . Suppose that B contains a path $P : u_0 \dots u_\ell$ such that u_0 and u_ℓ have degree at least 3 in B , and $u_1, \dots, u_{\ell-1}$ have degree 2 in B . Since $B - u_1$ is connected, the path P is contained in a cycle C such that u_0 and u_ℓ both have neighbors outside of $V(C)$, say u_0^1 and u_ℓ^1 , respectively. Let P_0 be a shortest path in $B - u_0$ between u_0^1 and $V(C) \setminus \{u_0\}$. Since the order of every cycle in G is a multiple of k , and, since k is odd, it follows that P_0 has length -1 modulo k , which implies that $B - V(C)$ contains a path P'_0 of order $(k - 1)/2$ starting in u_0^1 . Similarly, $B - V(C)$ contains a path P'_ℓ of order $(k - 1)/2$ starting in u_ℓ^1 . If G' is the subgraph of G induced by $V(C) \cup V(P'_0) \cup V(P'_\ell)$, then $\frac{n(C)}{k} \leq \nu_k(G') \leq \left\lfloor \frac{n(C) + n(P'_0) + n(P'_\ell)}{k} \right\rfloor = \frac{n(C)}{k}$. It follows that every minimum k -vertex cover X' of G' is also a minimum k -vertex cover of C , and, hence, as described in Lemma 2(ii). In view of P'_0 , P'_ℓ , and the subpaths of C not covered by X' , it follows that the vertices u_0 and u_ℓ must both belong to X' . This implies that the length

ℓ of P is a multiple of k . Let H be the multigraph that arises by replacing every path or cycle $u_0u_1 \dots u_{pk}$ of length pk such that u_0 and u_{pk} have degree at least 3 in B , and u_1, \dots, u_{pk-1} have degree 2 in B , by the path or cycle $u_0u_k \dots u_{pk}$ of length p . Clearly, $B = \text{Sub}_k(H)$.

Let K be a component of $G - V(H)$ that contains a vertex from $V(B) \setminus V(H)$. Let uv be an edge of H such that K intersects the subdivided edge uv . Since B is a block of G , the component K intersects $V(B) \setminus V(H)$ exactly in the interior of the subdivided edge uv . Let $P : uw_1 \dots w_{k-1}v$ be the path in G corresponding to the subdivided edge uv . Suppose, for a contradiction, that K contains a k -path. This implies that we may assume, by symmetry, that there is some $i \in [(k-1)/2]$, and a path $Q : x_1 \dots x_i$ in $K - V(B)$ such that x_i is adjacent to w_i . Let C be a cycle in B containing P . Similarly as above, we obtain the existence of a path R of order $(k-1)/2$ in $B - V(C)$ such that u is adjacent to an endvertex of R . If G' is the subgraph of G induced by $V(C) \cup V(Q) \cup V(R)$, then $\nu_k(G') = \frac{n(C)}{k}$. Therefore, every minimum k -vertex cover of G' is also a minimum k -vertex cover of C , and, hence, as described in Lemma 2(ii). In view of R and the subpaths of C not covered by X' , it follows that u must belong to X' . But now, $x_1 \dots x_i w_i \dots w_{k-1}$ is a k -path in $G' - X'$, which is a contradiction. Altogether, it follows that K contains no k -path, which implies that K is a tree without a k -path. Hence, B is as in (i) in the definition of \mathcal{H}'_k .

Next, we assume that B is a cycle $C : u_1 \dots u_{pk}$. For every i in $[pk]$, let p_i be the maximum length of a path in $G - (V(B) \setminus \{u_i\})$ starting in the vertex u_i . First, suppose that $\max\{p_1, \dots, p_{pk}\} \geq (k-1)/2$. By symmetry, we may assume that $p_1 \geq (k-1)/2$. Let $X = \{u_{1+(j-1)k} : j \in [p]\}$. Clearly, $B = \text{Sub}_k(H)$, where H is the cycle $u_1u_{1+k} \dots u_{1+(p-1)k}u_1$ with vertex set X .

Let K be a component of $G - V(H)$ that contains a vertex from $V(B) \setminus V(H)$. If K contains a k -path, then, by symmetry, we may assume that there is some index i in $[pk]$ such that $1 \leq (i-1) \bmod k \leq (k-1)/2$ and, p_i is at least $(i-1) \bmod k$. Now, G contains a subgraph G' that arises from B by attaching a path of order $(k-1)/2$ to u_1 , and a path of order $(i-1) \bmod k$ to u_i . As before $\nu_k(G') = \frac{n(B)}{k}$, and Lemma 2(ii) implies that every minimum k -vertex cover X' of G' must contain u_1 , and that $G' - X'$ still contains a k -path using the path attached to u_i , which is a contradiction. Altogether, it follows that K contains no k -path, which implies that K is a tree without a k -path. Hence, B is as in (i) in the definition of \mathcal{H}'_k .

Now, we may assume that $\max\{p_1, \dots, p_{pk}\} < (k-1)/2$. This implies that, for every i in $[pk]$, the component G_{u_i} of $G - (V(B) \setminus \{u_i\})$ that contains u_i , is a tree without a k -path. Let G' be the induced subgraph of G that arises from G by removing, for every i in $[pk]$, all of G_{u_i} except for a path of length p_i starting in the vertex u_i . By Lemma 7, the graph G' belong to \mathcal{H}'_k , which easily implies that also G belongs to \mathcal{H}'_k . Altogether, we obtain $\mathcal{G}'_k \subseteq \mathcal{H}'_k$.

Again, it follows easily from its definition that \mathcal{H}'_k is a hereditary class of graphs, and, hence, in order to show the reverse inclusion $\mathcal{H}'_k \subseteq \mathcal{G}'_k$, it suffices to show that $\nu_k(G) = \tau_k(G)$ for every connected graph G in \mathcal{H}'_k . This now follows very easily by induction on the order using Lemma 1 and Lemma 3, which completes the proof. \square

4 Conclusion

It is not difficult to extract from our results all minimal forbidden induced subgraphs for the graph classes \mathcal{G}_3 , \mathcal{G}_4 , and \mathcal{G}'_k for odd k at least 5. Furthermore, our results imply that the graphs in these classes can be recognized efficiently, and that there are simple combinatorial polynomial time algorithms that determine maximum k -matchings and minimum k -vertex covers for these graphs. Apart from extending our characterizations, a natural open problem concerns the complexity of recognizing the graphs in \mathcal{G}_k for general fixed k . We pose the following optimistic conjecture.

Conjecture 9 *For every fixed positive integer k , it can be decided in polynomial time whether a given graph belongs to \mathcal{G}_k .*

Lemma 2(i) easily implies that every graph in \mathcal{G}_k has minimum degree at most k . This implies that the graphs in \mathcal{G}_k are k -degenerate, which might be a useful property for their recognition.

For $k \in \{3, 4\}$, our results imply that $\nu_k(H) = \tau_k(H)$ for every not necessarily induced subgraph H of every graph G in \mathcal{G}_k . For $k = 1$, the same trivially holds, and, also for $k = 2$, the same holds, since graphs are bipartite if and only if all their not necessarily induced subgraphs are bipartite. We believe that these observations generalize, and pose the following conjecture.

Conjecture 10 *For every positive integer k , the set \mathcal{G}_k equals the set of all graphs G such that $\nu_k(H) = \tau_k(H)$ for every subgraph H of G .*

One proof of the König-Egerváry Theorem, as well as many polyhedral insights concerning matchings in bipartite graphs, rely on the total unimodularity of the vertex versus edge incidence matrices of bipartite graphs. Unfortunately, for integers k at least 3, the vertex versus k -path incidence matrices of the graphs in \mathcal{G}_k are not totally unimodular. If $G = \text{Sub}_3(H)$ for some graph H with a vertex u of degree at least 3 for instance, then considering three suitable 3-paths containing u as central vertex, and three suitable neighbors of u on these paths, implies that the vertex versus 3-path incidence matrix A of G contains the vertex versus edge incidence matrix of C_3 as a submatrix, that is, A is not totally unimodular.

References

- [1] R. Boliac, K. Cameron, V.V. Lozin, On computing the dissociation number and the induced matching number of bipartite graphs, *Ars Combinatorics* 72 (2004) 241-253.
- [2] B. Brešar, M. Jakovac, J. Katrenič, G. Semanišin, A. Taranenko, On the vertex k -path cover, *Discrete Applied Mathematics* 161 (2013) 1943-1949.
- [3] B. Brešar, F. Kardoš, J. Katrenič, G. Semanišin, Minimum k -path vertex cover, *Discrete Applied Mathematics* 159 (2011) 1189-1195.
- [4] E. Egerváry, Über kombinatorische Eigenschaften von Matrizen, *Matematikai és Fizikai Lapok* 38 (1931) 16-28.
- [5] M.R. Garey, D.S. Johnson, *Computers and Intractability: A Guide to the Theory of NP-Completeness*, W.H. Freeman and Co. pp. x+338, 1979.
- [6] F. Kardoš, J. Katrenič, I. Schiermeyer, On computing the minimum 3-path vertex cover and dissociation number of graphs, *Theoretical Computer Science* 412 (2011) 7009-7017.
- [7] J. Katrenič, A faster FPT algorithm for 3-path vertex cover, *Information Processing Letters* 116 (2016) 273-278.
- [8] D. König, Graphen und Matrices, *Matematikai és Fizikai Lapok* 38 (1931) 116-119.
- [9] X. Li, Z. Zhang, X. Huang, Approximation algorithms for minimum (weight) connected k -path vertex cover, *Discrete Applied Mathematics* 205 (2016) 101-108.
- [10] J. Tu, L. Wu, J. Yuan, L. Cui, On the vertex cover P_3 problem parameterized by treewidth, *Journal of Combinatorial Optimization* 34 (2017) 414-4

- [11] J. Tu, W. Zhou, A primal-dual approximation algorithm for the vertex cover P_3 problem, Theoretical Computer Science 412 (2011) 7044-7048.
- [12] M. Xiao, S. Kou, Exact algorithms for the maximum dissociation set and minimum 3-path vertex cover problems, Theoretical Computer Science 657 (2017) 86-97.
- [13] M. Yannakakis, Node-deletion problems on bipartite graphs, SIAM Journal on Computing 10 (1981) 310-327.