

Repetition avoidance in products of factors

Pamela Fleischmann, Pascal Ochem, Kamellia Reshadi

► **To cite this version:**

Pamela Fleischmann, Pascal Ochem, Kamellia Reshadi. Repetition avoidance in products of factors. Theoretical Computer Science, Elsevier, 2019, 791, pp.123-126. 10.1016/j.tcs.2019.04.013 . lirmm-02500480

HAL Id: lirmm-02500480

<https://hal-lirmm.ccsd.cnrs.fr/lirmm-02500480>

Submitted on 6 Mar 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Repetition avoidance in products of factors

Pamela Fleischmann* Pascal Ochem† Kamellia Reshadi‡

Abstract

We consider a variation on a classical avoidance problem from combinatorics on words that has been introduced by Mousavi and Shallit at DLT 2013. Let $\mathbf{pexp}_i(w)$ be the supremum of the exponent over the products of i factors of the word w . The repetition threshold $\text{RT}_i(k)$ is then the infimum of $\mathbf{pexp}_i(w)$ over all words $w \in \Sigma_k^\omega$. Mousavi and Shallit obtained that $\text{RT}_i(2) = 2i$ and $\text{RT}_2(3) = \frac{13}{4}$. We show that $\text{RT}_i(3) = \frac{3i}{2} + \frac{1}{4}$ if i is even and $\text{RT}_i(3) = \frac{3i}{2} + \frac{1}{6}$ if i is odd and $i \geq 3$.

Keywords: Words; Repetition avoidance.

Acknowledgements: This work was partially supported by the ANR project CoCoGro (ANR-16-CE40-0005).

1 Introduction

A *repetition* in a word w is a pair of words p and e such that pe is a factor of w , p is non-empty, and e is a prefix of pe . If pe is a repetition, then its *period* is $|p|$ and its *exponent* is $\frac{|pe|}{|p|}$. A word is α^+ -free (resp. α -free) if it contains no repetition with exponent β such that $\beta > \alpha$ (resp. $\beta \geq \alpha$).

Given $k \geq 2$, Dejean [2] defined the repetition threshold $\text{RT}(k)$ for k letters as the smallest α such that there exists an infinite α^+ -free word over a k -letter alphabet. Dejean initiated the study of $\text{RT}(k)$ in 1972 for $k = 2$ and $k = 3$. Her work was followed by a series of papers which determine the exact value of $\text{RT}(k)$ for any $k \geq 2$.

*Kiel University, Germany. fpa@informatik.uni-kiel.de

†LIRMM, Université de Montpellier, CNRS, Montpellier, France. ochem@lirmm.fr

‡Kiel University, Germany. kre@informatik.uni-kiel.de

- $\text{RT}(2) = 2$ [2];
- $\text{RT}(3) = \frac{7}{4}$ [2];
- $\text{RT}(4) = \frac{7}{5}$ [7];
- $\text{RT}(k) = \frac{k}{k-1}$, for $k \geq 5$ [1, 4, 8].

Mousavi and Shallit [5] have considered two notions related to the repetition threshold.

The first notion considers repetitions in conjugates of factors of the infinite word. A word is circularly r^+ -free if it does not contain a factor pxs such that sp is a repetition of exponent strictly greater than r . Let $\Sigma_k = \{0, 1, \dots, k-1\}$. The smallest real number r such that w is circularly r^+ -free is denoted by $\text{cexp}(w)$. Let $\text{RTC}(k)$ be the minimum of $\text{cexp}(w)$ over every $w \in \Sigma_k^\omega$.

The second notion considers repetitions in concatenations of a fixed number of factors of the infinite word. Let $\text{pexp}_i(w)$ be the smallest real number r such that every product of i factors of w is r^+ -free. Let $\text{RT}_i(k)$ be the minimum of $\text{pexp}_i(w)$ over every $w \in \Sigma_k^\omega$. Notice that $\text{RT}_i(k)$ generalizes the classical notion of repetition threshold which corresponds to the case $i = 1$, that is, $\text{RT}_1(k) = \text{RT}(k)$ for every $k \geq 2$.

Our first result shows that the case $i = 2$ corresponds to the first notion of repetition avoidance in conjugates.

Theorem 1. $\text{RT}_2(k) = \text{RTC}(k)$ for every $k \geq 2$.

Mousavi and Shallit [5] have considered the binary alphabet and obtained that $\text{RT}_i(2) = 2i$ for every $i \geq 1$. Our second result considers the ternary alphabet and gives the value of $\text{RT}_i(3)$ for every $i \geq 1$. This extends the result of Dejean [2] that $\text{RT}_1(3) = \frac{7}{4}$ and the result of Mousavi and Shallit [5] that $\text{RT}_2(3) = \frac{13}{4}$.

Theorem 2.

- $\text{RT}_i(3) = \frac{3i}{2} + \frac{1}{4}$ if $i = 1$ or i is even.
- $\text{RT}_i(3) = \frac{3i}{2} + \frac{1}{6}$ if i is odd and $i \geq 3$.

2 Proofs

Proof of Theorem 1.

The language of words in Σ_k^* avoiding circular repetitions of exponent at least e (or strictly greater than e) is a factorial language. As it is well-known [3], if a factorial language is infinite, then it contains a uniformly recurrent word w . By Proposition 14 in [5], $\text{pexp}_2(w) = \text{cexp}(w)$. This implies that $\text{RT}_2(k) = \text{RTC}(k)$. \square

To obtain the two equalities of Theorem 2, we show the two lower bounds and then the two upper bounds.

Proof of $\text{RT}_i(3) \geq \frac{3i}{2} + \frac{1}{4}$ for every even i .

Mousavi and Shallit [5] have proved that $\text{RT}_2(3) = \frac{13}{4}$, which settles the case $i = 2$. We have double checked their computation of the lower bound $\text{RT}_2(3) \geq \frac{13}{4}$. Suppose that i is a fixed even integer and that w_3 is an infinite ternary word. The lower bound for $i = 2$ implies that there exists two factors u and v such that $uv = t^e$ with $e \geq \frac{13}{4}$. Thus, the prefix t^3 of uv is also a product of two factors of w_3 . So we can form the i -terms product $(t^3)^{i/2-1}uv$ which is a repetition of the form t^x with exponent $x = 3\left(\frac{i}{2} - 1\right) + e \geq 3\left(\frac{i}{2} - 1\right) + \frac{13}{4} = \frac{3i}{2} + \frac{1}{4}$. This is the desired lower bound. \square

Proof of $\text{RT}_i(3) \geq \frac{3i}{2} + \frac{1}{6}$ for every odd $i \geq 3$.

Suppose that $i \geq 3$ is a fixed odd integer, that is, $i = 2j + 1$. Suppose that w_3 is a recurrent ternary word such that the product of i factors of w_3 is never a repetition of exponent at least $\frac{3i}{2} + \frac{1}{6} = 3j + \frac{5}{3}$. First, w_3 is square-free since otherwise there would exist an i -terms product of exponent $2i$. Also, w_3 does not contain two factors u and v with the following properties:

- $uv = t^3$,
- $u = t^e$ with $e \geq \frac{5}{3}$.

Indeed, this would produce the i -terms product $(uv)^j u$ which is a repetition of the form t^x with exponent $x = 3j + e \geq 3j + \frac{5}{3}$.

So if a , b , and c are distinct letters, then w_3 does not contain both $u = abcab$ and $v = cabc$ and w_3 does not contain both $u = abcbabc$ and $v = babcb$. A computer check shows that no infinite ternary square-free word satisfies this property. This proves the desired lower bound. \square

Proof of $\text{RT}_i(3) \leq \frac{3i}{2} + \frac{1}{4}$ for every even i .

Let i be any even integer at least 2. To prove this upper bound, it is sufficient to construct a ternary word w satisfying $\text{pexp}_i(w) \leq \frac{3i}{2} + \frac{1}{4}$. The ternary morphic word used in [5] to obtain $\text{RT}_2(3) \leq \frac{13}{4}$ seems to satisfy the property. However, it is easier for us to consider another construction. Let us show that the image of every $7/5^+$ -free word over Σ_4 by the following 45-uniform morphism satisfies $\text{pexp}_i \leq \frac{3i}{2} + \frac{1}{4}$.

$0 \mapsto 010201210212021012102010212012101202101210212$
 $1 \mapsto 010201210212012101202101210201021202101210212$
 $2 \mapsto 010201210120212012102120210121021201210120212$
 $3 \mapsto 010201210120210121021201210120212012102010212$

Recall that a word is (β^+, n) -free if it does not contain a repetition with period at least n and exponent strictly greater than β . First, we check that such ternary images are $\left(\frac{202}{135}^+, 36\right)$ -free using the method in [6]. By Lemma 2.1 in [6], it is sufficient to check this freeness property for the image of every $7/5^+$ -free word over Σ_4 of length smaller than $\frac{2 \times \frac{202}{135}}{\frac{202}{135} - 5} < 32$. Since $\frac{202}{135} < \frac{3}{2}$, the period of every repetition formed from i pieces and with exponent at least $\frac{3i}{2}$ must be at most 35. Then we check exhaustively by computer that the ternary images do not contain two factors u and v such that

- $uv = t^e$,
- $e > 3$,
- $9 \leq |t| \leq 35$.

Thus, the period of every repetition formed from i pieces and with exponent strictly greater than $\frac{3i}{2}$ must be at most 8. So we only need to check that $\text{pexp}_i \leq \frac{3i}{2} + \frac{1}{4}$ for i -terms products that are repetitions of period at most 8.

Now the period is bounded, but i can still be arbitrarily large, a priori. For every factor t of length at most 8, we define $\text{pexp}_{i,t}$ as the length of a largest factor of t^ω that is a i -terms product, divided by $|t|$. We actually consider conjugacy classes, since if t' is a conjugate of t , then $\text{pexp}_{i,t'} = \text{pexp}_{i,t}$. Let t be such a factor. If, for some even j , we have $\text{pexp}_{j+2,t} = \text{pexp}_{j,t} + 3$, then it means that by appending a 2-terms product to a j -terms product that

corresponds to a maximum factor of t^ω , that can only add a cube of period $|t|$. This implies that for every k , $\text{pexp}_{j+2k,t} = \text{pexp}_{j,t} + 3k$.

We have checked by computer that for every conjugacy class of words t of length at most 8, there exists a (small) even j such that $\text{pexp}_{j+2,t} = \text{pexp}_{j,t} + 3$. Thus we have $\text{pexp}_i \leq \frac{3i}{2} + \frac{1}{4}$ in all cases. \square

Proof of $\text{RT}_i(3) \leq \frac{3i}{2} + \frac{1}{6}$ for every odd $i \geq 3$.

Let us show that the image of every $7/5^+$ -free word over Σ_4 by the following 514-uniform morphism satisfies $\text{pexp}_i \leq \frac{3i}{2} + \frac{1}{6}$ for every odd $i \geq 3$.

0 \mapsto 01020120210120102120210201210120102012021020121021201020121012
 02102012102120210120102012102120102012021020121012010212021020
 12102120102012021012010212021020121021202101201020121021201020
 12101202102012102120210120102120210201210120102012021020121012
 01021202102012102120102012101202102012102120102012021012010212
 02102012101201020120210201210212021012010201210120210201210212
 01020120210201210120102120210201210212010201210120210201210212
 02101201021202102012101201020120210201210120102120210201210212
 021012010201210212

1 \mapsto 01020120210120102120210201210120102012021020121021201020121012
 02102012102120102012021020121012010212021020121021201020120210
 12010212021020121021202101201020121021201020121012021020121021
 20210120102120210201210120102012021020121021201020121012021020
 12101201021202102012102120210120102012102120102012021012010212
 02102012101201020120210201210120102120210201210212010201210120
 21020121021202101201021202102012101201020120210201210212021012
 01020121012021020121021201020120210201210120102120210201210212
 021012010201210212

2 \mapsto 01020120210120102120210201210120102012021020121021201020121012
 02102012101201021202102012102120102012021012010212021020121021
 20210120102012102120102012101202102012102120210120102120210201
 21012010201202102012101201021202102012102120102012101202102012
 10212010201202101201021202102012102120210120102012102120102012
 02102012101201021202102012102120102012021012010212021020121012
 01020120210201210212021012010201210212010201210120210201210212
 02101201021202102012101201020120210201210120102120210201210212
 021012010201210212

3 \mapsto 01020120210120102120210201210120102012021020121021201020121012
02102012101201021202102012102120102012021012010212021020121021
20210120102012101202102012102120102012021020121012010212021020
12102120102012101202102012102120210120102012102120102012021012
01021202102012101201020120210201210212010201210120210201210212
01020120210201210120102120210201210212021012010201210212010201
20210120102120210201210212010201210120210201210212021012010212
02102012101201020120210201210120102120210201210212021012010201
210120210201210212

First, we check that such ternary images are $\left(\frac{3^+}{2}, 45\right)$ -free using the method in [6]. By Lemma 2.1 in [6], it is sufficient to check this freeness property for the image of every $7/5^+$ -free word over Σ_4 of length smaller than $\frac{2 \times \frac{3}{2}}{\frac{3}{2} - \frac{1}{5}} = 30$. Thus, the period of every repetition formed from i pieces and with exponent strictly greater than $\frac{3i}{2}$ must be at most 44. Using the same argument as in the previous proof, we have checked by computer that for every conjugacy class of words t of length at most 44, there exists a (small) odd j such that $\text{pexp}_{j+2,t} = \text{pexp}_{j,t} + 3$. Thus we have $\text{pexp}_i \leq \frac{3i}{2} + \frac{1}{6}$ in all cases. \square

3 Concluding remarks

The next step would be to consider the 4-letter alphabet. Obviously, $\text{RT}_{i+1}(k) \geq \text{RT}_i(k) + 1$ for every $i \geq 1$ and $k \geq 2$. Mousavi and Shallit [5] verified that $\text{RT}_2(4) \geq \frac{5}{2}$, so that $\text{RT}_i(4) \geq i + \frac{1}{2}$ for every $i \geq 2$. We conjecture that this is best possible, i.e., that $\text{RT}_i(4) = i + \frac{1}{2}$ for every $i \geq 2$. However, a proof of an upper bound of the form $\text{RT}_i(4) \leq i + c$ cannot be similar to the proof of the upper bounds of Theorem 2. The multiplicative factor of i , which drops from $\frac{3}{2}$ when $k = 3$ to 1 when $k = 4$, forbids that the constructed word is the morphic image of any (unspecified) Dejean word over a given alphabet.

References

- [1] A. Carpi. On Dejean’s conjecture over large alphabets. *Theoret. Comput. Sci.*, 385(1–3):137–151, 2007.

- [2] F. Dejean. Sur un théorème de Thue, *J. Combin. Theory. Ser. A*, 13:90–99, 1972.
- [3] Pytheas Fogg. Substitutions in Dynamics, Arithmetics and Combinatorics. Springer Science & Business Media, 2002.
- [4] J. Moulin Ollagnier. Proof of Dejean’s conjecture for alphabets with 5, 6, 7, 8, 9, 10, and 11 letters. *Theoret. Comput. Sci.*, 95(2):187–205, 1992.
- [5] H. Mousavi and J. Shallit. Repetition avoidance in circular factors. *Developments in Language Theory 2013*, 384–395.
- [6] P. Ochem. A generator of morphisms for infinite words. *RAIRO - Theor. Inform. Appl.*, 40:427–441, 2006.
- [7] J.-J. Pansiot. A propos d’une conjecture de F. Dejean sur les répétitions dans les mots. *Discrete Appl. Math.*, 7(3):297–311, 1984.
- [8] M. Rao. Last cases of Dejean’s conjecture. *Theoret. Comput. Sci.*, 412(27):3010–3018, 2011.
- [9] A. Thue. Über unendliche Zeichenreihen. *Norske Vid. Selsk. Skr. I. Mat. Nat. Kl. Christiania*, 7:1–22, 1906.