



HAL
open science

The chromatic number and switching chromatic number of 2-edge-colored graphs of bounded degree

Fabien Jacques, Mickaël Montassier, Alexandre Pinlou

► **To cite this version:**

Fabien Jacques, Mickaël Montassier, Alexandre Pinlou. The chromatic number and switching chromatic number of 2-edge-colored graphs of bounded degree. BGW 2019 - 5th Bordeaux Graph Workshop, Oct 2019, Bordeaux, France. lirmm-02938635

HAL Id: lirmm-02938635

<https://hal-lirmm.ccsd.cnrs.fr/lirmm-02938635v1>

Submitted on 3 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The chromatic number and switching chromatic number of 2-edge-colored graphs of bounded degree

Fabien Jacques, Mickael Montassier, and Alexandre Pinlou

LIRMM, Université de Montpellier, CNRS, France

EXTENDED ABSTRACT

The notion of homomorphisms of 2-edge-colored graphs has already been studied as a way of extending classical results in graph coloring such as Hadwiger's conjecture. Guenin [5] introduced the notion of switching homomorphisms for its relation with a well-known conjecture of Seymour. In 2012, this notion has been further developed by Naserasr et al. [6] as it captures a number of well-known conjectures that can be reformulated using the definition of switching homomorphisms. In this extended abstract, we study homomorphisms of 2-edge colored graphs and switching homomorphisms of bounded degree graphs.

A *2-edge-colored graph* $G = (V, E, s)$ is a simple graph (V, E) with two kinds of edges: positive and negative edges. The signature $s : E(G) \rightarrow \{-1, +1\}$ assigns to each edge its sign. In the sequel, \mathcal{D}_k (resp. \mathcal{D}_k^c) denotes the class of 2-edge-colored graphs (resp. connected 2-edge-colored graphs) with maximum degree k .

Given two 2-edge-colored graphs G and H , the mapping $\varphi : V(G) \rightarrow V(H)$ is a *homomorphism* if φ maps every edge of G to an edge of H with the same sign. This can be seen as coloring the graph G by using the vertices of H as colors. The target graph H gives us the rules that this coloring must obey. If vertices 1 and 2 in H are connected with a positive (resp. negative) edge, then every pair of adjacent vertices in G colored with 1 and 2 must be connected with a positive (resp. negative) edge. The *chromatic number* $\chi_2(G)$ of a 2-edge-colored graph G is the order of a smallest 2-edge-colored graph H such that G admits a homomorphism to H . The chromatic number $\chi_2(\mathcal{C})$ of a class of 2-edge-colored graphs \mathcal{C} is the maximum of the chromatic numbers of the graphs in the class. This number can be infinite.

2-edge-colored graphs are, in some sense, similar to oriented graphs since a pair of vertices can be adjacent in two different ways in both kinds of graphs: with a positive or a negative edge in the case of 2-edge-colored graphs, with a toward or a backward arc in the oriented case.

The notion of homomorphism of oriented graphs has been introduced by Courcell [3] in 1994 and has been widely studied since then. Due to the similarity above-mentioned, we try to adapt techniques used to study the homomorphisms of oriented graphs of bounded degree to 2-edge-colored graphs of bounded degree. We also study switching homomorphisms of 2-edge-colored graphs in order to obtain results on signed graphs.

Switching a vertex v of a 2-edge-colored graph corresponds to reversing the signs of all edges incident to v .

Two 2-edge-colored graphs G and G' are *switching equivalent* if it is possible to turn G into G' after some number of switches. We call the classes created by this equivalence relation *switching classes* (note that switching classes are equivalent to the notion of signed graphs).

Given two 2-edge-colored graphs G and H , the mapping $\varphi : V(G) \rightarrow V(H)$ is a *switching homomorphism* if there is a graph G' switching equivalent to G such that φ maps every edge of G' to an edge of H with the same sign. The *switching chromatic number* $\chi_s(G)$ of a 2-edge-colored graph G is the order of a smallest 2-edge-colored graph H such that G admits a switching homomorphism to H .

Table 1 summarizes results on the chromatic number and switching chromatic number of the classes of (connected) 2-edge-colored graphs of bounded degree.

The first two lines of Table 1 are more or less folklore. Let us explain in the following the difference that exists between the connected case and the non-connected case for the

| | $\chi_2(\mathcal{D}_k)$ | $\chi_2(\mathcal{D}_k^c)$ | $\chi_s(\mathcal{D}_k)$ | $\chi_s(\mathcal{D}_k^c)$ |
|------------|---|-------------------------------|--|-----------------------------------|
| $k = 1$ | $\chi_2(\mathcal{D}_k) = 3$ | $\chi_2(\mathcal{D}_k^c) = 2$ | $\chi_s(\mathcal{D}_k) = \chi_s(\mathcal{D}_k^c) = 2$ | |
| $k = 2$ | $\chi_2(\mathcal{D}_k) = 6$ | $\chi_2(\mathcal{D}_k^c) = 5$ | $\chi_s(\mathcal{D}_k) = \chi_s(\mathcal{D}_k^c) = 4$ | |
| $k = 3$ | $8 \leq \chi_2(\mathcal{D}_k^c) \leq \chi_2(\mathcal{D}_k) \leq 11$ | | $6 \leq \chi_s(\mathcal{D}_k) \leq 7$ [1] | $\chi_s(\mathcal{D}_k^c) = 6$ [1] |
| $k = 4$ | $12 \leq \chi_2(\mathcal{D}_k^c) \leq \chi_2(\mathcal{D}_k) \leq 31$ | | $10 \leq \chi_s(\mathcal{D}_k^c) \leq \chi_s(\mathcal{D}_k) \leq 16$ | |
| $k \geq 5$ | $2^{\frac{k}{2}} \leq \chi_2(\mathcal{D}_k^c) \leq \chi_2(\mathcal{D}_k) \leq 2^{k+1}(k-1)^2$ [4] | | $\chi_s(\mathcal{D}_k^c) \leq \chi_s(\mathcal{D}_k) \leq 2^{k+1}(k-1)^2$ [4] | |

Table 1: Results on the chromatic number and switching chromatic number of the classes of (connected) 2-edge-colored graphs of bounded degree.

chromatic number of 2-edge-colored graphs with maximum degree 1 or 2. An edge of a 2-edge-colored graph has chromatic number 2 and thus $\chi_2(\mathcal{D}_1^c) = 2$; however, a 2-edge-colored graph with two non-adjacent edges, one positive and one negative, has chromatic number 3 (the target graph needs a positive and a negative edge, hence at least three vertices) and thus $\chi_2(\mathcal{D}_1) = 3$. We therefore have a difference between the chromatic numbers of connected and non-connected 2-edge-colored graphs with maximum degree 1. This difference does not exist for switching homomorphisms since a negative edge can be changed into a positive one after a switch. This difference (and lack thereof for switching homomorphisms) appears also in graphs with maximum degree 2. We have $\chi_2(\mathcal{D}_2) \geq 6$ since there is no 2-edge-colored graph on 5 vertices that can color all the four graphs depicted in Figure 1. However, every connected 2-edge-colored graph with maximum degree 2 admits a homomorphism to one of the two graphs on 5 vertices depicted in Figure 2 and thus $\chi_2(\mathcal{D}_2^c) \leq 5$. In order to color any graph of \mathcal{D}_2 , we need a target graph that contains both graphs depicted in Figure 2 as subgraphs. This is possible with 6 vertices so $\chi_2(\mathcal{D}_2) = 6$. We do not know yet if this is also the case for graphs with maximum degree at least 3.




Figure 1: Four examples of 2-edge-colored graphs with chromatic number 5.




Figure 2: Target graphs for connected 2-edge-colored graphs of maximum degree 2.

The last three lines of Table 1 are dedicated to graphs with maximum degree at least 3. Our main results are the following:

Theorem 1 *We have:*

- $8 \leq \chi_2(\mathcal{D}_3) \leq 11$,
- $12 \leq \chi_2(\mathcal{D}_4) \leq 31$,
- $10 \leq \chi_s(\mathcal{D}_4) \leq 16$.

In order to find an upper bound for a class of graphs, we need to find a target graph that can color every graph in the class. In the case of oriented homomorphisms, oriented graphs that are antiautomorphic, K_n -transitive for some n , or that have Property $P_{n,k}$ for some n and k are good candidates. We analogously define these properties in term of 2-edge-colored graphs.

A 2-edge-colored graph (V, E, s) is said to be *antiautomorphic* if it is isomorphic to $(V, E, -s)$.

A 2-edge-colored graph $G = (V, E, s)$ is said to be *K_n -transitive* if for every pair of cliques $\{u_1, u_2, \dots, u_n\}$ and $\{v_1, v_2, \dots, v_n\}$ in G such that for all $i \neq j$, $s(u_i u_j) = s(v_i v_j)$ there exists an automorphism that maps u_i to v_i for all i . For $n = 1, 2$, or 3 , we say that the graph is *vertex*, *edge*, or *triangle-transitive*, respectively.

A 2-edge-colored graph G has *Property $P_{k,n}$* if for every sequence of k distinct vertices (v_1, v_2, \dots, v_k) that induces a clique in G and for every sign vector $(\alpha_1, \alpha_2, \dots, \alpha_k) \in \{-1, +1\}^k$ there exist at least n distinct vertices $\{u_1, u_2, \dots, u_n\}$ such that $s(v_i u_j) = \alpha_i$ for $1 \leq i \leq k$ and $1 \leq j \leq n$.

Given an integer $q \equiv 1 \pmod{4}$, we consider the family of complete signed Paley graphs SP_q built from the field of order q which has the above-mentioned properties. The vertices of SP_q are the elements of the field of order q and $s(uv) = +1$ if $u - v$ is a square and $s(uv) = -1$ otherwise.

Lemma 2 ([7]) *Graph SP_q is vertex-transitive, edge-transitive, antiautomorphic, and has properties $P_{1, \frac{q-1}{2}}$ and $P_{2, \frac{q-5}{4}}$.*

Let us consider the following operation. Given a 2-edge-colored graph G , we create the *antitwinned graph* of G denoted by $\rho(G)$ as follows. Let G^{+1}, G^{-1} be two copies of G . The vertex corresponding to $v \in V(G)$ in G^i is denoted by v_i , $V(\rho(G)) = V(G^+) \cup V(G^-)$, $E(\rho(G)) = \{u_i v_j : uv \in E(G), i, j \in \{-1, +1\}\}$ and $s_{\rho(G)}(u_i v_j) = i \times j \times s_G(u, v)$.

Lemma 3 ([2]) *Let G and H be two 2-edge-colored graphs. The graph G admits a homomorphism to $\rho(H)$ if and only if it admits a switching homomorphism to H .*

In other words, if a 2-edge-colored graph admits a homomorphism to an antitwinned target graph on n vertices, then it also admits a switching homomorphism to a target graph on $\frac{n}{2}$ vertices. The family $\rho(SP_q)$ also are interesting target graphs (especially for bounding the switching chromatic number since they are antitwinned).

Lemma 4 ([7]) *The graph $\rho(SP_q)$ is vertex-transitive, antiautomorphic, and has properties $P_{1, q-1}$, $P_{2, \frac{q-3}{2}}$, and $P_{3, \max(\frac{q-9}{4}, 0)}$.*

One last family of interesting target graphs are the Tromp-Paley graphs (this construction due to Tromp (unpublished) has been widely used in the case of oriented homomorphisms). Let SP_q^+ be SP_q with an additional vertex that is connected to every other vertex with a positive edge. The Tromp-Paley graph $TR(SP_q)$ corresponds to $\rho(SP_q^+)$. This construction improves the properties of $\rho(SP_q)$ at the cost of having two more vertices. Since Tromp-Paley graphs are antitwinned, they are interesting for bounding the switching chromatic number.

Lemma 5 ([7]) *$TR(SP_q)$ is vertex-transitive, edge-transitive, antiautomorphic, and has properties $P_{1, q}$, $P_{2, \frac{q-1}{2}}$, and $P_{3, \frac{q-5}{4}}$.*

Bensmail et al. [1] recently proved that every 2-edge-colored graph with maximum degree 3 except the all positive and all negative K_4 admits a homomorphism to $TR(SP_5)$, hence $\chi_2(\mathcal{D}_3^c) \leq 12$, and $\chi_s(\mathcal{D}_3^c) \leq 6$ by Lemma 3. In the non-connected case, we can easily get $\chi_2(\mathcal{D}_3) \leq 14$ and thus $\chi_s(\mathcal{D}_3) \leq 7$ by Lemma 3 (it is possible to create an all positive K_4 and an all negative K_4 in $TR(SP_5)$ by adding two vertices). Their proof uses a computer to show that a minimal counter-example cannot contain some configurations and then concludes by using the properties of $TR(SP_5)$. Theorem 1 improves the upper bound of 14 to 11.

Let us give a sketch of proof of the first result of Theorem 1, namely $\chi_2(\mathcal{D}_3) \leq 11$.

Consider the graph SP_9^* obtained from SP_9 by adding two new vertices $0'$ and $1'$ as follows. Take the two vertices 0 and 1 of SP_9 (note that $s(01) = +1$), and link $0'$ and $1'$ to the vertices of SP_9 in the same way as 0 and 1 are, respectively; add an edge $0'1'$ with $s(0'1') = -1$; finally we add edges $00'$ and $11'$ with $s(00') = -1$ and $s(11') = +1$. We will prove that every graph from \mathcal{D}_3 admits a homomorphism to SP_9^* .

We first show that every connected 2-degenerate 2-edge-colored graph with maximum degree 3 admits a homomorphism to SP_9 by using its structural properties given by Lemma 2 (a unique exception exists and is separately treated).

Let G be a connected 3-regular 2-edge-colored graph. If G is all positive, then we color it using an all positive K_4 that SP_9^* contains as a subgraph. Assume now that G is not all positive. Let uv be a negative edge of G . We remove uv from G to create a new graph G' . Graph G' is 2-degenerate so it admits a homomorphism φ' to SP_9 . If $s(\varphi'(u)\varphi'(v)) = -1$, then φ' is also a homomorphism from G to SP_9 .

If $s(\varphi'(u)\varphi'(v)) = +1$, then by edge-transitivity of SP_9 we can recolor the vertices of G' such that $\varphi'(u) = 0$ and $\varphi'(v) = 1$. We can then extend φ' to a homomorphism φ of G to SP_9^* by recoloring u and v such that $\varphi(u) = 0'$ and $\varphi(v) = 1'$ since $s(0'1') = -1$.

Finally, if $\varphi'(u) = \varphi'(v)$, then by vertex-transitivity of SP_9 we can recolor the vertices of G' such that $\varphi'(u) = \varphi'(v) = 0$. We can then extend φ' to a homomorphism φ of G to SP_9^* by recoloring v such that $\varphi(v) = 0'$ since $s(00') = -1$.

We have proven that every graph in \mathcal{D}_3^c admits a homomorphism to SP_9^* which means that SP_9^* is universal for \mathcal{D}_3 . This concludes the proof.

To prove the two other upper bounds of Theorem 1, we use the same method on target graphs SP_{29} and $TR(SP_{13})$.




Figure 3: The graph SP_9 , non-edges are negative edges

References

- [1] J. Bensmail, T. Pierron, É. Sopena. Personal communication, 2019.
- [2] R. C. Brewster, T. Graves. Edge-switching homomorphisms of edge-coloured graphs. In **Discrete Mathematics** 309(18):5540-5546, 2009.
- [3] B. Courcelle. The monadic second order-logic of graphs VI : on several representations of graphs by relational structures. In **Discrete Applied Mathematics** 54:117-149, 1994.
- [4] S. Das, S. Nandi, S. Sen. On chromatic number of colored mixed graphs. In **Conf. on Algorithms and Discrete Applied Mathematics** 130-140, 2017.
- [5] B. Guenin. Packing odd circuit covers: A conjecture. Manuscript, 2005.
- [6] R. Naserasr, E. Rollova, and É. Sopena. Homomorphisms of signed graphs. In **Journal of Graph Theory** 79(3):178-212, 2015.
- [7] P. Ochem, A. Pinlou, S. Sen. Homomorphisms of 2-edge-colored triangle-free planar graphs. In **Journal of Graph Theory** 85(1):258-277, 2017.