



HAL
open science

r-hued coloring of planar graphs with girth at least 8

Hoang La, Mickaël Montassier, Alexandre Pinlou, Petru Valicov

► **To cite this version:**

Hoang La, Mickaël Montassier, Alexandre Pinlou, Petru Valicov. r-hued coloring of planar graphs with girth at least 8. BGW 2019 - 5th Bordeaux Graph Workshop, Oct 2019, Bordeaux, France. lirmm-02938645

HAL Id: lirmm-02938645

<https://hal-lirmm.ccsd.cnrs.fr/lirmm-02938645v1>

Submitted on 29 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

r -hued coloring of planar graphs with girth at least 8

Hoang La, Mickael Montassier, Alexandre Pinlou, and Petru Valicov

LIRMM, University of Montpellier, CNRS, France

EXTENDED ABSTRACT

In this paper, we consider simple undirected graphs without loops. A *proper k -coloring* of the vertices of a graph $G = (V, E)$ is an assignment of colors from 1 to k such that no two adjacent vertices have the same color. The *chromatic number* of G , denoted $\chi(G)$, is the smallest integer k so that G has a proper k -coloring. In 1969, Kramer and Kramer introduced the notion of *2-distance k -coloring* [23] which is a proper k -coloring such that no pair of vertices at distance 2 have the same color. The *2-distance chromatic number* of G , denoted $\chi^2(G)$, is the smallest integer k so that G has a 2-distance k -coloring. An example of 2-distance coloring is given in Figure 1a.




Figure 1: A graph G with $\chi^2(G) = 6$ and $\chi(G) = 3$

Note that for any graph G with maximum degree Δ , $\Delta + 1 \leq \chi^2(G) \leq \Delta^2 + 1$. The lower bound is trivial since, in a 2-distance coloring, every neighbor of a vertex v with degree Δ , and v itself must have a different color. As for the upper bound, a greedy algorithm shows that $\chi^2(G) \leq \Delta(G)^2 + 1$. Moreover, this bound is tight for some graphs, for example, Moore graphs of type $(\Delta, 2)$, which are graphs where all vertices have degree Δ , are at distance at most two from each other, and the total number of vertices is $\Delta^2 + 1$. The Moore graphs of type $(3, 2)$ and of type $(7, 2)$ are the Petersen graph and the Hoffman-Singleton graph respectively.

In this paper, we focus on *planar graphs* which are graphs that can be drawn in the plane without crossing the edges. One motivation to study this class of graphs is the following famous conjecture stating an upper bound which is linear in Δ :

Conjecture 1 (Wegner, 1977 [31]) *Let G be a planar graph with maximum degree Δ . Then,*

$$\chi^2(G) \leq \begin{cases} 7, & \text{if } \Delta \leq 3, \\ \Delta + 5, & \text{if } 4 \leq \Delta \leq 7, \\ \lfloor \frac{3\Delta}{2} \rfloor + 1, & \text{if } \Delta \geq 8. \end{cases}$$

Wegner showed that the upper bounds of the conjecture are tight. For instance when $\Delta \geq 8$ consider the Wegner graph obtained as follows : take a triangle xyz , multiply each edge $\frac{3}{2}\Delta$ times, subdivide them once, then add an edge between x and y . The case $\Delta \leq 3$ of Conjecture 1 was proved independently by Thomassen [29] and by Hartke *et al.* [17]. For $\Delta \geq 8$, Havet *et al.* [18] proved that the bound is $\frac{3}{2}\Delta(1 + o(1))$, where $o(1)$ is as $\Delta \rightarrow \infty$.

The coefficient before Δ becomes 1 when the *girth*, the length of a shortest cycle, of the graph increases. Extensive researches have been done in this case, and many results have taken the following form: *every planar graph G of girth $g \geq g_0$ and $\Delta(G) \geq \Delta_0$ satisfies $\chi^2(G) \leq \Delta + c(g_0, \Delta_0)$, where $c(g_0, \Delta_0)$ is a constant depending only on g_0 and Δ_0 .* Table 1 shows all known such results on the 2-distance chromatic number of planar graphs with fixed girth, up to our own knowledge.

$g_0 \setminus \chi^2(G)$	$\Delta + 1$	$\Delta + 2$	$\Delta + 3$	$\Delta + 4$	$\Delta + 5$	$\Delta + 6$	$\Delta + 7$	$\Delta + 8$
3	X			$\Delta = 3$ [29, 17]				
4	X							
5	X	$\Delta \geq 10$ [1]	$\Delta \geq 339$ [15]	$\Delta \geq 312$ [14]	$\Delta \geq 15$ [8]	$\Delta \geq 12$ [7]	$\Delta \neq 7, 8$ [14]	all Δ [30]
6	X	$\Delta \geq 17$ [4]	$\Delta \geq 9$ [7]		all Δ [10]			
7	$\Delta \geq 16$ [19]			$\Delta = 4$ [12]				
8	$\Delta \geq 10$ [19] $\Delta \geq 9$ (Corollary 4)		$\Delta = 5$ [6]					
9	$\Delta \geq 8$ [3]	$\Delta = 5$ [6]	$\Delta = 3$ [13]					
10	$\Delta \geq 6$ [19]							
11		$\Delta = 4$ [12]						
12	$\Delta = 5$ [19]	$\Delta = 3$ [5]						
13								
14	$\Delta \geq 4$ [2]							
...								
22	$\Delta = 3$ [19]							

Table 1: The latest results with a coefficient 1 before Δ in the upper bound of χ^2 .

For example, the result from line "7" and column " $\Delta + 1$ " from Table 1 reads as follows : "every planar graph G of girth at least 7 and of Δ at least 16 satisfies $\chi^2(G) \leq \Delta + 1$ ". The crossed out cases in the first column correspond to the fact that, for $g_0 \leq 6$, there are planar graphs G with $\chi^2(G) = \Delta + 2$ for arbitrarily large Δ [16]. The lack of results for $g \geq 4$ is due to the fact that the Wegner graph without xy has girth 4, and $\chi^2 = \lfloor \frac{3\Delta}{2} \rfloor - 1$ for all Δ .

The "2-distance" condition in 2-distance colorings requires that vertices at distance at most two have different colors. In other words, all neighbors of the same vertex must have different colors. Recently, this condition was generalized and the notion of r -hued coloring was introduced [26]. Let $r, k \geq 1$ be two integers. An r -hued k -coloring of the vertices of G is a proper k -coloring of the vertices, such that all vertices are r -hued. A vertex is r -hued if the number of colors in its neighborhood $N_G(v) = \{x| xv \in E\}$ is at least $\min\{d_G(v), r\}$. The r -hued chromatic number of G , denoted $\chi_r(G)$, is the smallest integer k so that G has an r -hued k -coloring. It is indeed a generalization of 2-distance colorings which correspond to the case $r \geq \Delta$, as all vertices in the same neighborhood will have different colors. More generally, its link to proper coloring and 2-distance coloring resides in the following equation:

$$\chi(G) = \chi_1(G) \leq \chi_2(G) \leq \dots \leq \chi_\Delta(G) = \chi_{\Delta+1}(G) = \dots = \chi^2(G) \quad (1)$$

Examples of r -hued colorings are given in Figure 1b and Figure 1c.

Similar to the 2-distance chromatic number, the r -hued chromatic number is linear in r when it comes to planar graphs. In 2014, Song *et al.* proposed a generalization of Conjecture 1:

Conjecture 2 (Song *et al.*, 2014 [27]) *Let G be a planar graph. Then,*

$$\chi_r(G) \leq \begin{cases} r + 3, & \text{if } 1 \leq r \leq 2, \\ r + 5, & \text{if } 3 \leq r \leq 7, \\ \lfloor \frac{3r}{2} \rfloor + 1, & \text{if } r \geq 8. \end{cases}$$

Note that Conjecture 2 implies Conjecture 1 except for the case $r = 3$. Moreover, the only extremal known examples reaching the upper bounds of Conjecture 2 are the same as for Conjecture 1. It is less clear what would be the expected upper bound when $r < \Delta$. In 2018, Song and Lai [28] proved that, if $r \geq 8$, then every planar graph G verifies $\chi_r(G) \leq 2r + 16$. Similar to 2-distance coloring, the coefficient before r in this upper bound becomes 1 for graphs with a higher girth. Table 2 shows all known results of the following form: *let r and r_0 be integers such that $r \geq r_0$, all planar graph G of girth $g(G) \geq g_0$ satisfies $\chi_r(G) \leq r + c(g_0, r_0)$, where $c(g_0, r_0)$ is a constant depending only on g_0 and r_0 .* The result from the "9" line and " $r + 1$ " column reads "for $r \geq 8$, all planar graph G of girth at least 9 satisfies $\chi_r(G) \leq r + 1$ ".

Since $r + 1$ is a trivial lower bound for χ_r , we study the class of planar graphs verifying $\chi_r = r + 1$ and show the following:

$g_0 \setminus \chi_r(G)$	$r+1$	$r+2$	$r+3$	$r+4$	$r+5$	$r+6$	$r+7$...	$r+10$
3		$r = 2[20]^1$	$r = 2[20]$	$r = 2[22]$			$r = 3[25]$		
4									
5					$r \geq 15[8]$				all $r[8]$
6					$r \geq 3[24]$				
7		$r = 2[22]$		$r = 3[21]$					
8	$r \geq 9$ (Theorem 3)								
9	$r \geq 8[9]$		$r = 3[21]$						
10	$r \geq 6[9]$								
11									
12	$r \geq 5[9]$								
13									
14		$r = 3[11]$							

Table 2: The latest results with a coefficient 1 before r in the upper bound of χ_r .

Theorem 3 *If G is a planar graph with $g(G) \geq 8$, then $\chi_r(G) = r + 1$ for $r \geq 9$.*

Our proof uses the discharging method and exploits planarity arguments.

For $r \geq \Delta$, Theorem 3 gives an improvement of a result on 2-distance coloring published in [19] (see Table 1):

Corollary 4 *If G is a planar graph with $g(G) \geq 8$ and $\Delta(G) \geq 9$, then $\chi^2(G) = \Delta(G) + 1$.*

References

- [1] M. Bonamy, D. Cranston, L. Postle. Planar graphs of girth at least five are square $(\Delta + 2)$ -choosable. In **J. Comb. Theory, Series B**. 134:218–238, 2019.
- [2] M. Bonamy, B. Lévêque, A. Pinlou. 2-distance coloring of sparse graphs. In **Discrete Math.**. 38:155–160, 2011.
- [3] M. Bonamy, B. Lévêque, A. Pinlou. 2-distance coloring of sparse graphs. In **J. Graph Theory**. 77(3), 2014.
- [4] M. Bonamy, B. Lévêque, A. Pinlou. Graphs with maximum degree $\Delta \geq 17$ and maximum average degree less than 3 are list 2-distance $(\Delta + 2)$ -colorable. In **Discrete Math.**. 317:19–32, 2014.
- [5] O.V. Borodin, A.O. Ivanova. List 2-facial 5-colorability of plane graphs with girth at least 12. In **Discrete math.**. 312:306–314, 2012.
- [6] Y. Bu, X. Lv, X. Yan. The list 2-distance coloring of a graph with $\Delta(G) = 5$. In **Discrete Math., Algo. and Appl.**. 7(2), 2015.
- [7] Y. Bu, C. Shang. List 2-distance coloring of planar graphs without short cycles. In **Discrete Math., Algo. and Appl.**. 8(1), 2016.
- [8] Y. Bu, J. Zhu. Channel Assignment with r -Dynamic Coloring. 12th International Conference, AAIM 2018, Dallas, TX, USA, December 3-4, 2018, Proceedings. 36–48, 2018.
- [9] Y. Bu, J. Zhu. List r -dynamic coloring of graphs with small maximum average degree. In **Discrete Appl. Math.**. 258:254–263, 2019.
- [10] Y. Bu, X. Zhu. An optimal square coloring of planar graphs. In **J. Comb. Optim.**. 24:580, 2012.
- [11] J. Cheng, H. -J. Lai, K. J. Lorenzen, R. Luo, J. C. Thompson, C. -Q. Zhang. r -hued coloring of sparse graphs. In **Discrete Appl. Math.**. 237:75–81, 2018.

- [12] D. Cranston, R. Erman, R. Skrekovski. Choosability of the square of a planar graph with maximum degree four. In **Australian J. Comb.** 59(1):86–97, 2014.
- [13] D. Cranston, S. Kim. List-coloring the square of a subcubic graph. In **J. Graph Theory**. 1:65–87, 2008.
- [14] W. Dong, W. Lin. On 2-distance coloring of plane graphs with girth 5. In **Discrete Appl. Math.** 217:495–505, 2017.
- [15] W. Dong, B. Xu. 2-distance coloring of planar graphs with girth 5. In **J. Comb. Optim.** 34:1302, 2017.
- [16] Z. Dvořák, D. Král, P. Nejedlý, R. Škrekovski. Coloring squares of planar graphs with girth six. In **European J. Comb.** 29(4):838–849, 2008.
- [17] S. G. Hartke, S. Jahanbekam, B. Thomas. The chromatic number of the square of subcubic planar graphs. arXiv 1604.06404, 2018.
- [18] F. Havet, J. van den Heuvel, C. McDiarmid, B. Reed. List colouring squares of planar graphs. arXiv 0807.3233, 2017.
- [19] A. O. Ivanova. List 2-distance $(\Delta + 1)$ -coloring of planar graphs with girth at least 7. In **J. Appl. Ind. Math.** 5(2):221–230, 2011.
- [20] S. -J. Kim, S. J. Lee, W. -J. Park. Dynamic coloring and list dynamic coloring of planar graphs. In **Discrete Appl. Math.** 161:2207–2212, 2013.
- [21] S. -J. Kim, B. Park. List 3-dynamic coloring of graphs with small maximum average degree. In **Discrete Math.** 341:1406–1418, 2018.
- [22] S. -J. Kim, W. -J. Park. List dynamic coloring of sparse graphs. In **Comb. Optim. and Appl.** 6831:156–162, 2011.
- [23] F. Kramer, H. Kramer. Un problème de coloration des sommets d’un graphe. In **Comptes Rendus Mathématique, Académie des Sciences, Paris**. 268:46–48, 1969.
- [24] H. -J. Lai, H. Song, J. -L. Wu. On r -hued coloring of planar graphs with girth at least 6. In **Discrete Appl. Math.** 198:251–263, 2016.
- [25] S. Loeb, T. Mahoney, B. Reiniger, J. Wise. Dynamic coloring parameters for graphs with given genus. In **Discrete Appl. Math.** 235:129–141, 2018.
- [26] B. Montgomery. Ph.D. dissertation. West Virginia University, 2001.
- [27] Y. Chen, S. -H. Fan, H. -J. Lai, H. Song, L. Sun. On r -hued coloring of K_4 -minor free graphs. In **Discrete Math.** 315-316:47–52, 2014.
- [28] H. -J. Lai, H. Song. Upper bound of r -hued colorings of planar graphs. In **Discrete Appl. Math.** 243:262–369, 2018.
- [29] C. Thomassen. The square of a planar cubic graph is 7-colorable. In **J. Comb. Theory, Series B**. 128:192–218, 2018.
- [30] W. Dong, W. Lin. An improved bound on 2-distance coloring plane graphs with girth 5. In **J. Comb. Optim.** 32(2):645–655, 2016.
- [31] G. Wegner. Graphs with given diameter and a coloring problem. Technical report, University of Dortmund, 1977.