



HAL
open science

Not all planar graphs are in PURE-4-DIR

Daniel Gonçalves

► **To cite this version:**

Daniel Gonçalves. Not all planar graphs are in PURE-4-DIR. Journal of Graph Algorithms and Applications, 2020, 24 (3), pp.293-301. 10.7155/jgaa.00533 . lirmm-03003499


HAL Id: lirmm-03003499

<https://hal-lirmm.ccsd.cnrs.fr/lirmm-03003499v1>

Submitted on 13 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Not all planar graphs are in PURE-4-DIR

*Daniel Gonçalves*¹

¹LIRMM, Univ Montpellier, CNRS, Montpellier, France.

Abstract

We prove that some planar graphs are not intersection graphs of segments if only four slopes are allowed for the segments, and if parallel segments do not intersect. This refutes a conjecture of D. West [14].

Submitted: October 2019	Reviewed: February 2020	Revised: February 2020	Accepted: May 2020
	Final: May 2020	Published: June 2020	
Article type: Concise paper		Communicated by: Martin Nöllenburg	

1 Introduction

The *intersection graphs of segments* class SEG is a widely studied class of graphs. Any graph $G \in \text{SEG}$ admits a representation in the plane, where each vertex $v \in V(G)$ corresponds to a segment \mathbf{v} , and where two segments \mathbf{u} and \mathbf{v} intersect if and only if $uv \in E(G)$. The class SEG can be refined by bounding the number of slopes used in the representation. A graph belongs to k -DIR if it has an intersection representation with segments using at most k different slopes. Actually for any $k \geq 1$ [11],

$$k\text{-DIR} \subsetneq (k+1)\text{-DIR} \subsetneq \text{SEG}.$$

One should be aware that here the set of allowed slopes is not set, and that for $k \geq 4$ the choice of the slopes would lead to different graph classes [6]. Let us also introduce the class PURE- k -DIR of graphs having a *pure* k -DIR representation, that is where parallel segments do not intersect.

The relation between planar graphs and these intersections graphs goes back to the PhD thesis of E.R. Scheinerman [13]. He proved that outerplanar graphs belong to PURE-3-DIR and he conjectured that every planar graph belongs to SEG. This conjecture was confirmed by J. Chalopin and the author [1], and more recently by the author, L. Iseemann, and C. Pennarun [8] using a simpler method.

D. West conjectured a strengthening of Scheinerman's conjecture [14]. He asked whether every planar graph has a PURE-4-DIR representation. In a pure representation parallel segments form a stable set. Thus this conjecture would also strengthen the 4-color theorem. Actually planar graphs with a smaller chromatic number have such property. Indeed for any $k \leq 3$, k -colored planar graphs belong to PURE- k -DIR [4, 5, 7, 9]. We here refute West's conjecture.

Theorem 1 *There exist planar graphs that do not belong to PURE-4-DIR.*

In the following section we study the PURE-4-DIR representations of some planar gadget. We then prove Theorem 1 in Section 3 using signed planar graphs that are not 4-colorable [10] (in the sense of signed graphs).

2 Planar gadget

Consider any set of four slopes $\{s_0, s_1, s_2, s_3\}$, ordered such that drawing four concurrent lines $\ell_0, \ell_1, \ell_2, \ell_3$ with these slopes and going around the intersection point one successively crosses $\ell_0, \ell_1, \ell_2, \ell_3, \ell_0, \ell_1, \ell_2$, and ℓ_3 . As there exists a linear mapping that maps s_0 to the horizontal slope, and s_2 to the vertical slope, we have that a graph has a 4-DIR (resp. a PURE-4-DIR) representation using slopes $\{s_0, s_1, s_2, s_3\}$ if and only if it has a 4-DIR (resp. a PURE-4-DIR) representation using the horizontal s_- , some negative slope s_\setminus , the vertical s_\mid , and some positive slope s_\sloppy . Thus in the following we only consider the slopes s_- , s_\setminus , s_\mid , and s_\sloppy . In the pictures s_\setminus , and s_\sloppy form 45° angles with s_- and s_\mid ,

but all the claims hold for any negative slope s_{\setminus} , and for any positive slope $s_{/}$. Let us now study the PURE-4-DIR representations of the graph H depicted in Figure 1.




Figure 1: The graph H , and two of its PURE-4-DIR representations.

Consider a PURE-4-DIR representation of H where the vertices x and y are represented with segments \mathbf{x} and \mathbf{y} of slopes s_{-} and $s_{|}$. As the considered representation is pure, we have that the segments \mathbf{z}_i with i even have the same slope among s_{\setminus} and $s_{/}$, and that the other slope is the one of all the segments \mathbf{z}_i with i odd. Let $\ell_{\mathbf{x}}$ and $\ell_{\mathbf{y}}$ be the lines supporting \mathbf{x} and \mathbf{y} , respectively. Note that the intersection point of $\ell_{\mathbf{x}}$ and $\ell_{\mathbf{y}}$, \mathbf{p} , belongs to exactly one of \mathbf{x} and \mathbf{y} . Indeed, it cannot belong to both as \mathbf{x} and \mathbf{y} should not intersect. One the other hand, if it belongs to none of them, say that \mathbf{x} and \mathbf{y} are on the left and above \mathbf{p} respectively, then there cannot be segments of slope s_{\setminus} intersecting both of them, a contradiction. We can thus define triangles Δ_{ij} as follows.

Definition 2 Given a PURE-4-DIR representation of H where the vertices x and y are represented with segments \mathbf{x} and \mathbf{y} of slopes s_{-} and $s_{|}$, Δ_{ij} , with $1 \leq i = j - 1 < 8$, is the triangle formed by the segments \mathbf{z}_i , \mathbf{z}_j , and \mathbf{x} (resp. and \mathbf{y}) if the intersection point of $\ell_{\mathbf{x}}$ and $\ell_{\mathbf{y}}$, \mathbf{p} , belongs to \mathbf{x} (resp. \mathbf{y}).

Remark 3 Consider any PURE-4-DIR representation of H where the vertices x and y are represented with segments \mathbf{x} and \mathbf{y} of slopes s_{-} and $s_{|}$, and where Δ_{45} is bounded by \mathbf{z}_4 , \mathbf{z}_5 , and \mathbf{x} . Then, for every vertex z that is adjacent to both x and y , but to none of z_4 and z_5 , we have either that

- Δ_{45} contains $\mathbf{y} \cap \mathbf{z}$, the intersection point of \mathbf{y} and \mathbf{z} , or that
- Δ_{45} does not intersect \mathbf{z} .

The following lemma tells us that both cases occur among \mathbf{z}_1 and \mathbf{z}_8 .

Lemma 4 Consider any PURE-4-DIR representation of H where the vertices x and y are represented with segments \mathbf{x} and \mathbf{y} of slopes s_{-} and $s_{|}$, and where Δ_{45} is bounded by \mathbf{z}_4 , \mathbf{z}_5 , and \mathbf{x} . Then either

- Δ_{45} contains $\mathbf{y} \cap \mathbf{z}_1$ and does not intersect \mathbf{z}_8 , or
- Δ_{45} contains $\mathbf{y} \cap \mathbf{z}_8$ and does not intersect \mathbf{z}_1 .

Proof: Let us consider the triangles Δ_{12} , Δ_{45} , and Δ_{78} , that are all bounded by \mathbf{x} .

Claim 5 *One of Δ_{45} and Δ_{12} (resp. one of Δ_{45} and Δ_{78} , and one of Δ_{12} and Δ_{78}) contains the other.*

This is obvious as otherwise \mathbf{z}_4 or \mathbf{z}_5 would intersect \mathbf{z}_1 or \mathbf{z}_2 (resp. \mathbf{z}_4 or \mathbf{z}_5 would intersect \mathbf{z}_7 or \mathbf{z}_8 , and \mathbf{z}_1 or \mathbf{z}_2 would intersect \mathbf{z}_7 or \mathbf{z}_8).

Claim 6 *Triangles Δ_{12} and Δ_{78} cannot both contain (resp. be contained in) Δ_{45} .*

Towards a contradiction assume that the triangles Δ_{12} and Δ_{78} both contain (resp. are contained in) Δ_{45} . By symmetry of H , assume that $\Delta_{45} \subsetneq \Delta_{78} \subsetneq \Delta_{12}$ (resp. $\Delta_{12} \subsetneq \Delta_{78} \subsetneq \Delta_{45}$). Note that in that case, the segment \mathbf{z}_3 that has to intersect \mathbf{x} , \mathbf{y} , \mathbf{z}_2 , and \mathbf{z}_4 necessarily intersects \mathbf{z}_8 , a contradiction.

By the above two claims, either $\Delta_{12} \subsetneq \Delta_{45} \subsetneq \Delta_{78}$ or $\Delta_{78} \subsetneq \Delta_{45} \subsetneq \Delta_{12}$. By Remark 3, this clearly implies the lemma. \square

3 Proof of Theorem 1

A *signed graph* G is a graph where the edges are signed by a mapping $\sigma : E(G) \rightarrow \{-, +\}$. A signed graph is $2k$ -colorable if there exists a mapping $c : V(G) \rightarrow \{-k, 1-k, \dots, -1, 1, \dots, k-1, k\}$ such that for each edge $uv \in E(G)$, $c(u) \neq \sigma(uv)c(v)$. This definition of signed coloring appeared in [12] and note that it is distinct from the one in [15].

Our construction is based on the construction G of Kardoš and Narboni [10] depicted in Figure 2. This signed plane graph is not 4-colorable, and this contradicts a conjecture of E. Máčajová, A. Raspaud, and M. Škoviera [12]. Note that every non-signed plane graph is 4-colorable in the classical sense. We hence have the following.

Remark 7 *For every classical 4-coloring c , with colors $-2, -1, +1, +2$, of the plane graph induced by the positive (i.e. black) edges of G , one of the negative (i.e. red) edges uv is such that $c(u) = -c(v)$.*

We define the (non-signed) graph G^H from G by replacing each negative (i.e. red) edge uv by a copy of H , identifying the vertices x and y of H with the vertices u and v respectively, and by then triangulating the obtained plane graph without creating a new edge among the original vertices of G (See Figure 2 right).

Towards a contradiction consider a PURE-4-DIR representation of G^H . Consider the 4-coloring c of G^H , provided by the mapping that associates the slopes $s_{|}$, s_{\setminus} , $s_{/}$, and s_{-} , with $-2, -1, +1$, and $+2$ respectively. By Remark 7, let us consider the copy of H such that $c(x) = -c(y)$. We can assume that the segments \mathbf{x} and \mathbf{y} have slopes s_{-} and $s_{|}$, respectively (i.e. we assume that $c(x) = -c(y) = 2$). If the segments \mathbf{x} and \mathbf{y} have a different pair of slopes




Figure 2: Left: The construction G of Kardoš and Narboni [10], where black and red edges are positive and negative, respectively. Right: Replacing each red edge with a copy of H , and triangulating (with dashed edges).

(i.e. if $c(x) = -c(y) = -2$ or if $c(x) = -c(y) = \pm 1$), there exists a linear mapping of the plane that would turn the given PURE-4-DIR representation into a PURE-4-DIR representation with the mentioned slopes for \mathbf{x} and \mathbf{y} . Let us also assume, by symmetry of vertices x and y in H , that Δ_{45} is bounded by \mathbf{x} , \mathbf{z}_4 , and \mathbf{z}_5 .

By Lemma 4, we have some constraints on the locations of \mathbf{z}_1 and \mathbf{z}_8 , and let us assume by symmetry of vertices z_1 and z_8 in H , that Δ_{45} contains the point $y \cap z_1$ and does not intersect the segment z_8 . As G^H is a triangulation, there is a $z_1 z_8$ -path P contained in $N(y) \setminus \{z_2, \dots, z_7\}$, where $N(y)$ is the set of y 's neighbors in G^H (see Figure 3). Moreover by Remark 3, P has a subpath P' whose ends, z' and z'' are common neighbors of x and y , while none of the internal vertices is in $N(x)$, and such that Δ_{45} contains the point $y \cap z'$, and does not intersect the segment z'' .

As \mathbf{z}'' lies outside Δ_{45} , and as none of the internal vertices of P' is in $N(x)$, $N(z_4)$ or $N(z_5)$, all the corresponding segments lie outside Δ_{45} too. In particular, this is the case for the neighbor of z' in P' , w . Note that \mathbf{w} should intersect \mathbf{y} and \mathbf{z}' , so it cannot use the same slope, but it can use slope s_- . In that case \mathbf{w} cannot intersect \mathbf{y} and \mathbf{z}' (see Figure 3), a contradiction. Thus G^H does not admit a PURE-4-DIR representation.

4 Conclusion

It now remains open whether plane graphs belong to PURE- k -DIR, for some constant $k > 4$. Actually it is even open whether every plane graph with




Figure 3: Left: The 4-DIR representation of H with Δ_{45} , y and z' . The red segments are possible positions for w , not intersecting Δ_{45} , and with slopes s_- or s_+ . In none of the cases w intersects y and z' . Right: Neighborhood of y in G^H . The path P corresponds to bold edges and the common neighbors of x and y are z_1, z_8, a and b . So the ends of P' should belong to $\{z_1, z_8, a, b\}$.

n vertices belongs to PURE- $(n - f(n))$ -DIR, for some function f tending to infinity. Another question is whether plane graphs belong to k -DIR, for some $k > 0$. In a non-pure k -DIR representation, parallel segments induce an interval representation. Actually, if every planar graph belongs to k -DIR, for some $k > 0$, we can ask parallel segments to induce just a path forest. Indeed, given a plane graph G , let G' be the graph G where for each vertex v , a leaf v' has been attached. Note that G' is planar, and that its k -DIR representations are such that for each $v \in V(G)$ the segment \mathbf{v} has a point (in $\mathbf{v} \cap \mathbf{v}'$) that does not belong to any other segment \mathbf{u} for $u \in V(G)$. The interval representations where each interval has a private point clearly correspond to path forests. As some plane graphs do not admit a vertex partition into two path forests [2,3] we have that some planar graphs do not admit a 2-DIR representation. An anonymous referee provided us an example, described below, that does not admit a 3-DIR representation. Up to our knowledge it is open to know whether every plane graph belongs to 4-DIR.

Consider the planar graph obtained from K_4 by stacking, for each face (including the external face), a new vertex connected to the three incident vertices. Proceed to another step of vertex stacking in every face. As observed earlier, if planar graphs belong to 3-DIR, then our graph has a 3-DIR representation such that each slope induces a path forest. Consider the initial K_4 and denote its vertices v_1, v_2, v_3, v_4 .

If the corresponding four segments use the three different slopes, say $\mathbf{v}_1, \mathbf{v}_2$ are horizontal, \mathbf{v}_3 is vertical, and \mathbf{v}_4 is diagonal. In that case, note that \mathbf{v}_3 and \mathbf{v}_4 intersect \mathbf{v}_1 and \mathbf{v}_2 on their common part. Consider the vertex w stacked onto v_1, v_3, v_4 . Note that w cannot be horizontal as it should intersect \mathbf{v}_1 and \mathbf{v}_3 but avoid \mathbf{v}_2 . Note that w cannot be vertical neither, as it should intersect

\mathbf{v}_3 and \mathbf{v}_1 but avoid \mathbf{v}_2 . Similarly, \mathbf{w} cannot be diagonal.

If the corresponding four segments \mathbf{v}_1 , \mathbf{v}_2 , \mathbf{v}_3 , and \mathbf{v}_4 use only two slopes, say \mathbf{v}_1 , \mathbf{v}_2 are horizontal, and \mathbf{v}_3 , \mathbf{v}_4 are vertical. Note that necessarily \mathbf{v}_1 , \mathbf{v}_2 , \mathbf{v}_3 , and \mathbf{v}_4 all meet in a common point. Consider again the vertex w stacked onto v_1, v_3, v_4 . the segment \mathbf{w} can neither be horizontal, nor vertical as it avoids \mathbf{v}_2 . But then v_1, v_3, v_4, w form a K_4 with three slopes and the above argument applies.

Acknowledgments

The author would like to thank P. Ochem and A. Pinlou for indicating him the references on signed graphs [10,12]. He would also like to thank the anonymous referee that provided the example presented in the conclusion.

References

- [1] J. Chalopin and D. Gonçalves. Every planar graph is the intersection graph of segments in the plane: extended abstract. In *Proceedings of the forty-first annual ACM symposium on Theory of computing SODA '09*, pages 631–638, 2009. doi:10.1145/1536414.1536500.
- [2] G. Chartrand and H. Kronk. The point-arboricity of planar graphs. *Journal of the London Mathematical Society*, 44(1):612–616, 1969. doi:10.1112/jlms/s1-44.1.612.
- [3] L. Cowen, R. Cowen, and D. Woodall. Defective colorings of graphs in surfaces: Partitions into subgraphs of bounded valency. *Journal of Graph Theory*, 10(2):187–195, 1986. doi:10.1002/jgt.3190100207.
- [4] J. Czyzowicz, E. Kranakis, and J. Urrutia. A simple proof of the representation of bipartite planar graphs as the contact graphs of orthogonal straight line segments. *Inform. Process. Lett.*, 66(3):125–126, 1998. doi:10.1016/S0020-0190(98)00046-5.
- [5] H. de Fraysseix, P. O. de Mendez, and J. Pach. Representation of planar graphs by segments. In *Intuitive geometry (Szeged, 1991)*, *Colloq. Math. Soc. János Bolyai*, volume 63, pages 109–117, 1994.
- [6] J. Černý, D. Král, H. Nyklová, and O. Pangrác. On intersection graphs of segments with prescribed slopes. In *Proc. Graph Drawing '01, LNCS*, volume 2265, pages 261–271, 2002. doi:10.1007/3-540-45848-4_21.
- [7] D. Gonçalves. 3-colorable planar graphs have an intersection segment representation using 3 slopes. In *Proc. of WG 2019, LNCS*, volume 11789, pages 351–363, 2019. doi:10.1007/978-3-030-30786-8_27.
- [8] D. Gonçalves, L. Isenmann, and C. Pennarun. Planar graphs as L-intersection or L-contact graphs. In *Proc. of 29th Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 172–184, 2018. doi:10.1137/1.9781611975031.12.
- [9] I.-A. Hartman, I. Newman, and R. Ziv. On grid intersection graphs. *Discrete Math.*, 87(1):41–52, 1991. doi:10.1016/0012-365X(91)90069-E.
- [10] F. Kardoš and J. Narboni. On the 4-color theorem for signed graphs. *ArXiv*, 1906.05638, 2019.
- [11] J. Kratochvíl and J. Matoušek. Intersection graphs of segments. *Journal of Combinatorial Theory, Series B*, 62(2):289–315, 1994. doi:10.1006/jctb.1994.1071.
- [12] E. Máčajová, A. Raspaud, and M. Škoviera. The chromatic number of a signed graph. *Elec. J. of Combinatorics*, 23(1):1–14, 2016.

- [13] E. Scheinerman. *Intersection classes and multiple intersection parameters of graphs*. PhD thesis, Princeton University, 1984.
- [14] D. West. Open problems. *SIAM J. Discrete Math. Newslett.*, 2(1):10–12, 1991.
- [15] T. Zaslavsky. Signed graph coloring. *Discrete Mathematics*, 39(2):215–228, 1982. doi:10.1016/0012-365X(82)90144-3.