



HAL
open science

Homomorphisms of planar (m, n) -colored-mixed graphs to planar targets

Fabien Jacques, Pascal Ochem

► **To cite this version:**

Fabien Jacques, Pascal Ochem. Homomorphisms of planar (m, n) -colored-mixed graphs to planar targets. *Discrete Mathematics*, 2021, 344 (12), pp.#112600. 10.1016/j.disc.2021.112600. lirmm-03371159

HAL Id: lirmm-03371159

<https://hal-lirmm.ccsd.cnrs.fr/lirmm-03371159>

Submitted on 16 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial | 4.0 International License

Homomorphisms of planar (m, n) -colored-mixed graphs to planar targets

Fabien Jacques and Pascal Ochem*
LIRMM, Université de Montpellier, and CNRS. France

Abstract

An (m, n) -colored-mixed graph $G = (V, A_1, A_2, \dots, A_m, E_1, E_2, \dots, E_n)$ is a graph having m colors of arcs and n colors of edges. We do not allow two arcs or edges to have the same endpoints. A homomorphism from an (m, n) -colored-mixed graph G to another (m, n) -colored-mixed graph H is a morphism $\varphi : V(G) \rightarrow V(H)$ such that each edge (resp. arc) of G is mapped to an edge (resp. arc) of H of the same color (and orientation). An (m, n) -colored-mixed graph T is said to be $P_g^{(m,n)}$ -universal if every graph in $P_g^{(m,n)}$ (the planar (m, n) -colored-mixed graphs with girth at least g) admits a homomorphism to T .

We show that planar $P_g^{(m,n)}$ -universal graphs do not exist for $2m + n \geq 3$ (and any value of g) and find a minimal (in the number vertices) planar $P_g^{(m,n)}$ -universal graphs in the other cases.

1 Introduction

The concept of homomorphisms of (m, n) -colored-mixed graph was introduced by J. Nesěřil and A. Raspaud [1] in order to generalize homomorphisms of k -edge-colored graphs and oriented graphs.

An (m, n) -colored-mixed graph $G = (V, A_1, A_2, \dots, A_m, E_1, E_2, \dots, E_n)$ is a graph having m colors of arcs and n colors of edges. We do not allow two arcs or edges to have the same endpoints and we do not allow loops. The case $m = 0$ and $n = 1$ corresponds to simple graphs, $m = 1$ and $n = 0$ to oriented graphs and $m = 0$ and $n = k$ to k -edge-colored graphs. For the case $m = 0$ and $n = 2$ (2-edge-colored graphs) we refer to the two types of edges as *blue* and *red* edges.

A *homomorphism* from an (m, n) -colored-mixed graph G to another (m, n) -colored-mixed graph H is a mapping $\varphi : V(G) \rightarrow V(H)$ such that every edge (resp. arc) of G is mapped to an edge (resp. arc) of H of the same color (and orientation). If G admits a homomorphism to H , we say that G is H -colorable since this homomorphism can be seen as a coloring of the vertices of G using the vertices of H as colors. The edges and arcs of H (and their colors) give us the rules that this coloring must follow. Given a class of graphs \mathcal{C} , a graph is \mathcal{C} -universal if for every graph $G \in \mathcal{C}$ is H -colorable. The class $P_g^{(m,n)}$ contains every planar (m, n) -colored-mixed graph with girth at least g . Graph \vec{C}_6^2 is the graph with vertex set $\{0, 1, 2, 3, 4, 5\}$ such that uv is an arc if and only if $v - u \equiv 1 \pmod{6}$ or $v - u \equiv 2 \pmod{6}$.

In this paper, we consider some planar $P_g^{(m,n)}$ -universal graphs with few vertices. They are depicted in Figures 1 and 2. The known results about this topic are as follows.

Theorem 1.

1. K_4 is a planar $P_3^{(0,1)}$ -universal graph. This is the four color theorem.
2. K_3 is a planar $P_4^{(0,1)}$ -universal graph. This is Grötzsch's Theorem [2].
3. \vec{C}_6^2 is a planar $P_{16}^{(1,0)}$ -universal graph [3].

*This work is supported by the ANR project HOSIGRA (ANR-17-CE40-0022)

Our first result shows that, in addition to the case of $(0,1)$ -graphs covered by Theorems 1.1 and 1.2, our topic is actually restricted to the cases of oriented graphs (i.e., $(m,n) = (1,0)$) and 2-edge-colored graphs (i.e., $(m,n) = (0,2)$).

Theorem 2. *For every $g \geq 3$, there exists no planar $P_g^{(m,n)}$ -universal graph if $2m + n \geq 3$.*

As Theorems 1.1 and 1.2 show for $(0,1)$ -graphs, there might exist a trade-off between minimizing the girth g and the number of vertices of the universal graph, for a fixed pair (m,n) . For oriented graphs, Theorem 1.3 tries to minimize the girth. For oriented graphs and 2-edge-colored graphs, we choose instead to minimize the number of vertices of the universal graph.

Theorem 3.

1. \vec{T}_5 is a planar $P_{28}^{(1,0)}$ -universal graph on 5 vertices.
2. T_6 is a planar $P_{22}^{(0,2)}$ -universal graph on 6 vertices.

The following results shows that Theorem 3 is optimal in terms of the number of vertices of the universal graph.

Theorem 4.

1. For every $g \geq 3$, there exists an oriented bipartite cactus graph (i.e., K_4^- minor-free graph) with girth at least g and oriented chromatic number at least 5.
2. For every $g \geq 3$, there exists a 2-edge-colored bipartite outerplanar graph (i.e., $(K_4^-, K_{2,3})$ minor-free graph) with girth at least g that does not map to a planar graph with at most 5 vertices.

Most probably, Theorem 3 is not optimal in terms of girth. The following constructions give lower bounds on the girth.

Theorem 5.

1. There exists an oriented bipartite 2-outerplanar graph with girth 14 that does not map to \vec{T}_5 .
2. There exists a 2-edge-colored planar graph with girth 11 that does not map to T_6 .
3. There exists a 2-edge-colored bipartite planar graph with girth 10 that does not map to T_6 .




Figure 1: The $P_{28}^{(1,0)}$ -universal graph \vec{T}_5 .




Figure 2: The $P_{22}^{(0,2)}$ -universal graph T_6 .

Next, we obtain the following complexity dichotomies:

Theorem 6.

1. For any fixed girth $g \geq 3$, either every graph in $P_g^{(1,0)}$ maps to \vec{T}_5 or it is NP-complete to decide whether a graph in $P_g^{(1,0)}$ maps to \vec{T}_5 . Either every bipartite graph in $P_g^{(1,0)}$ maps to \vec{T}_5 or it is NP-complete to decide whether a bipartite graph in $P_g^{(1,0)}$ maps to \vec{T}_5 .
2. Either every graph in $P_g^{(0,2)}$ maps to T_6 or it is NP-complete to decide whether a graph in $P_g^{(1,0)}$ maps to T_6 . Either every bipartite graph in $P_g^{(0,2)}$ maps to T_6 or it is NP-complete to decide whether a bipartite graph in $P_g^{(1,0)}$ maps to T_6 .

Finally, we can use Theorem 6 with the non-colorable graphs in Theorem 5.

Corollary 7.

1. Deciding whether a bipartite graph in $P_{14}^{(1,0)}$ maps to \vec{T}_5 is NP-complete.
2. Deciding whether a graph in $P_{11}^{(0,2)}$ maps to T_6 is NP-complete.
3. Deciding whether a bipartite graph in $P_{10}^{(0,2)}$ maps to T_6 is NP-complete.

A 2-edge-colored path or cycle is said to be *alternating* if any two adjacent edges have distinct colors.

Proposition 8 (folklore).

- Every planar simple graph on n vertices has at most $3n - 6$ edges.
- Every planar simple graph satisfies $(\text{mad}(G) - 2) \cdot (g(G) - 2) < 4$.

2 Proof of Theorem 3

We use the discharging method for both results in Theorem 3. The following lemma will handle the discharging part. We call a vertex of degree n an n -vertex and a vertex of degree at least n an n^+ -vertex. If there is a path made only of 2-vertices linking two vertices u and v , we say that v is a weak-neighbor of u . If v is a neighbor of u , we also say that v is a weak-neighbor of u . We call a (weak-)neighbor of degree n an n -(weak-)neighbor.

Lemma 9. *Let k be a non-negative integer. Let G be a graph with minimum degree 2 such that every 3-vertex has at most k 2-weak-neighbors and every path contains at most $\frac{k+1}{2}$ consecutive 2-vertices. Then $\text{mad}(G) \geq 2 + \frac{2}{k+2}$. In particular, G cannot be a planar graph with girth at least $2k + 6$.*

Proof. Let G be as stated. Every vertex has an initial charge equal to its degree. Every 3^+ -vertex gives $\frac{1}{k+2}$ to each of its 2-weak-neighbors. Let us check that the final charge $ch(v)$ of every vertex v is at least $2 + \frac{2}{k+2}$.

- If $d(v) = 2$, then v receives $\frac{1}{k+2}$ from each of its 3-weak-neighbors. Thus $ch(v) = 2 + \frac{2}{k+2}$.
- If $d(v) = 3$, then v gives $\frac{1}{k+2}$ to each of its 2-weak-neighbors. Thus $ch(v) \geq 3 - \frac{k}{k+2} = 2 + \frac{2}{k+2}$.
- If $d(v) = d \geq 4$, then v has at most $\frac{k+1}{2}$ 2-weak-neighbors in each of the d incident paths. Thus $ch(v) \geq d - d \left(\frac{k+1}{2}\right) \left(\frac{1}{k+2}\right) = \frac{d}{2} \left(1 + \frac{1}{k+2}\right) \geq 2 + \frac{2}{k+2}$.

This implies that $\text{mad}(G) \geq 2 + \frac{2}{k+2}$. Finally, if G is planar, then the girth of G cannot be at least $2k + 6$, since otherwise $(\text{mad}(G) - 2) \cdot (g(G) - 2) \geq \left(2 + \frac{2}{k+2} - 2\right) (2k + 6 - 2) = \left(\frac{2}{k+2}\right) (2k + 4) = 4$, which contradicts Proposition 8. \square

2.1 Proof of Theorem 3.1

We prove that the oriented planar graph \vec{T}_5 on 5 vertices from Figure 1 is $P_{28}^{(1,0)}$ -universal by contradiction. Assume that G is an oriented planar graph with girth at least 28 that does not admit a homomorphism to \vec{T}_5 and is minimal with respect to the number of vertices. By minimality, G cannot contain a vertex v with degree at most one since a \vec{T}_5 -coloring of $G - v$ can be extended to G . Similarly, G does not contain the following configurations.

- A path with 6 consecutive 2-vertices.
- A 3-vertex with at least 12 2-weak-neighbors.

Suppose that G contains a path $u_0u_1u_2u_3u_4u_5u_6u_7$ such that the degree of u_i is two for $1 \leq i \leq 6$. By minimality of G , $G - u_1, u_2, u_3, u_4, u_5, u_6$ admits a \vec{T}_5 -coloring φ . We checked on a computer that for any $\varphi(v_0)$ and $\varphi(v_6)$ in $V(\vec{T}_5)$ and every possible orientation of the 7 arcs u_iu_{i+1} , we can always extend φ into a \vec{T}_5 -coloring of G , a contradiction.

Suppose that G contains a 3-vertex v with at least 12 2-weak-neighbors. Let u_1, u_2, u_3 be the 3⁺-weak-neighbors of v and let l_i be the number of common 2-weak-neighbors of v and u_i , i.e., 2-vertices on the path between v and u_i . Without loss of generality and by the previous discussion, we have $5 \geq l_1 \geq l_2 \geq l_3$ and $l_1 + l_2 + l_3 \geq 12$. So we have to consider the following cases:

- **Case 1:** $l_1 = 5, l_2 = 5, l_3 = 2$.
- **Case 2:** $l_1 = 5, l_2 = 4, l_3 = 3$.
- **Case 3:** $l_1 = 4, l_2 = 4, l_3 = 4$.

By minimality, the graph G' obtained from G by removing v and its 2-weak-neighbors admits a \vec{T}_5 -coloring φ . Let us show that in all three cases, we can extend φ into a \vec{T}_5 -coloring of G to get a contradiction.

With an extensive search on a computer we found that if a vertex v is connected to a vertex u colored in $\varphi(u)$ by a path made of l 2-vertices ($0 \leq l \leq 5$) then v can be colored in:

- at least 1 color if $l = 0$,
- at least 2 colors if $l = 1$,
- at least 2 colors if $l = 2$ (the sets $\{c, d, e\}$ and $\{b, c, d\}$ are the only sets of size 3 that can be forbidden from v),
- at least 3 colors if $l = 3$,
- at least 4 colors if $l = 4$ and
- at least 4 colors if $l = 5$ (only the sets $\{b\}$, $\{c\}$, and $\{e\}$ can be forbidden from v).

In Case 1, u_3 forbids at most 3 colors from v since $l_3 = 2$. If it forbids less than 3 colors, we will be able to find a color for v since u_1 and u_2 forbid at most 1 color from v . The only sets of 3 colors that u_3 can forbid are $\{b, c, d\}$ and $\{c, d, e\}$. Since u_1 and u_2 can each only forbid b, c or e , we can always find a color for v .

In Case 2, u_1 and u_2 each forbid at most one color and u_3 forbids at most 2 colors so there remains at least one color for v .

In Case 3, u_1, u_2 , and u_3 each forbid at most one color, so there remains at least two colors for v .

We can always extend φ into a \vec{T}_5 -coloring of G , a contradiction.

So G contains at most 5 consecutive 2-vertices and every 3-vertex has at most 11 2-weak-neighbors. Using Lemma 9 with $k = 11$ contradicts the fact that the girth of G is at least 28.

2.2 Proof of Theorem 3.2

We prove that the 2-edge-colored planar graph T_6 on 6 vertices from Figure 2 is $P_{22}^{(0,2)}$ -universal by contradiction. Assume that G is a 2-edge-colored planar graphs with girth at least 22 that does not admit a homomorphism to T_6 and is minimal with respect to the number of vertices. By minimality, G cannot contain a vertex v with degree at most one since a T_6 -coloring of $G - v$ can be extended to G . Similarly, G does not contain the following configurations.

- A path with 5 consecutive 2-vertices.
- A 3-vertex with at least 9 2-weak-neighbors.

Suppose that G contains a path $u_0u_1u_2u_3u_4u_5u_6$ such that the degree of u_i is two for $1 \leq i \leq 5$. By minimality of G , $G - u_1, u_2, u_3, u_4, u_5$ admits a T_6 -coloring φ . We checked on a computer that for any $\varphi(v_0)$ and $\varphi(v_6)$ in $V(T)$ and every possible colors of the 6 edges u_iu_{i+1} , we can always extend φ into a T_6 -coloring of G , a contradiction.

Suppose that G contains a 3-vertex v with at least 9 2-weak-neighbors. Let u_1, u_2, u_3 be the 3⁺-weak-neighbors of v and let l_i be the number of common 2-weak-neighbors of v and u_i , i.e., 2-vertices on the path between v and u_i . Without loss of generality and by the previous discussion, we have $4 \geq l_1 \geq l_2 \geq l_3$ and $l_1 + l_2 + l_3 \geq 9$. So we have to consider the following cases:

- **Case 1:** $l_1 = 3, l_2 = 3, l_3 = 3$.
- **Case 2:** $l_1 = 4, l_2 = 3, l_3 = 2$.
- **Case 3:** $l_1 = 4, l_2 = 4, l_3 = 1$.

By minimality of G , the graph G' obtained from G by removing v and its 2-weak-neighbors admits a T_6 -coloring φ . Let us show that in all three cases, we can extend φ into a T_6 -coloring of G to get a contradiction.

With an extensive search on a computer we found that if a vertex v is connected to a vertex u colored in $\varphi(u)$ by a path P made of l 2-vertices ($0 \leq l \leq 4$) then v can be colored in:

- at least 1 color if $l = 0$ (the sets a, c, d, e, f and b, c, d, e, f of colors are the only sets of size 5 that can be forbidden from v for some $\varphi(u) \in T$ and edge-colors on P),
- at least 2 colors if $l = 1$ (the sets a, b, c, f and b, c, e, f are the only sets of size 4 that can be forbidden from v),
- at least 3 colors if $l = 2$ (the sets b, c, f, c, e, f and d, e, f are the only sets of size 3 that can be forbidden from v),
- at least 4 colors if $l = 3$ (the set c, b is the only set of size 2 that can be forbidden from v), and
- at least 5 colors if $l = 4$ (the sets c and f are the only sets of size 1 that can be forbidden from v).

Suppose that we are in Case 1. Vertices u_1, u_2 , and u_3 each forbid at most 2 colors from v since $l_1 = l_2 = l_3 = 3$. Suppose that u_1 forbids 2 colors. It has to forbid colors c and f (since it is the only pair of colors that can be forbidden by a path made of 3 2-vertices). If u_2 or u_3 also forbids 2 colors, they will forbid the exact same pair of colors. We can therefore assume that they each forbid 1 color from v . There are 6 available colors in T_6 , so we can always find a color for v and extend φ to a T_6 -coloring of G , a contradiction. We proceed similarly for the other two cases.

So G contains at most 4 consecutive 2-vertices and every 3-vertex has at most 8 2-weak-neighbors. Then Lemma 9 with $k = 8$ contradicts the fact that the girth of G is at least 22.

3 Proof of Theorem 4.1

We construct an oriented bipartite cactus graph with girth at least g and oriented chromatic number at least 5. Let g' be such that $g' \geq g$ and $g' \equiv 4 \pmod{6}$. Consider a circuit $v_1, \dots, v_{g'}$. Clearly, the oriented chromatic number of this circuit is 4 and the only tournament on 4 vertices it can map to is the tournament \vec{T}_4 induced by the vertices a, b, c , and d in \vec{T}_5 . Now we consider the cycle $C = w_1, \dots, w_{g'}$ containing the arcs $w_{2i-1}w_{2i}$ with $1 \leq i \leq g'/2$, $w_{2i+1}w_{2i}$ with $1 \leq i \leq g'/2 - 1$, and $w_{g'}w_1$.

Suppose for contradiction that C admits a homomorphism φ such that $\varphi(w_1) = d$. This implies that $\varphi(w_2) = a$, $\varphi(w_3) = d$, $\varphi(w_4) = a$, and so on until $\varphi(w_{g'}) = a$. Since $\varphi(w_{g'}) = a$ and $\varphi(w_1) = d$, $w_{g'}w_1$ should map to ad , which is not an arc of \vec{T}_4 , a contradiction.

Our cactus graph is then obtain from the circuit $v_1, \dots, v_{g'}$ and g' copies of C by identifying every vertex v_i with the vertex w_1 of a copy of C . This cactus graph does not map to \vec{T}_4 since one of the v_i would have to map to d and then the copy of C attached to v_i would not be \vec{T}_4 -colorable.

4 Proof of Theorem 4.2

We construct a 2-edge-colored bipartite outerplanar graph with girth at least g that does not map to a 2-edge-colored planar graph with at most 5 vertices. Let g' be such that $g' \geq g$ and $g' \equiv 2 \pmod{4}$. Consider an alternating cycle $C = v_0, \dots, v_{g'-1}$. For every $0 \leq i \leq g' - 3$, we add $g' - 2$ 2-vertices $w_{i,1}, \dots, w_{i,g'-2}$ that form the path $P_i = v_i w_{i,1} \dots w_{i,g'-2} v_{i+1}$ such that the edges of P_i get the color distinct from the color of the edge $v_i v_{i+1}$. Let G be the obtained graph. The 2-edge-colored chromatic number of C is 5. So without loss of generality, we assume for contradiction that G admits a homomorphism φ to a 2-edge-colored planar graph H on 5 vertices. Let us define $\mathcal{E} = \bigcup_{i \text{ even}} \varphi(v_i)$ and $\mathcal{O} = \bigcup_{i \text{ odd}} \varphi(v_i)$. Since C is alternating, $\varphi(v_i) \neq \varphi(v_{i+2})$ (indices are modulo g'). Since $g' \equiv 2 \pmod{4}$, there is an odd number of v_i with an even (resp. odd) index. Thus, $|\mathcal{E}| \geq 3$ and $|\mathcal{O}| \geq 3$. Therefore we must have $\mathcal{E} \cap \mathcal{O} \neq \emptyset$.

Notice that every two vertices v_i and v_j in G are joined by a blue path and a red path such that the lengths of these paths have the same parity as $i - j$. Thus, the blue (resp. red) edges of H must induce a connected spanning subgraph of H . Since $|V(H)| = 5$, H contains at least 4 blue (resp. red) edges. Since red and blue edges play symmetric roles in G and since $|E(H)| \leq 9$ by Proposition 8, we assume without loss of generality that H contains exactly 4 blue edges. Moreover, these 4 blue edges induce a tree. In particular, the blue edges induce a bipartite graph which partitions $V(H)$ into 2 parts. Thus, every v_i with even index is mapped into one part of $V(H)$ and every v_i with odd index is mapped into the other part of $V(H)$. So $\mathcal{E} \cap \mathcal{O} = \emptyset$, which is a contradiction.

5 Proof of Theorem 2

Let T be a $P_g^{(m,n)}$ -universal planar graph for some g that is minimal with respect to the subgraph order.

By minimality of T , there exists a graph $G \in P_g^{(m,n)}$ such that every color in T has to be used at least once to color G . Without loss of generality, G is connected, since otherwise we can replace G by the connected graph obtained from G by choosing a vertex in each component of G and identifying them. We obtain a graph G' from G as follows:

For each edge or arc uv in G , we keep uv in G' and we add $4m + n$ paths starting at u and ending at v made of vertices of degree 2:

- For each type of edge, we add a path made of $g - 1$ edges of this type.
- For each type of arc, we add two paths made of $g - 1$ arcs of this type such that the paths alternate between forward and backward arcs. We make the paths such that u is the tail of the first arc of one path and the head of the first arc of the other path.

- Similarly, for each type of arc we add two paths made of g arcs of this type such that the paths alternate between forward and backward arcs. We make the paths such that u is the tail of the first arc of one path and the head of the first arc of the other path.

Notice that G' is in $P_g^{(m,n)}$ and thus admits a homomorphism φ to T . Since G is a connected subgraph of G' and every color in T has to be used at least once to color G , we can find for each pair of vertices (c_1, c_2) in T and each type of edge a path (v_1, v_2, \dots, v_l) in G' made only of edges of this type such that $\varphi(v_1) = c_1$ and $\varphi(v_l) = c_2$.

This implies that for every pair of vertices (c_1, c_2) in T and each type of edge, there exists a walk from c_1 to c_2 made of edges of this type. Therefore, for $1 \leq j \leq n$, the subgraph induced by $E_j(T)$ is connected and contains all the vertices of T . So $E_j(T)$ contains a spanning tree of T . Thus T contains at least $|V(T)| - 1$ edges of each type.

Similarly, we can find for each pair of vertices (c_1, c_2) in T and each type of arc a path of even length $(v_1, v_2, \dots, v_{2l-1})$ in G' made only of arcs of this type, starting with a forward arc and alternating between forward and backward arcs such that $\varphi(v_1) = c_1$ and $\varphi(v_l) = c_2$. We can also find a path of the same kind with odd length.

This implies that for every pair of vertices (c_1, c_2) in T and each type of arc there exist a walk of odd length and a walk of even length from c_1 to c_2 made of arcs of this type, starting with a forward arc and alternating between forward and backward arcs. Let p be the maximum of the length of all these paths. Given one of these walks of length l , we can also find a walk of length $l + 2$ that satisfies the same constraints by going through the last arc of the walk twice more. Therefore, for every $l \geq p$, every pair of vertices (c_1, c_2) in T , and every type of arc, it is possible to find a homomorphism from the path P of length l made of arcs of this type, starting with a forward arc and alternating between forward and backward arcs to T such that the first vertex is colored in c_1 and the last vertex is colored in c_2 .

We now show that this implies that $|A_j(T)| \geq 2|V(T)| - 1$ for $1 \leq j \leq m$. Let P be a path $(v_1, v_2, \dots, v_p, v_{p+1})$ of length p starting with a forward arc and alternating between forward and backward arcs of the same type. We color v_1 in some vertex c of T . Let C_i be the set of colors in which vertex v_i could be colored. We know that $C_1 = c$ and C_2 is the set of direct successors of c . Set C_3 is the set of direct predecessors of vertices in C_2 so $C_1 \subseteq C_3$ and, more generally, $C_i \subseteq C_{i+2}$. Let uv be an arc in T . If $u \in C_i$ with i odd, then $v \in C_{i+1}$. If $v \in C_i$ with i even then $u \in C_{i+1}$. We can see that uv is capable of adding at most one vertex to a C_i (and every C_j with $j \equiv i \pmod{2}$ and $i \leq j$). We know that $C_{p+1} = V(T)$ hence T contains at least $2|V(T)| - 1$ arcs of each type.

Therefore, the underlying graph of T contains at least $m(2|V(T)| - 1) + n(|V(T)| - 1) = (2m + n)|V(T)| - m - n$ edges, which contradicts Proposition 8 for $2m + n \geq 3$.

6 Proof of Theorem 5.1

We construct an oriented bipartite 2-outerplanar graph with girth 14 that does not map to \vec{T}_5 .

The oriented graph X is a cycle on 14 vertices v_0, \dots, v_{13} such that the tail of every arc is the vertex with even index, except for the arc $v_{13}v_0$. Suppose for contradiction that X has a \vec{T}_5 -coloring h such that no vertex with even index maps to b . The directed path $v_{12}v_{13}v_0$ implies that $h(v_{12}) \neq h(v_0)$. If $h(v_0) = a$, then $h(v_1) \in \{b, c\}$ and $h(v_2) = a$ since $h(v_2) \neq b$. By contagion, $h(v_0) = h(v_2) = \dots = h(v_{12}) = a$, which is a contradiction. Thus $h(v_0) \neq a$. If $h(v_0) = c$, then $h(v_1) = d$ and $h(v_2) = c$ since $h(v_2) \neq b$. By contagion, $h(v_0) = h(v_2) = \dots = h(v_{12}) = c$, which is a contradiction. Thus $h(v_0) \neq c$. So $h(v_0) \notin \{a, b, c\}$, that is, $h(v_0) \in \{d, e\}$. Similarly, $h(v_{12}) \in \{d, e\}$. Notice that \vec{T}_5 does not contain a directed path xyz such that x and z belong to

$\{d, e\}$. So the path $v_{12}v_{13}v_0$ cannot be mapped to \vec{T}_5 . Thus X does not have a \vec{T}_5 -coloring h such that no vertex with even index maps to b .

Consider now the path P on 7 vertices p_0, \dots, p_6 with the arcs $\overrightarrow{p_1p_0}, \overrightarrow{p_1p_2}, \overrightarrow{p_3p_2}, \overrightarrow{p_4p_3}, \overrightarrow{p_5p_4}, \overrightarrow{p_5p_6}$. It is easy to check that there exists no \vec{T}_5 -coloring h of P such that $h(p_0) = h(p_6) = b$.

We construct the graph Y as follows: we take 8 copies of X called $X_{\text{main}}, X_0, X_2, X_4, \dots, X_{12}$. For every couple $(i, j) \in \{0, 2, 4, 6, 8, 10, 12\}^2$, we take a copy $P_{i,j}$ of P , we identify the vertex p_0 of $P_{i,j}$ with the vertex v_i of X_{main} and we identify the vertex p_6 of $P_{i,j}$ with the vertex v_j of H_i .

So Y is our oriented bipartite 2-outerplanar graph with girth 14. Suppose for contradiction that Y has a \vec{T}_5 -coloring h . By previous discussion, there exists $i \in \{0, 2, 4, 6, 8, 10, 12\}$ such that the vertex v_i of X_{main} maps to b . Also, there exists $j \in \{0, 2, 4, 6, 8, 10, 12\}$ such that the vertex v_j of X_i maps to b . So the corresponding path $P_{i,j}$ is such that $h(p_0) = h(p_6) = b$, a contradiction. Thus Y does not map to \vec{T}_5 .

7 Proof of Theorem 5.2

We construct a 2-edge-colored 2-outerplanar graph with girth 11 that does not map to T_6 . We take 12 copies X_0, \dots, X_{11} of a cycle of length 11 such that every edge is red. Let $v_{i,j}$ denote the j^{th} vertex of X_i . For every $0 \leq i \leq 10$ and $0 \leq j \leq 10$, we add a path consisting of 5 blue edges between $v_{i,11}$ and $v_{j,i}$.

Notice that in any T_6 -coloring of a red odd cycle, one vertex must map to c . So we suppose without loss of generality that $v_{0,11}$ maps to c . We also suppose without loss of generality that $v_{0,0}$ maps to c . The blue path between $v_{0,11}$ and $v_{0,0}$ should map to a blue walk of length 5 from c to c in T_6 . Since T_6 contains no such walk, our graph does not map to T_6 .

8 Proof of Theorem 5.3

We construct a 2-edge-colored bipartite 2-outerplanar graph with girth 10 that does not map to T_6 . By Theorem 4.2, there exists a bipartite outerplanar graph M with girth at least 10 such that for every T_6 -coloring h of M , there exists a vertex v in M such that $h(v) = c$.

Let X be the graph obtained as follows. Take a main copy Y of M . For every vertex v of Y , take a copy Y_v of M . Since Y_v is bipartite, let A and B the two independent sets of Y_v . For every vertex w of A , we add a path consisting of 5 blue edges between v and w . For every vertex w of B , we add a path consisting of 4 edges colored (blue, blue, red, blue) between v and w .

Notice that X is indeed a bipartite 2-outerplanar graph with girth 10. We have seen in the previous proof that T_6 contains no blue walk of length 5 from c to c . We also check that T_6 contains no walk of length 4 colored (blue, blue, red, blue) from c to c . By the property of M , for every T_6 -coloring h of X , there exist a vertex v in Y and a vertex w in Y_v such that $h(v) = h(w) = c$. Then h cannot be extended to the path of length 4 or 5 between v and w . So X does not map to T_6 .

9 Proof of Theorem 6.1

Let g be the largest integer such that there exists a graph in $P_g^{(1,0)}$ that does not map to \vec{T}_5 . Let $G \in P_g^{(1,0)}$ be a graph that does not map to \vec{T}_5 and such that the underlying graph of G is minimal with respect to the homomorphism order.

Let G' be obtained from G by removing an arbitrary arc v_0v_3 and adding two vertices v_1 and v_2 and the arcs v_0v_1, v_2v_1, v_2v_3 . By minimality, G' admits a homomorphism φ to \vec{T}_5 . Suppose for contradiction that $\varphi(v_2) = c$. This implies that $\varphi(v_1) = \varphi(v_3) = d$. Thus φ provides a \vec{T}_5 -coloring of G , a contradiction. So $\varphi(v_2) \neq c$ and, similarly, $\varphi(v_2) \neq e$.

Given a set S of vertices of \vec{T}_5 , we say that we force S if we specify a graph H and a vertex $v \in V(H)$ such that for every vertex $x \in V(\vec{T}_5)$, we have $x \in S$ if and only if there exists a

\vec{T}_5 -coloring φ of H such that $\varphi(v) = x$. Thus, with the graph G' and the vertex v_2 , we force a non-empty set $\mathcal{S} \subset V(\vec{T}_5) \setminus \{c, e\} = \{a, b, d\}$.

We use a series of constructions in order to eventually force the set $\{a, b, c, d\}$ starting from \mathcal{S} . Recall that $\{a, b, c, d\}$ induces the tournament \vec{T}_4 . We thus reduce \vec{T}_5 -coloring to \vec{T}_4 -coloring, which is NP-complete for subcubic bipartite planar graphs with any given girth [4].

These constructions are summarized in the tree depicted in Figure 3. The vertices of this forest contain the non-empty subsets of $\{a, b, d\}$ and a few other sets. In this tree, an arc from S_1 to S_2 means that if we can force S_1 , then we can force S_2 . Every arc has a label indicating the construction that is performed. In every case, we suppose that S_1 is forced on the vertex v of a graph H_1 and we construct a graph H_2 that forces S_2 on the vertex w .




Figure 3: Forcing the set $\{a, b, c, d\}$.

- Arcs labelled "out": The set S_2 is the out-neighborhood of S_1 in \vec{T}_5 . We construct H_2 from H_1 by adding a vertex w and the arc vw . Thus, S_2 is indeed forced on the vertex w of H_2 .
- Arcs labelled "in": The set S_2 is the in-neighborhood of S_1 in \vec{T}_5 . We construct H_2 from H_1 by adding a vertex w and the arc wv . Thus, S_2 is indeed forced on the vertex w of H_2 .
- Arc labelled "Z": Let g' be the smallest integer such that $g' \geq g$ and $g' \equiv 4 \pmod{6}$. We consider a circuit $v_1, \dots, v_{g'}$. For $2 \leq i \leq g'$, we take a copy of H_1 and we identify its vertex v with v_i . We thus obtain the graph H_2 and we set $w = v_2$. Let φ be any T_6 -coloring of H_2 . By construction, $\{\varphi(v_2), \dots, \varphi(v_{g'})\} \subset S_1 = \{a, b, d\}$. A circuit of length $\not\equiv 0 \pmod{3}$ cannot map to the 3-circuit induced by $\{a, b, d\}$, so $\varphi(v_1) \in \{c, e\}$. If $\varphi(v_1) = c$ then $\varphi(v_2) = d$ and if $\varphi(v_1) = e$ then $\varphi(v_2) = a$. Thus $S_2 = \{ad\}$.

10 Proof of Theorem 6.2

Let g be the largest integer such that there exists a graph in $P_g^{(0,2)}$ that does not map to T_6 . Let $G \in P_g^{(0,2)}$ be a graph that does not map to T_6 and such that the underlying graph of G is minimal with respect to the homomorphism order.

Let G' be obtained from G by subdividing an arbitrary edge v_0v_3 twice to create the path $v_0v_1v_2v_3$ such that the edges v_0v_1 and v_1v_2 are red and the edge v_2v_3 gets the color of the original edge v_0v_3 . By minimality, G' admits a homomorphism φ to T_6 . Suppose for contradiction




Figure 4: Forcing a good set.

that $\varphi(v_1) = f$. This implies that $\varphi(v_0) = \varphi(v_2) = b$. Thus φ provides a T_6 -coloring of G , a contradiction.

Given a set S of vertices of T_6 , we say that we force S if we specify a graph H and a vertex $v \in V(H)$ such that for every vertex $x \in V(T_6)$, we have $x \in S$ if and only if there exists T_6 -coloring φ of H such that $\varphi(v) = x$. Thus, with the graph G' and the vertex v_1 , we force a non-empty set $\mathcal{S} \subset V(T_6) \setminus \{f\} = \{a, b, c, d, e\}$.

Recall that the core of a graph is the smallest subgraph which is also a homomorphic image. We say that a subset S of $V(T_6)$ is *good* if the core of the subgraph induced by S is isomorphic to the graph T_4 which is a a clique on 4 vertices such that both the red and the blue edges induce a path of length 3. We use a series of constructions in order to eventually force a good set starting from \mathcal{S} . We thus reduce T_6 -coloring to T_4 -coloring, which is NP-complete for subcubic bipartite planar graphs with any given girth [5].

These constructions are summarized in the forest depicted in Figure 4. The vertices of this forest are the non-empty subsets of $\{a, b, c, d, e\}$ together with a few auxiliary sets of vertices containing f . In this forest, an arc from S_1 to S_2 means that if we can force S_1 , then we can force S_2 . Every set with no outgoing arc is good. We detail below the construction that is performed for each arc. In every case, we suppose that S_1 is forced on the vertex v of a graph H_1 and we construct a graph H_2 that forces S_2 on the vertex w .

- Blue arcs: The set S_2 is the blue neighborhood of S_1 in T_6 . We construct H_2 from H_1 by adding a vertex w adjacent to v such that vw is blue. Thus, S_2 is indeed forced on the vertex w of H_2 .
- Red arcs: The set S_2 is the red neighborhood of S_1 in T_6 . The construction is as above except that the edge vw is red.
- Dashed blue arcs: The set S_2 is the set of vertices incident to a blue edge contained in the subgraph induced by S_1 in T_6 . We construct H_2 from two copies of H_1 by adding a blue edge between the vertex v of one copy and the vertex v of the other copy. Then w is one of the vertices v .
- Dashed red arcs: The set S_2 is the set of vertices incident to a red edge contained in the subgraph induced by S_1 in T_6 . The construction is as above except that the added edge is red.
- Arc labelled "X": Let $g' = 2 \lceil g/2 \rceil$. We consider an even cycle $v_1, \dots, v_{g'}$ such that $v_1 v_{g'}$ is red and the other edges are blue. For every vertex v_i , we take a copy of H_1 and we identify its vertex v with v_i . We thus obtain the graph H_2 and we set $w = v_1$. Let φ be any T_6 -coloring of H_2 . In any T_6 -coloring of H_2 , the cycle $v_1, \dots, v_{g'}$ maps to a 4-cycle with exactly one red edge contained in the subgraph of T_6 induced by $S_1 = \{a, b, c, d, e\}$. These 4-cycles are $aedb$ with red edge ae and $cdba$ with red edge cd . Since w is incident to the red edge in the cycle $v_1, \dots, v_{g'}$, w can be mapped to a, e, c , or d but not to b . Thus $S_2 = \{a, c, d, e\}$.
- Arc labelled "Y": We consider an alternating cycle v_0, \dots, v_{8g-1} . For every vertex v_i , we take a copy of H_1 and we identify its vertex v with v_i . We obtain the graph H_2 by adding the vertex x adjacent to v_0 and v_{4g+2} such that xv_0 and xv_{4g+2} are blue. We set $w = v_0$. In any T_6 -coloring φ of H_2 , the cycle $v_1, \dots, v_{g'}$ maps to the alternating 4-cycle $acde$ contained in $S_1 = \{a, c, d, e\}$ such that $\varphi(v_i) = \varphi(v_{i+4 \pmod{8g}})$. So, a priori, either $\{\varphi(v_0), \varphi(v_{4g+2})\} = \{a, d\}$ or $\{\varphi(v_0), \varphi(v_{4g+2})\} = \{c, e\}$. In the former case, we can extend φ to H_2 by setting $\varphi(x) = b$. In the latter case, we cannot color x since c and e have no common blue neighbor in T_6 . Thus, $\{\varphi(v_0), \varphi(v_{4g+2})\} = \{a, d\}$ and $S_2 = \{a, d\}$.

References

- [1] J. Nešetřil and A. Raspaud. Colored homomorphisms of colored mixed graphs. *Journal of Combinatorial Theory, Series B*, 80(1):147–155, 2000.
- [2] H. Grötzsch. Ein dreifarbensatz für dreikreisfreie netze auf der kugel. *Wiss. Z. Martin-Luther-Universität Halle-Wittenberg Math.-Natur. Reihe*, 8:109–120, 1959.
- [3] O.V. Borodin, A.V. Kostochka, J. Nešetřil, A. Raspaud, and É. Sopena. On universal graphs for planar oriented graphs of a given girth. *Discrete Mathematics*, 188(1):73–85, 1998.
- [4] G. Guegan and P. Ochem. Complexity dichotomy for oriented homomorphism of planar graphs with large girth. *Theoretical Computer Science*, 596:142–148, 2015.
- [5] N. Movarraei and P. Ochem. Oriented, 2-edge-colored, and 2-vertex-colored homomorphisms. *Information Processing Letters*, 123:42–46, 2017.