



HAL
open science

On counting orientations for graph homomorphisms and for dually embedded graphs using the Tutte polynomial of matroid perspectives

Emeric Gioan

► **To cite this version:**

Emeric Gioan. On counting orientations for graph homomorphisms and for dually embedded graphs using the Tutte polynomial of matroid perspectives. ICGT 2022 - 11th International Colloquium on Graph Theory and Combinatorics, Jul 2022, Montpellier, France. lirmm-03868791

HAL Id: lirmm-03868791

<https://hal-lirmm.ccsd.cnrs.fr/lirmm-03868791>

Submitted on 24 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On counting orientations for graph homomorphisms and for dually embedded graphs using the Tutte polynomial of matroid perspectives¹²

Emeric Gioan — LIRMM, Université de Montpellier, CNRS, Montpellier, France

Abstract

An (oriented) matroid perspective (or morphism, or strong map, or quotient) is an ordered pair of (oriented) matroids satisfying some structural relationship. In this presentation, we will focus on the case of graphs, where two notable types of perspectives can be considered: graph homomorphisms, and dually embedded graphs on a surface. The Tutte polynomial of such a perspective is a classical polynomial (also called Las Vergnas polynomial in the case of dually embedded graphs), whose coefficients and (some) evaluations are known to count pairs of orientations of certain types. In this presentation, we show how coefficients and (other) evaluations of the polynomial also count pairs of orientations of certain types where some edge orientations are fixed, as well as some equivalence classes of pairs of orientations of certain types. These properties appear when the edge set is linearly ordered.

Dedicated to the memories of Claude Berge and Michel Las Vergnas, as Claude Berge used to be the thesis advisor of Michel Las Vergnas who used to be my own thesis advisor.

1 Perspectives for graphs

This paper briefly highlights applications to graph orientations of some results recently published in [5]. In general, an (oriented) matroid perspective (or morphism, or strong map, or quotient, up to unimportant variants) is an ordered pair $M \rightarrow N$ of (oriented) matroids on the same ground set satisfying some structural relationship, that can be characterized in various ways, such as: any circuit of M is a (conformal) union of circuits of N , or, equivalently, any cocircuit of N is a (conformal) union of cocircuits of M (see [11] or [16, Section 7.3] on matroid perspectives, and [1, Section 7.7] on oriented matroid perspectives). In the case of graphs, two notable types of perspectives can be considered, coming from graph homomorphisms and from graph embeddings. When we consider cycles and cuts of a graph, we always consider them as subsets of edges (vertices are not relevant here). We denote by $M_c(G)$ the cycle matroid of the graph G (whose circuits and cocircuits are inclusion-minimal cycles and cuts of G , respectively).

First situation. Consider an homomorphism between two graphs G and H , where vertices of G are identified to get vertices of H , while keeping a natural bijection between edges of G and H . Then any cycle of G corresponds to a union of cycles of H , and any cut of H corresponds to a union of cuts of G . (In what follows, we write “is” rather than “corresponds to”.) Then the two cycle matroids of these graphs form a perspective $M_c(G) \rightarrow M_c(H)$. One can focus on the following usual construction: given a graph $\overline{G} = (V, \overline{E})$ and $A \subseteq \overline{E}$, the two graphs $G = \overline{G} \setminus A$ and $H = \overline{G} / A$ form such an homomorphism. See Figure 1.




Figure 1: Perspective $M_c(G) \rightarrow M_c(H)$ from a graph homomorphism (with $G = \overline{G} \setminus e$ and $H = \overline{G} / e$).

Second situation. Given a graph H embedded on a closed surface, and H^* its embedded dual, we have that $M_c(H^*)^*$ (the dual matroid of the dual embedding of H) and $M_c(H)$ form a perspective $M_c(H^*)^* \rightarrow M_c(H)$. This is a classical result from [2] (the above definition is satisfied as cuts of H^* correspond to unions of cycles of H). Observe that if H is planar and embedded in the sphere, then $M_c(H^*)^* = M_c(H)$ (as matroid duality is consistent with planar graph duality) and then the perspective is trivial (that is, $M = N$). See an example in Figure 2.

¹Research supported by the ANR Grant DISTANCIA (Metric Graph Theory, ANR-17-CE40-0015) and the ANR Grant DIGRAPHS (Directed graphs, ANR-19-CE48-0013).

²Manuscript for the 11th International Colloquium on Graph Theory and combinatorics, Montpellier, France, ICGT 2022.




Figure 2: Perspective $M_c(H^*)^* \rightarrow M_c(H)$ from an embedding of H (blue) and H^* (red) in the torus. Observe that, in this example, $M_c(H^*)^*$ yields the same matroid as that of the graph G from Figure 1. Hence, this example illustrates both situations (but the two situations are not equivalent in general, as H^* might not be a planar graph and thus $M_c(H^*)^*$ might not be a graphic matroid).

In each situation, considering an orientation of a graph in the pair directly yields a consistent orientation for the other graph (use \overline{G} for homomorphisms, and use a \rightleftarrows rule for embeddings), and this pair of orientations effectively forms an oriented matroid perspective. We will consider the set of orientations of a perspective as the set 2^E , provided that a reference orientation, say $M \rightarrow N$, is given and then all orientations are obtained by reorienting some $A \subseteq E$ from this reference orientation, yielding $-_A M \rightarrow -_A N$.

2 Classical properties of the Tutte polynomial

The Tutte polynomial of a matroid perspective $M \rightarrow N$ is a 3-variable polynomial defined in [12, 14] in terms of rank functions. Ever since, this polynomial has been mainly studied by Michel Las Vergnas in a series of papers, including [13, 15]. See [6] for a recent survey. In the particular case where $M = N$, one retrieves the usual and well-known Tutte polynomial of the matroid M . In the above second situation for graphs, this polynomial was surveyed, deepened and named Las Vergnas polynomial in [3]. When the third variable is set to 1 (the third variable is not relevant in terms of orientations), it is given by:
$$T(M, N; x, y, 1) = \sum_{A \subseteq E} (x-1)^{r(N)-r_N(A)} (y-1)^{|A|-r_M(A)}.$$

Concerning orientations, the main theorem of [13] states that, for an oriented matroid perspective $M \rightarrow N$ on a linearly ordered set E , its Tutte polynomial satisfies

$$T(M, N; x, y, 1) = \sum_{A \subseteq E} \left(\frac{x}{2}\right)^{|O^*(-_A N)|} \left(\frac{y}{2}\right)^{|O(-_A M)|}, \text{ where}$$

$$\begin{aligned} O(M) &= \left\{ \min(C) \mid C \text{ positive circuit of } M \right\} && \text{(for a general oriented matroid } M) \\ &= \left\{ \min(C) \mid C \text{ directed cycle of } G \right\} && \text{(when } M = M_c(G), \text{ first situation)} \\ &= \left\{ \min(C) \mid C \text{ directed cut of } H^* \right\} && \text{(when } M = M_c(H^*)^*, \text{ second situation),} \\ O^*(N) &= \left\{ \min(C) \mid C \text{ positive cocircuit of } N \right\} && \text{(for a general oriented matroid } N) \\ &= \left\{ \min(C) \mid C \text{ directed cut of } H \right\} && \text{(when } N = M_c(H), \text{ in the two situations).} \end{aligned}$$

The sets $O(M)$ and $O^*(N)$ are known as the sets of *active* and *dual-active* elements of M and N , respectively. Note that they are two dual notions, as $O(M^*) = O^*(M)$. Observe that $O(M) = \emptyset$ if and only if M is acyclic, and that $O^*(N) = \emptyset$ if and only if N is totally cyclic (that is, strongly connected for a connected digraph). Let us denote by $T_{i,j}$ the coefficient of $x^i y^j$ of T . This number is an invariant (it does not depend on the ordering), and the above theorem can be summed up as:

$$T_{i,j} = 1/2^{i+j} \times \#\{\text{orientations with } i \text{ dual-active elements and } j \text{ active elements}\}.$$

See [7] for a survey and geometric interpretations. Notably, we get the following counting results:

Orientations $-_A M \rightarrow -_A N$ with	are counted by
no condition	$T(M, N; 2, 2, 1)$
acyclic $-_A M$	$T(M, N; 2, 0, 1)$
totally cyclic $-_A N$	$T(M, N; 0, 2, 1)$
acyclic $-_A M$ and totally cyclic $-_A N$	$T(M, N; 0, 0, 1)$

Naturally, the useful translation of the terms “acyclic” and “totally cyclic” depends on the situation considered for graphs. For instance, for connected graphs, $T(M, N; 0, 0, 1) = T_{0,0}$ counts, in the first situation, the number of orientations such that G is acyclic and H is strongly connected, and, in the second situation, the number of orientations such that both H and H^* are strongly connected.

3 More involved notions and results of the same flavour

The results below from [5] are obtained by means of active partitions which refine active elements, and by means of a partition of the set of orientations into activity classes. These notions for digraphs and oriented matroids were defined and used in the context of the active bijection in a series of papers, including [4, 8, 9, 10] (the Tutte polynomial also counts objects related to spanning trees or bases, see [7] for a survey). This corresponds to trivial perspectives (i.e., $M = N$). These notions are generalized in [5] for perspectives, where one can also find a 4-variable expansion formula for the Tutte polynomial, which can be seen as an algebraic counterpart of these constructions, and which was announced in [15]. The interpretations in the two situations for graphs follow the terminology presented in the previous section.

Definition 1. Let $M \rightarrow N$ be a perspective on a linearly ordered set E and $A \subseteq E$. The reorientation $-_A M \rightarrow -_A N$ is called *active-fixed* if no active element of $-_A M$ belongs to A , and *dual-active-fixed* if no dual-active element of $-_A N$ belongs to A . In other words, (dual-)active-fixed means that every (dual-)active element has the same orientation as in the reference orientation.

Definition 2. Let $M \rightarrow N$ be a perspective on a linearly ordered set E with $O(M) = \{g_1, \dots, g_j\}_<$ and $O^*(N) = \{h_1, \dots, h_i\}_<$. The *active filtration* of $M \rightarrow N$ is sequence of sets

$$\emptyset = G_j \subset \dots \subset G_1 \subset G_0 \subseteq H_0 \subset H_1 \subset \dots \subset H_i = E$$

defined by: $G_j = \emptyset$; for $0 \leq k \leq j-1$, G_k is the union of all positive circuits of M whose smallest element is greater than or equal to g_{k+1} (use directed cycles of G in the first graph situation, or directed cuts of H^* in the second graph situation); $E \setminus H_i = \emptyset$; and, for $0 \leq k \leq i-1$, $E \setminus H_k$ is the union of all positive cocircuits of N whose smallest element is greater than or equal to h_{k+1} (use directed cuts of H in the two graph situations). Observe that $H_0 \setminus G_0$ is the (possibly empty) set of elements belonging to positive cocircuits of M and positive circuits of N .

The successive differences of these sets form a partition of E , named the *active partition* of $M \rightarrow N$, with j *cyclic parts* contained in G_0 , i *acyclic parts* contained in $E \setminus H_0$, and a (possibly empty) *hybrid part* $H_0 \setminus G_0$. The smallest elements of the (a)cyclic parts are the (dual-)active elements.

Then, the *activity class* of $M \rightarrow N$ is the set of 2^{i+j} reorientations obtained by arbitrarily reorienting cyclic and/or acyclic parts of the active partition of $M \rightarrow N$ (or, equivalently, subsets G_k , $0 \leq k \leq j-1$, and/or $E \setminus H_k$, $0 \leq k \leq i-1$, given by its active filtration). Observe that the hybrid part is not reoriented.

Theorem 3. *Let $M \rightarrow N$ be a perspective on a linearly ordered set E . Given a reorientation of $M \rightarrow N$, all reorientations in the same activity class as this reorientation share the same active filtration/partition. Activity classes of orientations form a partition of the set 2^E of orientations (into boolean lattices, actually). In each activity class of orientations; there is one and only one orientation which is active-fixed and dual-active-fixed. We also get the following counting results.*

$$\begin{aligned} T_{i,j} &= \#\{\text{active-fixed dual-active-fixed orientations with } i \text{ dual-active elements and } j \text{ active elements}\} \\ &= \#\{\text{activity classes of orientations with } i \text{ dual-active elements and } j \text{ active elements}\} \end{aligned}$$

Orientations $-_A M \rightarrow -_A N$ with	Activity classes with	are counted by
acyclic $-_A M$ and dual-active-fixed $-_A N$	acyclic $-_A M$	$T(M, N; 1, 0, 1)$
active-fixed $-_A M$ and totally cyclic $-_A N$	totally cyclic $-_A N$	$T(M, N; 0, 1, 1)$
active-fixed $-_A M$		$T(M, N; 2, 1, 1)$
dual-active-fixed $-_A N$		$T(M, N; 1, 2, 1)$
active-fixed $-_A M$ and dual-active-fixed $-_A N$	no condition	$T(M, N; 1, 1, 1)$
	acyclic $-_A M$ and tot. cyclic $-_A N$	$T(M, N; 0, 0, 1)$

Let us illustrate Theorem 3 on the example of Figures 1 and 2. See Figure 3. We have

$$T(M, N; x, y, 1) = 4\left(\frac{x}{2}\right)^2 + 8\left(\frac{x}{2}\right) + 2\left(\frac{y}{2}\right) + 2 = x^2 + 4x + y + 2,$$

$$\begin{aligned} T(M, N; 2, 2, 1) &= 16, & T(M, N; 1, 1, 1) &= 8, & T(M, N; 0, 0, 1) &= 2, \\ T(M, N; 2, 1, 1) &= 15, & T(M, N; 2, 0, 1) &= 14, & T(M, N; 1, 0, 1) &= 7, \\ T(M, N; 1, 2, 1) &= 9, & T(M, N; 0, 2, 1) &= 4, & T(M, N; 0, 1, 1) &= 3. \end{aligned}$$

$-_A M$	$-_A N$	$O(-_A M)$		$-A$	active-fixed		dual-active-fixed		$-_A M$	$-_A N$	$O(-_A M)$		$-A$	active-fixed		dual-active-fixed	
		cyclic parts	acyclic parts		hybrid part	opp.	opp.	cyclic parts			acyclic parts	hybrid part		opp.	opp.		
		\emptyset	13	$-\emptyset$	\times	\times	\times	\times			\emptyset	3	-2	\times	\times	\times	\times
		\emptyset	12+34	\emptyset	\times	\cdot	\cdot	\cdot			\emptyset	34	12	\times	\cdot	\cdot	\cdot
		\emptyset	1	-4	\times	\times	\times	\times			1	\emptyset	-24	\times	\times	\cdot	\times
		\emptyset	12	34	\times	\cdot	\cdot	\cdot			1234	\emptyset	\emptyset	\cdot	\times	\times	\times
		\emptyset	1	-3	\times	\times	\times	\times			\emptyset	\emptyset	-23	\times	\times	\times	\times
		\emptyset	12	34	\times	\cdot	\cdot	\cdot			\emptyset	\emptyset	1234	\times	\times	\times	\times
		\emptyset	13	-34	\times	\cdot	\cdot	\cdot			\emptyset	3	-234	\times	\cdot	\times	\cdot
		\emptyset	12+34	\emptyset	\times	\cdot	\cdot	\cdot			\emptyset	34	12	\times	\times	\times	\times

Figure 3: List of orientations for the perspective of Figures 1 and 2. We use the natural ordering $1 < 2 < 3 < 4$. Only half the orientations are represented (the other half are the opposites, with the same active elements, dual-active elements, and active partitions). The reference orientation is the first of the list (with $A = \emptyset$). At each orientation, we indicate the set A of reoriented edges, the active and dual-active elements, and the active partition below. Next, we indicate (with a \times) if it is active-fixed, dual-active-fixed, and the same information below for its opposite. Let us detail the $T(M, N; 1, 1, 1) = 8$ activity classes. For the two light grey cells on the left: these orientations and their opposites are all in the same activity class of size 4 given by $A \in \{\emptyset, 34, 12, 1234\}$. For the dark grey cell on the right: this orientation and its opposite are those whose hybrid part is equal to E , and each forms an activity class of size 1 (they are enumerated by $T(M, N; 0, 0, 1) = 2$). Every other activity class has size 2: it consists of one orientation represented in a cell and its opposite (when the hybrid part is empty, that is, for $A \in \{24, 13\}$), or of one orientation and the opposite of an orientation represented in another cell (e.g., for $A \in \{4, 124\} = \{4, E\}3$), or of two represented orientations (for $A \in \{2, 234\}$), or of the opposite orientations of these two (for $A \in \{134, 1\}$).

References

- [1] A. Björner, M. Las Vergnas, B. Sturmfels, N. White, and G. Ziegler. *Oriented Matroids*, volume 46 of *Encyclopedia of Mathematics and Its Applications*. Cambridge University Press, second edition, 1999.
- [2] J. Edmonds. On the surface duality of linear graphs. *J. Res. Nat. Bur. Stand.*, 69:121–123, 1965.
- [3] J.A. Ellis-Monaghan and I. Moffatt. The Las Vergnas polynomial for embedded graphs. *Eur. J. Comb.*, 50:97–114, 2015. *Combinatorial Geometries: Matroids, Oriented Matroids and Applications. Special Issue in Memory of M. Las Vergnas*.
- [4] E. Gioan. *Correspondance naturelle entre bases et réorientations des matroïdes orientés*. Ph.D. Thesis, Univ. Bordeaux 1, 2002.
- [5] E. Gioan. On Tutte polynomial expansion formulas in perspectives of matroids and oriented matroids. *Disc. Math.*, 345(7), 112796, 2022.
- [6] E. Gioan. The Tutte polynomial of matroid perspectives. In J. Ellis-Monaghan and I. Moffatt, editors, *Handbook of the Tutte Polynomial and Related Topics*, CRC Monographs and Research Notes in Mathematics. In press.
- [7] E. Gioan. The Tutte polynomial of oriented matroids. In J. Ellis-Monaghan and I. Moffatt, editors, *Handbook of the Tutte Polynomial and Related Topics*, CRC Monographs and Research Notes in Mathematics. In press.
- [8] E. Gioan and M. Las Vergnas. Activity preserving bijections between spanning trees and orientations in graphs. *Discrete Mathematics*, 298:169–188, 2005.
- [9] E. Gioan and M. Las Vergnas. The active bijection for graphs. *Advances in Applied Mathematics*, 104:165–236, 2019.
- [10] E. Gioan and M. Las Vergnas. The active bijection - 2.b - Decomposition of activities for oriented matroids, and general definitions of the active bijection. *Submitted, preprint available at arXiv:1807.06578*.
- [11] J. Kung. Strong maps. In N. White, editor, *Theory of matroids*, volume 26 of *Encyclopedia of Mathematics and Its Applications*. Cambridge University Press, 1986.
- [12] M. Las Vergnas. On the Tutte polynomial of a morphism of matroids. *Annals of Discrete Mathematics*, 8:7–20, 1980.
- [13] M. Las Vergnas. The Tutte polynomial of a morphism of matroids II. Activities of orientations. In J.A. Bondy and U.S.R. Murty, editors, *Progress in Graph Theory*, pages 367–380. Academic Press, Toronto, Canada, 1984.
- [14] M. Las Vergnas. The Tutte polynomial of a morphism of matroids I. Set-pointed matroids and matroid perspectives. *Ann. Inst. Fourier, Grenoble*, 49(3):973–1015, 1999.
- [15] M. Las Vergnas. The Tutte polynomial of a morphism of matroids 6. A multi-faceted counting formula for hyperplane regions and acyclic orientations. 2012. Unpublished, preliminary preprint available at arXiv 1205.5424.
- [16] J.G. Oxley. *Matroid Theory*. Oxford Graduate Texts in Mathematics. Oxford University Press, 2011 (2nd ed.).