



**HAL**  
open science

# Structural and algorithmic aspects of partial orderings of graphs

Jean-Florent Raymond

► **To cite this version:**

Jean-Florent Raymond. Structural and algorithmic aspects of partial orderings of graphs. Discrete Mathematics [cs.DM]. Université de Montpellier; University of Warsaw, 2016. English. NNT: . tel-01486769

**HAL Id: tel-01486769**

**<https://hal-lirmm.ccsd.cnrs.fr/tel-01486769>**

Submitted on 10 Mar 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

University of Warsaw  
Faculty of Mathematics, Informatics, and Mechanics  
and  
University of Montpellier  
LIRMM, école doctorale I2S

Jean-Florent Raymond

# Structural and algorithmic aspects of partial orderings of graphs

*PhD dissertation*

Supervisors

dr hab. Marcin Kamiński  
Institute of Informatics  
University of Warsaw

prof. dr hab. Dimitrios M. Thilikos  
AlGCo project team  
CNRS, LIRMM  
Department of Mathematics  
National and Kapodistrian University of Athens

October 2016

Author's declaration:  
I hereby declare that this dissertation is my own work.

Date: .....

*Jean-Florent Raymond*

Supervisors' declaration:  
The dissertation is ready to be reviewed.

Date: .....

*dr hab. Marcin Kamiński*

.....  
*prof. dr hab. Dimitrios M. Thilikos*

This thesis is centered around the study of exclusion theorems in Graph Theory and their applications. Toward this, we obtain structural decompositions for graphs excluding a given sub-structure, or bounds on certain invariants of these graphs. A first application is the characterization of some well-quasi-ordered graph classes defined by excluded sub-structures. A second application is the study of the relations between invariants related to problems of packing and covering. We establish new connections between these invariants and present algorithmic applications of these results.

*Keywords:* Graph Theory, exclusion theorems, forbidden substructures, packing and covering, well-quasi-ordering, combinatorial dichotomies.

*2000 MSC:* 05C35, 05C70, 06A06, 68R10, 05C83.



---

# CONTENTS

<b>Contents</b>	<b>5</b>
<b>Acknowledgements</b>	<b>9</b>
<b>Une introduction en français</b>	<b>11</b>
Beaux préordres . . . . .	14
La propriété d’Erdős–Pósa . . . . .	16
Aperçu des résultats et organisation de la thèse . . . . .	20
<b>1 Introduction</b>	<b>23</b>
Well-quasi-ordering . . . . .	26
The Erdős–Pósa property . . . . .	28
Overview of the results and organization of the thesis . . . . .	31
<b>2 General definitions</b>	<b>33</b>
2.1 Basics . . . . .	33
2.2 Orders . . . . .	33
2.3 Graphs . . . . .	34
2.3.1 Basics . . . . .	34
2.3.2 Special graphs and graph classes . . . . .	36
2.4 Graph operations and orderings . . . . .	38
2.4.1 Local operations . . . . .	38
2.4.2 Containment models . . . . .	39
2.4.3 Graph orderings . . . . .	41
2.5 Tree-like decompositions and width parameters . . . . .	43
<b>3 Graphs and well-quasi-ordering</b>	<b>49</b>
3.1 Preliminaries on well-quasi-orders . . . . .	49
3.2 Notions specific to this chapter . . . . .	56
3.3 Raising well-quasi-orders . . . . .	58

3.4	A high-level view of the proofs . . . . .	61
3.5	The bestiary . . . . .	63
3.5.1	Antichains for induced minors . . . . .	63
3.5.2	Antichains for contractions . . . . .	65
3.6	Induced minors and well-quasi-ordering . . . . .	66
3.6.1	The dichotomy theorem . . . . .	68
3.6.2	Graphs not containing $\hat{K}_4$ . . . . .	71
3.6.3	Graphs not containing Gem . . . . .	87
3.7	Contractions and well-quasi-ordering . . . . .	95
3.7.1	On graphs with no diamond . . . . .	95
3.7.2	Well-quasi-ordering clique-cactus graphs . . . . .	98
3.7.3	The dichotomy . . . . .	100
3.7.4	Canonical antichains and contractions . . . . .	102
3.8	Multigraph contractions and well-quasi-ordering . . . . .	104
3.8.1	Gluing graphs . . . . .	106
3.8.2	Well-quasi-ordering graphs without big bonds . . . . .	110
3.8.3	Canonical antichains and multigraph contractions . . . . .	112
<b>4</b>	<b>Exclusion theorems</b> . . . . .	<b>115</b>
4.1	Clique-majors in graphs of large $\theta_r$ -girth . . . . .	115
4.1.1	The quest for large clique-majors . . . . .	116
4.1.2	Definitions specific to this section . . . . .	117
4.1.3	Finding small $\theta_r$ -majors . . . . .	119
4.1.4	Structural results on graphs with large $\theta_r$ -girth . . . . .	121
4.1.5	Remarks on the $H$ -girth . . . . .	128
4.2	Degree and $\theta_r$ -packings . . . . .	128
4.2.1	On maximum degree and edge-disjoint packings . . . . .	128
4.2.2	On the minimum degree and vertex-disjoint packings . . . . .	131
4.3	Treewidth and excluded majors . . . . .	132
4.3.1	Our tools . . . . .	135
4.3.2	Excluding a wheel . . . . .	137
4.3.3	Excluding a double wheel . . . . .	140
4.3.4	Excluding a graph of pathwidth at most 2 . . . . .	143
4.3.5	Excluding a yurt graph . . . . .	148
4.3.6	Excluding a union of $k$ disjoint copies of $\theta_r$ . . . . .	148
4.3.7	Lower bounds . . . . .	151
4.4	Immersion of planar subcubic graphs in graphs of large tree-cut width . . . . .	152
<b>5</b>	<b>The Erdős-Pósa property</b> . . . . .	<b>157</b>
5.1	Introduction . . . . .	157
5.2	Notions related to the Erdős-Pósa property . . . . .	159
5.3	General tools for proving Erdős-Pósa type results . . . . .	159
5.3.1	The Erdős-Pósa property from graph decompositions . . . . .	160
5.3.2	The Erdős-Pósa property from girth . . . . .	165
5.4	Applications to selected classes of graphs . . . . .	166

5.4.1	Wheels, yurts, and graphs of pathwidth at most two . . . . .	167
5.4.2	Pumpkins . . . . .	167
5.4.3	Double pumpkins . . . . .	169
5.4.4	Planar subcubic graphs . . . . .	171
5.5	Negative results . . . . .	177
5.6	Summary of results . . . . .	182
5.6.1	Results in terms of containment relations . . . . .	182
5.6.2	Results in terms of graph classes . . . . .	184
<b>6</b>	<b>From the Erdős–Pósa property to approximation</b>	<b>193</b>
6.1	Introduction . . . . .	193
6.2	Definitions specific to this section . . . . .	195
6.2.1	Basic definitions . . . . .	195
6.2.2	Boundaried graphs . . . . .	196
6.2.3	Encodings, signatures, and folios . . . . .	199
6.3	The reduction procedure . . . . .	201
6.4	Approximation meets the Erdős–Pósa property . . . . .	207
6.4.1	Reduce or progress . . . . .	208
6.4.2	Approximation algorithms . . . . .	209
<b>7</b>	<b>Conclusion</b>	<b>211</b>
7.1	General contribution . . . . .	211
7.2	Perspectives and problems . . . . .	213
7.2.1	On the $H$ -girth . . . . .	213
7.2.2	On the Erdős–Pósa property . . . . .	214
7.2.3	On edge-protrusions . . . . .	215
7.2.4	On well-quasi-ordering . . . . .	215
7.3	Publications . . . . .	216
7.3.1	On the Erdős–Pósa property . . . . .	216
7.3.2	On well-quasi-ordering . . . . .	217
7.3.3	On exclusion results . . . . .	217
7.3.4	On other topics . . . . .	217
	<b>Bibliography</b>	<b>219</b>
	<b>List of Figures</b>	<b>237</b>
	<b>List of Tables</b>	<b>241</b>
	<b>Index</b>	<b>243</b>



---

# ACKNOWLEDGEMENTS

I cannot start this section without thanking my two advisors Marcin Kamiński and Dimitrios Thilikos. Thank you Marcin and Dimitrios for your constant availability and for the freedom you gave me during these three years under your guidance. I enjoyed the different topics we studied together, the blackboard sessions in Warsaw, Montpellier and Paris, as well as your advice and discussions about the academic world and life in general. Thank you Marcin for your valuable help with the Polish paperwork.

Then, I would like to thank all the members of my defence committee: Stéphane Bessy, Fedor Fomin, Petr Golovach, Laurent Imbert, Łukasz Kowalik, Nicolas Nisse, Jerzy Tiuryn, and Nicolas Trotignon. Thank you for accepting to be part of my committee and for travelling to Montpellier the day of my defense. Thank you Petr and Nicolas N. for accepting to review this thesis, for your interest in my work, for your careful reading of my manuscript and for the improvements you suggested. In general, thanks to all those who suggested improvements to the manuscript.

I also want to thank Agnieszka Graff who accepted to be my examiner for the complementary discipline and who provided me with good reading.

Je remercie l'équipe AlGCo du LIRMM pour son ambiance, sa dynamique, pour la proximité qui y règne entre thésards et permanents, pour les repas (festins ?) d'équipe, pour les voyages que vous m'avez permis de faire. Thanks also to the Algorithms team of the University of Warsaw. Merci aux autres thésards montpelliérains pour avoir distrait mes repas par vos débats toujours différents : Julien Baste, François Dross, Valentin Garnero, Guillaume Guégan, Sabrina Ouazzani, Swann Rocher, et j'en oublie sans doute. Thanks also to my officemates in Warsaw.

Thanks to all my coauthors and all those with whom I have had interesting research discussions, in particular Aistis Atminas, Florian Barbero, Jarosław Błasiok, Nicolas Bousquet, Dimitris Chatzidimitriou, Vincent Cohen-Addad, François Dross, Archontia Giannopoulou, Gwenaél Joret, O-joung Kwon, Guillaume Lagarde, Marcin Kamiński, Michał Pilipczuk, Ignasi Sau, Dimitrios Thilikos, Théophile Trunck, and Marcin Wrochna. Remerciements à Théophile pour m'avoir re-conditionné à manger à heures fixes quand il était à Varsovie.

Que soient ici remerciés tous les professeurs qui m'ont donné envie d'aller en cours et d'étudier, comme Richard Ganaye, Daniel Hirschkoff, Alexandre Miquel, Éric Thierry et Stéphan Thomassé. Je me dois aussi de remercier Binh-Minh Bui-Xuân et Philippe Trébuchet qui m'ont offert ma toute première expérience de recherche en théorie des graphes pendant mon stage de L3.

Merci à tous ceux qui m'ont aidé à vivre avec ma phobie administrative. Je pense d'abord à Nicolas Serrurier, qui m'a régulièrement aidé (toujours avec la sourire) pour la cotutelle, les réinscriptions et à chaque fois que j'avais une question, à Caroline Ycre, Caroline Lebrun et Laurie Laverne. Merci à Marianne Huchard pour son aide déterminante dans la validation de mon jury. Dziękuję to the members of the SOB office of our department in Warsaw, Daniel Adamiak, Magda Szwaranowicz, and Iwona Zielińska, who continuously helped me to declare my trips and use my grant. Many thanks to Marcin Wrochna who printed the version of the manuscript that has been sent to the reviewers. Thanks also to the (Polish) National Science Center (NCN) for awarding me with a Preludium grant which funded most of my trips. I would also like to thank the contributors of the tikz drawing package.

Je me dois également de remercier ici ceux qui m'ont accompagné dans des activités non liées à la recherche. Merci à l'équipe de grimpeurs avides du LIRMM (ou d'ailleurs) Paolo Boscaro, Laurie Bougeois, Daniel Gonçalves, Bruno Grenet, Mickaël Montassier, Christophe Paul, Max, Nico et tous les autres pour la dynamique qu'ils maintiennent. Thanks to the happy climbers I met in Warsaw, Łukasz Kowalik, Michał Skrzypczak, and Elisa Therani. Merci aux joyeux spéléos du CLPA avec qui j'ai partagé explorations et moments de convivialité dans les nombreux avens du Larzac, en particulier Mickaël Bordes, Jean-Marie Catuogno, Rémy Cavin, Yann Guessard, Ludovic Leterme, Jean-Claude Molière, Sylvain Piry, Nicolas Tarlier, Thomas Valding et Claude Viladomat.

Merci à mes amis de prépa et de l'ENS de Lyon (et d'Uppsala), pour vos discussions, vos visites et votre hospitalité : Paul-Elliot Anglès d'Auriac, Basile Augé, Vincent Cohen-Addad, Emmanuelle Deléglise, Loïc Cotlb-Froidevaux, Lolo(ïc) Gaillard, Maxime Herda, Lucca Hirschi, Guillaume Lagarde, Mémé(lanie) Séon, Isabelle Schummers, Thomas Sibut et tous les autres, à mes coolocataires rondelesiens Grégoire Vauquiez et Michele Botti pour tous ces bons moments passés ensemble, à Chloé Armatnat et Yoann Genolini pour toutes nos aventures au bout d'une corde ou à skis et à Héloïse Verdan pour nos pérégrinations en train, en vélo, à pieds ou en stop et pour nos discussions sur l'art et sur la vie. Merci à ma famille pour son soutien, à mes parents, mes sœurs, mes grands parents, mes oncles et tantes et surtout merci à ma dulcinée, parce qu'on se comprend.

---

# UNE INTRODUCTION EN FRANÇAIS

This chapter is a translation of **Chapter 1** to French language.

Imaginons une collection d'objets, possiblement infinie. Dans cette collection, certains objets peuvent être plus désirables que d'autres. Il peut aussi arriver que deux objets soient incomparables, c'est à dire qu'aucun ne soit plus désirable que l'autre. Une illustration est donnée dans la **figure 1**, où les objets ordonnés sont des fruits. Dans un tel diagramme, qu'on appelle diagramme de Hasse, un élément  $x$  est supérieur (pour l'ordre considéré) à un élément  $y$  s'il y a un chemin de haut en bas reliant  $x$  à  $y$ . Par exemple, on voit sur le diagramme de la **figure 1** que la cerise est plus désirable que le kiwi, car le chemin cerise-abricot-kiwi va de bas en haut. Il faut noter que certains fruits, comme la framboise et la mirabelle, ne sont pas comparables.




FIGURE 1 : Quelques fruits ordonnés par désirabilité.

Fixons un objet  $x$  et considérons tous les objets qui ne sont pas plus désirables que  $x$ . Cela inclut les objets qui sont moins désirables que  $x$ , mais aussi ceux qui n'y sont pas comparables. Les fruits de la **figure 1** dont le nom est souligné sont ceux qui ne sont pas plus désirables que l'abricot. Un thème central de cette thèse, pour diverses collections d'objets et notions de *désirabilité*, est l'étude des objets qui ne sont pas plus désirables que  $x$ . À quoi ressemblent-ils ? Quelle est leur *structure* ? Quelles

sont leurs propriétés? Pour continuer l'exemple de la [figure 1](#), on peut remarquer qu'un point commun à tous les fruits qui ne sont pas plus désirables que l'abricot est d'avoir des pépins<sup>1</sup>.

Les objets que nous considérons dans cette thèse sont les graphes et le concept de désirabilité est remplacé par divers ordres partiels : sous-graphes, sous-graphes induits, mineurs, mineurs topologiques, mineurs induits, immersions, etc., qui la plupart du temps expriment qu'un graphe est *contenu* comme sous-structure d'un autre. La [figure 2](#) présente les (classes d'isomorphisme des) graphes qui peuvent être obtenus en subdivisant les arêtes du graphe complet à cinq sommets (dessiné au bas de la figure), ordonné(e)s par l'ordre correspondant. Bien entendu, cette figure est incomplète car il y a une infinité de tels graphes.




FIGURE 2 : Subdivisions de  $K_5$ .

Un des premiers théorèmes à établir les propriétés d'une classe de graphes définie par sous-structure interdite est la caractérisation des graphes planaires par Kuratowski (voir la [figure 3](#) pour un dessin des graphes mentionnés).

**Théorème 0.1** (Théorème de Kuratowski, 1930 [[Kur30](#)]<sup>2</sup>). *Un graphe est planaire<sup>3</sup> si et seulement si il ne contient pas de sous-graphe qui est une subdivision de  $K_5$  ou  $K_{3,3}$ .*

<sup>1</sup>Un fruit de ce diagramme montre que la présence de pépins n'est pas une caractérisation des fruits qui ne sont pas plus désirables que l'abricot (lequel?).

<sup>2</sup>D'après [[Bur78](#)], ce résultat a été indépendamment obtenu par Pontryagin, mais jamais publié.

<sup>3</sup>On dit qu'un graphe est *planaire* s'il peut être dessiné dans le plan sans croisement d'arêtes.




FIGURE 3 : Les graphes de Kuratowski  $K_5$  et  $K_{3,3}$  (de gauche à droite).

Une particularité notable de ce résultat est qu’il décrit une propriété topologique, être plane, en termes purement combinatoires : l’absence de certaines sous-structures dans le graphe. Souvent présenté comme un précurseur de la théorie des graphes moderne, le théorème de Kuratowski a été l’initiateur d’une longue lignée de résultats sur les classes définies par *exclusion* d’une sous-structure donnée. L’un d’entre eux est le théorème de Turán, qui donne un majorant au nombre d’arêtes d’un graphe ne contenant par un grand sous-graphe complet.

**Théorème 0.2** (Théorème de Turán, 1941 [Tur41]). *Soient  $p \geq 2$  un entier et  $G$  un graphe qui ne contient pas de sous-graphe complet à  $p$  sommets. Si  $G$  a  $n$  sommets, alors son nombre d’arêtes est au plus :*

$$\frac{p-2}{2(p-1)} \cdot n^2.$$

Turán a également obtenu une description des graphes avec un nombre maximal d’arêtes (à nombre de sommets fixé) parmi ceux qui ne contiennent pas un sous-graphe complet d’une taille donnée. Il faut noter que, contrairement au théorème de Kuratowski, le théorème de Turán donne une borne sur un invariant de graphes (ici, le nombre d’arêtes). C’est un autre type de théorèmes sur les classes de graphes excluant une sous-structure, que nous considérons dans cette thèse.

Nous présentons désormais un troisième type de théorèmes sur ces classes, qui décrivent comment leurs éléments peuvent être construits à partir de blocs de base. Dans la longue et fructueuse suite de papiers Graph Minors, Robertson et Seymour ont donné une description générale des graphes ne contenant pas un gros graphe complet comme mineur [RS03]. Ce théorème fournit une description structurelle de ces graphes : à quoi ils ressemblent et comment ils peuvent être décomposés. Informellement, Robertson et Seymour ont prouvé que ces graphes peuvent être obtenus en attachant ensemble sous forme arborescente des graphes qui peuvent être “presque” plongés dans une surface de genre borné. On ne donnera pas ici d’énoncé plus précis de ce théorème car cela nécessiterait d’introduire beaucoup trop de définitions. Mentionnons tout de même qu’outre sa valeur combinatoire, ce résultat est une pierre angulaire de la preuve du Théorème de Robertson et Seymour que nous rencontrerons d’ici quelques paragraphes.

Les trois théorèmes énoncés ci-dessus donnent différentes conclusions à propos des classes considérées : le premier fournit une information topologique, le second une majoration d’un invariant pour tous les graphes de la classe et le troisième la structure des graphes. Notre but en énonçant ces théorèmes est de présenter les types de résultats qu’on peut obtenir à propos des classes définies par exclusion de sous-structure.

Cette thèse est centrée autour des théorèmes d'exclusion. Nous présentons maintenant deux sujet connexes et qui recevront notre attention dans cette thèse : les beaux préordres et la propriété d'Erdős–Pósa.

## Beaux préordres

Revenons à notre exemple initial d'une collection d'objets dont certains sont plus désirables que d'autres. Supposons qu'on nous demande de choisir un objet dans cette collection. Quelles propriétés garantissent qu'un tel choix est toujours aisé? Si la collection contient plusieurs objets qui sont deux à deux incomparables, on peut choisir parmi eux en lançant un dé. Toutefois, cela n'est plus possible si ces objets incomparables sont en nombre infini, en supposant qu'on veut un choix équitable. Il est aussi possible que la collection contienne une séquence infinie d'objets de plus en plus désirables : dans ce cas on ne peut pas trouver un objet le plus désirable parmi eux (voir la [figure 4](#)).



Un nombre infini d'objets incomparables. Une séquence croissante infinie.

FIGURE 4 : Deux situations où le choix est difficile.

Cela nous amène à considérer des collections où ces deux comportements sont bannis. C'est plus ou moins la définition d'un beau préordre. Formellement, un beau préordre est un préordre (i.e. une relation binaire réflexive et transitive) qui ne contient pas de collection infinie d'éléments deux à deux incomparables, ni de séquence décroissante infinie<sup>4</sup>. Ce concept est une extension de la notion de bon ordre, qui s'applique usuellement à des ordres totaux (voir [Mos06, Chapitre 7] pour une introduction aux bons ordres), aux particularités des ordres partiels, où deux éléments ne sont pas toujours comparables. Dans un *bon ordre*, seules les séquences décroissantes infinies sont interdites, car il n'y a pas de collection d'éléments incomparables qui possède plus d'un élément. L'intérêt porté aux bons ordres en mathématiques est partiellement dû au fait qu'ils sont utilisés, souvent implicitement, dans plusieurs techniques de démonstration répandues, comme la récurrence (ou plus généralement l'induction), les démonstrations par contre-exemple minimal et la technique de descente infinie qui y est liée. Les bons ordres peuvent aussi être utilisés pour s'assurer qu'un programme termine (ou, similairement, qu'un système de réécriture termine) : si chaque état de l'exécution du programme est inférieure à l'état précédent pour un

<sup>4</sup>Bien que cette définition diffère légèrement de l'exemple donné ci-dessus où les séquences *croissantes* infinies sont interdites, c'est la même chose si on considère l'ordre dual où  $x$  est au moins  $y$  ssi  $y$  est au moins  $x$  dans l'ordre original.

bon ordre précautionneusement choisi, alors le programme terminera en un nombre fini d'étapes. Cela est facilité par l'utilisation de types de données récurrents (comme les listes et les arbres), qui sont souvent bonnement ordonnés [MP67, Bur69]. Les premières occurrences du concept de beau préordre sont apparues au début des années 1950 dans les publications de Higman [Hig52] et Erdős et Rado [ER52]. Avant cela, Vazsonyi avait conjecturé que toute séquence infinie d'arbres finis contient une paire d'arbres qui sont comparables par plongement homéomorphique, ce qui revient à demander s'ils forment un beau préordre. Un autre précurseur du concept de beau préordre peut être trouvé dans le travail de Kurepa, d'après [Kru72].

En théorie des graphes, il est courant de considérer des classes qui sont *closer* pour un certain ordre, c'est à dire où tout graphe qui est inférieur à un graphe de la classe en est aussi membre. Par exemple, la classe des graphes sans cycles est close pour la relation de sous-graphe, car aucun sous-graphe d'un graphe sans cycles ne contient de cycle. L'importance des beaux préordres en théorie des graphes et en algorithmique est due au fait suivant : dans un beau préordre, le complémentaire d'une classe close a un nombre fini d'éléments minimaux. Cela découle de la définition, car ces éléments étant minimaux, ils sont deux à deux incomparables. Ainsi, afin de vérifier si un élément  $x$  appartient à une classe close d'un beau préordre, il suffit de s'assurer que  $x$  n'est pas supérieur à  $y$ , pour un nombre fini d'objets  $y$  (les éléments minimaux du complémentaire). En d'autres termes, toute classe close d'un beau préordre peut être caractérisée par la négation d'un nombre fini d'inégalités.

Un des plus considérables résultats ayant trait aux beaux préordres de graphes est certainement le théorème de Robertson et Seymour déjà mentionné plus haut.

**Théorème 0.3** (Théorème de Robertson et Seymour, [RS04]). *La relation de mineur est un beau préordre sur les graphes finis.*

Ses conséquences, importantes en combinatoire et théorie des graphes, touchent aussi des domaines connexes comme l'algorithmique et ses liens avec la logique et la complexité computationnelle. La démonstration du théorème de Robertson et Seymour est longue et repose sur le théorème de structure de Robertson et Seymour. Pour les raisons mentionnées ci-dessus, un corollaire du **théorème 0.3** est que toute classe de graphe close par mineur peut être caractérisée par un nombre fini de mineurs exclus. D'une certaine manière, le théorème de Robertson et Seymour donne un théorème du même type que le théorème de Kuratowski pour toute classe qui est close par mineur. Il faut remarquer que la liste finie de graphes exclus n'est pas donnée par le théorème, qui ne stipule que son existence. Complété par un algorithme qui décide en temps polynomial si un graphe fixé est un mineur de l'entrée (également issu de Graph Minors [RS95]), le théorème de Robertson et Seymour implique aussi (encore une fois, purement existentiellement) que les problèmes de décision associés à des classes de graphes closes par mineur peuvent être résolus en temps polynomial. Le lecteur pourra se référer à [Lov06, Joh87] pour avoir une vue d'ensemble sur la théorie des mineurs et ses conséquences algorithmiques.

Les autres préordres usuels sur les graphes ne sont pas aussi généreux : il ne sont en général pas de beaux préordres. Ce fait est illustré par **figure 5** : aucun cycle est un sous-graphe d'un plus grand cycle. Cela nous conduit à poser la question d'identifier





FIGURE 5 : Les cycles forment un ensemble infini de graphes deux à deux incompatibles pour l'ordre de sous-graphe.

les classes où ces préordres sont beaux.

Dans cette thèse, nous répondons partiellement à cette question pour plusieurs préordres des graphes, ce qui complète les résultats existants. Un rôle de premier plan dans nos démonstrations est joué par des théorèmes de décomposition, qui décrivent la structure des graphes excluant une certaine sous-structure. En effet, de tels graphes ont souvent une forme ou une structure spécifique qui peut alors être utilisée pour les découper en objets plus simples et finalement les ordonner.

## La propriété d'Erdős–Pósa

Nous considérons maintenant un sujet apparemment indépendant et montrons ce qui le lie aux théorèmes d'exclusion. Pour commencer avec un exemple concret, considérons un jeu pour enfant (répandu par [Mod], voir également [GOB06]). Considérons une grille qui représente un jardin. Ce jardin est envahi par des rats, qui sont représentés par des formes identiques faites de cases adjacentes avec une orientation fixée. Le but du jeu est de protéger le jardin en plaçant des pièges (qui occupent une case chacun) de manière à ce que tout rat présent dans le jardin touche un piège. Un rat et un piège sont dessinés sur la figure 6.



Un rat




Un piège


FIGURE 6 : Rats et pièges.

Bien entendu, on pourrait mettre un piège sur chaque case, mais le but est d'utiliser le moins de pièges que possible. En guise d'exemple, la figure 7 présente deux situations : un jardin avec cinq rats (peut-on en avoir plus?) et un jardin protégé par neuf pièges (peut-on en utiliser moins?). Il faut noter que ce jeu a un grand nombre de variantes : chaque choix d'un jardin (qui peut être une plus grande grille ou une forme plus biscornue) et d'un rat donne un nouveau jeu.

Cet exemple nous permet de définir deux nombres : le premier est le nombre maximum  $p$  de rats qui peuvent habiter le jardin sans partager une case et le second



Une invasion de rats



Le jardin protégé par des pièges

FIGURE 7 : La chasse au rat.

est le nombre minimum  $c$  de pièges à même de protéger le jardin. Observons que  $c$  vaut toujours au moins  $p$ . En effet, si  $p$  rats sont présents dans le jardin, alors le protéger demande à placer, au minimum, un piège par rat. Mais il y a des situations où plus de  $p$  pièges sont nécessaires, ce qui soulève les questions suivantes :

- (1) Quelle est le lien entre  $c$  et  $p$ ? En particulier, peut-on majorer  $c$  par une fonction de  $p$ ?
- (2) Peut-on facilement calculer les valeurs de  $p$  et de  $c$ ?

Il n'est pas difficile de voir que, si on considère  $p$  rats dans le jardin, on peut protéger ce dernier en plaçant un piège sur chaque case occupée par un rat. Comme chaque rat occupe trois cases, on obtient  $c \leq 3p$ . Peut-on faire mieux? Une variante de ce jeu est de protéger le jardin de plusieurs espèces avec des formes différentes, par exemple rats et serpents, ces derniers étant définis comme suite de trois cases consécutives. On suppose que les pièges utilisés sont aussi efficaces contre les serpents. Le problème change : comme la [figure 8](#) le suggère, une disposition de pièges protégeant contre les rats ne protège pas toujours contre les serpents. Par ailleurs, le nombre maximum de rats et de serpents qui peuvent habiter le jardin est potentiellement plus grand que  $p$ .

De retour aux graphes, on peut définir des invariants similaires. Étant donné un graphe  $G$  et une classe de graphes  $\mathcal{H}$ , le *nombre de packing* de  $\mathcal{H}$  dans  $G$  est le nombre maximum de graphes de  $\mathcal{H}$  (répétitions permises) qui peuvent être trouvés dans  $G$  sans se superposer. Ici,  $G$  joue le rôle du jardin et  $\mathcal{H}$  celui de la liste des animaux indésirables. De même, le *nombre de couverture* de  $\mathcal{H}$  dans  $G$  est la taille minimale d'un ensemble de sommets (correspondant à des pièges) qui touchent chaque occurrence d'un graphe de  $\mathcal{H}$  dans  $G$ . Les nombres de packing et de couverture incarnent les nombres  $p$  et  $c$ , respectivement. Les mêmes questions que ci-dessus peuvent être posées dans ce contexte. Un exemple d'un théorème classique répondant à la question (1) est le théorème de Kőnig.




FIGURE 8 : Une solution pour rats n'est pas toujours une solution pour serpents.

**Théorème 0.4** (Theorème de König, 1931 [Kön31]). *Dans tout graphe biparti, le nombre maximal d'arêtes sommet-disjointes est égal au nombre minimal de sommets touchant toutes les arêtes.*

Ce théorème peut être reformulé comme suit : les nombres de packing et de couverture de  $\{K_2\}$  sont égaux dans les graphes bipartis. Pour une classe  $\mathcal{H}$  de graphes, il se peut que ces nombres ne soient pas égaux, mais que le nombre de couverture de  $\mathcal{H}$  soit majoré par une fonction de son nombre de packing. Dans ce cas, on dit que  $\mathcal{H}$  a la propriété d'Erdős-Pósa. Ce nom tire son origine du résultat suivant, prouvé dans les années 1960 par Erdős et Pósa.

**Théorème 0.5** (Théorème d'Erdős-Pósa, 1965 [EP65]). *Il existe une fonction  $f: \mathbb{N} \rightarrow \mathbb{N}$  telle que, pour tout graphe  $G$  et tout entier naturel  $k$ , soit  $G$  contient  $k$  cycles deux à deux sommet-disjointes, soit il a un ensemble de  $f(k)$  sommets dont la suppression rend le graphe acyclique.*

En d'autres termes, ce théorème révèle que la classe des cycles a la propriété d'Erdős-Pósa. Cependant, toutes les classes de graphes n'ont pas cette propriété. Par exemple, c'est le cas de la classe des cycles impairs, comme énoncé ci-dessous.

**Théorème 0.6** ([DL88, Ree99]). *Il existe une famille de graphes  $\{G_i\}_{i \in \mathbb{N}}$  telle que pour tout  $i \in \mathbb{N}$ , le nombre de packing des cycles impairs dans  $G_i$  est 1 tandis que le nombre de couverture est au moins  $i$ .*

Une illustration d'un graphe d'une telle famille est donnée dans la figure 9. Il a été démontré que ce graphe ne contient pas deux cycles impairs sommet-disjointes. Cependant, on peut facilement vérifier qu'en privant le graphe de trois sommets on garde toujours un cycle impair intact, quels que soient ces sommets.

Les résultats de la forme du Théorème d'Erdős-Pósa sont intéressants car ils connectent des invariants issus de problèmes apparemment orthogonaux. Ils nous invitent à explorer la dualité des problèmes de packing et de couverture. Mentionnons également




FIGURE 9 : Un *mur d'Escher* de hauteur 4.

qu'ils ont trouvé des applications en algorithmique (voir par exemple [FLMS12, Corollary 2] pour une preuve simple utilisant le théorème d'Erdős–Pósa) et en bioinformatique [ARS16, ADG04, Ara08]. Ainsi, de considérables efforts ont été déployés pour identifier les classes qui ont la propriété d'Erdős–Pósa, c'est à dire répondre à la question (1). Il se trouve que dans de nombreuses situations, une réponse à cette dernière peut être obtenue en considérant des deux cas suivants :

1. le graphe considéré a un nombre de packing grand ;
2. un certain invariant structurel du graphe (le plus souvent la largeur arborescente ou un paramètre "arborescent" alternatif) est majoré.

Le premier cas permet de conclure immédiatement, tandis que l'information structurelle que la majoration du second cas apporte permet de construire une petite couverture. Il faut toutefois s'assurer, pour appliquer ce schéma de preuve, qu'un de ces deux cas se présente pour les graphes considérés. Les théorèmes d'exclusion du type du Théorème de Turán nous fournissent une telle dichotomie : quand une sous-structure est absente du graphe, la majoration d'un certain invariant est donnée.

La question (2) peut elle aussi être posée dans le contexte des graphes. Cependant, pour de nombreuses classes de graphes, calculer les nombres de packing et de couverture est un problème NP-dur [GJ79]. Les deux options naturelles pour les attaquer sont la complexité paramétrée et les algorithmes d'approximation (voir par exemple [FLMS12, FLM<sup>+</sup>16]).

Dans cette thèse, nous continuons la longue lignée de recherche sur la propriété d'Erdős–Pósa en répondant à la question (1) pour diverses classes de graphes. Nous considérons également les problèmes algorithmiques de la question (2), pour lesquels nous concevons un algorithme d'approximation pour certaines classes de graphes. Ces algorithmes reposent sur des théorèmes d'exclusion, mais aussi sur le fait que les classes considérées ont la propriété d'Erdős–Pósa.

## Aperçu des résultats et organisation de la thèse

La plupart des définitions et notions utilisées dans cette thèse sont définies dans le [chapitre 2](#).

**Beaux préordres.** Le [chapitre 3](#) est dédié aux questions liées aux beaux préordres. Dans ce chapitre, nous présentons des outils généraux pour prouver des résultats sur les beaux préordres ([section 3.3](#)). Nous utilisons ensuite ces techniques ainsi que les antichaînes de la [section 3.5](#) pour démontrer trois résultats :

1. une caractérisation des classes de graphes définies par un mineur induit exclu qui sont bellement préordonnés par la relation de mineur induit, dans la [section 3.6](#);
2. une caractérisation similaire pour la relation de contraction, dans la [section 3.7](#);  
et
3. une caractérisation des classes de multigraphes qui sont closes par contraction et bellement préordonnées par la relation de contraction, dans la [section 3.8](#).

Ces trouvailles complètent les résultats existants sur ces préordres de graphes qui ne sont pas beaux en général. Elles sont obtenues via des théorèmes de décomposition pour les graphes excluant une sous-structure, pour la plupart prouvés dans les sections correspondantes.

**Théorèmes d'exclusion.** Dans le [chapitre 4](#) nous nous penchons sur les théorèmes d'exclusion qui majorent un invariant des graphes excluant une sous-structure. Nous obtenons :

1. une majoration d'un paramètre similaire à la maille des graphes excluant un grand graphe complet comme mineur, dans la [section 4.1](#);
2. une majoration du degré des graphes qui excluent une grande collection de multiarêtes comme mineur, dans [section 4.2](#);
3. une majoration (basse<sup>5</sup>) de la largeur arborescente des graphes excluant divers graphes planaires comme mineurs, dans [section 4.3](#); et
4. une majoration de la largeur de coupe arborescente des graphes excluant un graphe planaire sous-cubique comme immersion, dans [section 4.4](#).

La plupart de ces résultats sont utilisés dans les chapitres suivants.

---

<sup>5</sup>C'est à dire plus basse que la majoration donnée par les théorèmes généraux.

**La propriété d'Erdős-Pósa.** Le [chapitre 5](#) traite des liens entre les invariants de packing et de couverture. Il se découpe en trois parties :

1. une présentation de techniques générales pour prouver des résultats de type Erdős-Pósa, soit à partir de décompositions arborescentes, soit en utilisant des invariants comme la maille, [section 5.3](#) ;
2. une application de ces techniques à plusieurs classes de graphes dans la [section 5.4](#), le plus souvent en utilisant les théorèmes d'exclusion démontrés dans le [chapitre 4](#) ;
3. un résumé des résultats positifs et négatifs sur la propriété d'Erdős-Pósa, la [section 5.6](#).

**Applications algorithmiques.** Nous présentons dans le [chapitre 6](#) un algorithme d'approximation pour calculer les nombres de packing et de couverture de certaines classes de graphes. La démonstration définit une notion d'équivalence de graphes vis-à-vis du problème considéré, ce qui permet de réduire un graphe dès qu'une certaine sous-structure apparaît. En utilisant un résultat de la [section 4.1](#), on peut alors soit continuer à réduire le graphe, soit utiliser les arguments de la [section 5.3](#) pour conclure.

**Conclusion.** Nous concluons cette thèse par le [chapitre 7](#) qui revient sur les résultats présentés, énonce quelques problèmes ouverts et liste les publications obtenues durant la préparation de la thèse.



# CHAPTER 1

---

## INTRODUCTION

Imagine a (possibly infinite) collection of objects. In this collection, some objects may be more desirable than others. It might also happen that two objects are not comparable, that is, neither of them is more desirable than the other. An illustration is given in [Figure 1.1](#), where the ordered objects are fruits. In such a diagram, called a Hasse diagram, an element  $x$  is greater (for the considered order) than an element  $y$  if there is a top-bottom path from  $x$  to  $y$ . For instance, the diagram of [Figure 1.1](#) shows that a cherry is more desirable than a kiwi, because the path cherry-apricot-kiwi never goes up. Note that some fruits (like raspberry and mirabelle) are not comparable.




Figure 1.1: Some fruits ordered by desirability.

For a particular object  $x$  of the considered collection, let us consider all the objects that are not more desirable than  $x$ . This includes objects that are less desirable than  $x$ , but also those that are not comparable with it. The fruits of [Figure 1.1](#) that are underlined are those that are not more desirable than the apricot. A central theme in this thesis, for various collections of objects and notions of *desirability*, is the study of these objects that are not more desirable than  $x$ . How do they look like? What is their *structure*? What are their properties? Continuing the example of [Figure 1.1](#), a feature shared by all fruits that are not more desirable than the apricot is that they

have multiple seeds<sup>1</sup>.

The objects that we consider in this thesis are graphs and the desirability concept is expressed by several partial orderings of graphs: subgraphs, induced subgraphs, minors, topological minors, induced minors, immersions, etc., that most often express that a graph is *contained* as a substructure of an other. Figure 1.2 presents (isomorphism classes of) graphs that can be obtained by subdividing edges of the complete graph on five vertices, depicted at the bottom of the figure, ordered by the corresponding order. Naturally, this picture is incomplete as there are infinitely many such graphs.




Figure 1.2: Subdivisions of  $K_5$ .

One of the first theorems that established the properties of a class of graphs defined by forbidden substructures is the characterization of planar graphs by Kuratowski (see Figure 1.3 for a picture of the mentioned graphs).

**Theorem 1.1** (Kuratowski's Theorem, 1930 [Kur30]<sup>2</sup>). *A graph is planar<sup>3</sup> iff it does not contain a subgraph that is a subdivision of  $K_5$  or  $K_{3,3}$ .*

<sup>1</sup>A fruit in this diagram shows that the presence of multiple seeds is not a characterization of fruits that are not more desirable than the apricot (which one?).

<sup>2</sup>According to [Bur78], this result has been independently obtained by Pontryagin, but never published.

<sup>3</sup>We say that a graph is *planar* if it can be drawn on the plane without crossing edges.




Figure 1.3: Kuratowski's graphs  $K_5$  and  $K_{3,3}$  (from left to right).

A feature that made this result famous is that it describes a topological property, being planar, in purely combinatorial terms: the absence of a substructure in the graph. Often cited as a precursor of modern Graph Theory, Kuratowski's Theorem has been followed by a long line of results on classes defined by *excluding* a given substructure. One of them is Turán's Theorem, which gives an upper-bound on the number of edges of a graph not containing a large complete subgraph.

**Theorem 1.2** (Turán's Theorem, 1941 [Tur41]). *Let  $p \geq 2$  be an integer, and let  $G$  be a graph that does not contain a complete subgraph on  $p$  vertices. If  $G$  has  $n$  vertices, then its number of edges is at most:*

$$\frac{p-2}{2(p-1)} \cdot n^2.$$

Turán also provided a precise description of the graphs with the maximum number of edges that do not contain a complete subgraph of a given size. We should note that, unlike Kuratowski's Theorem, Turán's theorem provides a bound on a graph invariant (here, the number of edges). This is an other type of theorems on graphs excuding a substructure that we consider in this thesis. Let us now present a third type of such theorems, which describes how graphs of a class excuding a substructure can be constructed from basic blocks.

In the long and fruitful series of papers Graph Minors, Robertson and Seymour gave a general depiction of the graphs not containing a large complete graph as a minor [RS03]. This result, known as the Graph Minor Structure Theorem, provides a structural description of these graphs: what they look like and how they can be decomposed. Informally, they proved that these graphs look like graphs that are "almost embedded" in surfaces of bounded genus glued together in a tree-like fashion. We do not give a precise statement of this theorem here as it would require to introduce many more definitions. Let us mention that in addition to its combinatorial value, this result is a cornerstone of the proof of the Graph Minor Theorem [RS04] that we will meet again in a few paragraphs.

The three theorems stated above yielded different conclusions about the considered classes: the first one provides us with topological information, the second one upper-bounds an invariant for all graphs in this class, and with the third one we learn about the structure of the graphs. Our purposes in introducing these theorems are to present possible outcomes of results on classes defined by excluded substructures. Those considered in this thesis are that are bounds on invariants and structural decompositions of graphs.

This thesis is centered around exclusion theorems. We now present two related

topics that will receive attention in this thesis: well-quasi-orderings and the Erdős–Pósa property.

## Well-quasi-ordering

Let us go back to our initial example of a collection of objects, some of which are more desirable than others. Imagine now that we are asked to choose one object in this collection. What properties ensure that such a choice is always easy? If the collection contains several desirable objects that are pairwise not comparable, we can choose among them by rolling a dice. However this is not possible when these non-comparable objects are infinitely many, assuming that we want a fair choice. An other situation that may occur is when the collection contains an infinite sequence of objects that are ever more desirable: in this case we cannot find a most desirable object among them (see [Figure 1.4](#)).




Figure 1.4: Two situation where the choice is hard.

This leads us to consider collections where these two behaviors are forbidden. This is more or less the definition of a well-quasi-order. Formally, a *well-quasi-order* is a quasi-order (that is, a reflexive and transitive binary relation) that contains neither infinite collections of incomparable elements, nor infinite decreasing sequences<sup>4</sup>. This concept is an extension of that of a well-order, usually defined for total orders (see [\[Mos06, Chapter 7\]](#) for an introduction to well-ordering), to the setting of partial orders, where two elements are not always comparable. In a *well-order*, only infinite decreasing sequences are forbidden, as there is no collection of incomparable elements. The popularity of well-orders in mathematics is partly due to the fact that they are used (often implicitly) in several widely-spread proof techniques such as induction, proofs by minimal counterexample, and the related infinite descent technique. Well-orders can also be used to ensure program termination (or similarly, the termination of a rewriting system): if every state of the program execution is lower than its anterior step for some carefully chosen well-order, then the program will eventually terminate. This is facilitated by the use of recursively defined data types (as lists and trees), which are often well-ordered [\[MP67, Bur69\]](#). The first mentions of the concept of well-quasi-order appeared in the early 50's in papers of Higman [\[Hig52\]](#) and Erdős

<sup>4</sup>While this definition slightly differs from the example we give above where infinite *increasing* sequences are forbidden, it is quite the same if we consider the dual order where  $x$  is at least  $y$  iff  $y$  is at least  $x$  in the original order.

and Rado [ER52]. Prior to that, Vazsonyi conjectured that any infinite sequence of finite trees contains a pair of trees that are comparable under homeomorphic embedding, which is equivalent to well-quasi-ordering. An other forerunner of the concept of well-quasi-ordering can be found in the work of Kurepa, according to [Kru72].

In Graph Theory, one often deals with classes that are *closed* with respect to some ordering, i.e. every graph that is lower than a graph of the class also belongs to it. For instance, the class of acyclic graphs is closed with respect to the subgraph relation, because no subgraph of an acyclic graph contains a cycle. The importance of well-quasi-orders in Graph Theory and Algorithms is due to the following fact: in a well-quasi-order, the complementary of a closed class has finitely many minimal elements. This follows from the definition, as these elements, being minimal, are incomparable. Therefore, in order to decide if an object  $x$  belongs to a closed subclass of a well-quasi-order, one simply needs to check that  $x$  is not greater than  $y$ , for finitely many objects  $y$  (the minimal elements of the complementary). In other words, every closed subclass of a well-quasi-order can be characterized by a finite number of (negated) inequalities.

One of the most considerable results on well-quasi-ordering in Graph Theory is arguably the already mentioned Graph Minor Theorem of Robertson and Seymour.

**Theorem 1.3** (Graph Minor Theorem, [RS04]). *Finite graphs are well-quasi-ordered by the minor relation.*

The above theorem is considered as one of the most striking results in modern combinatorics. It had a strong impact, in Combinatorics and Graph Theory, but also in related fields as Algorithms and their connections to Logic and Computational Complexity. The proof of the Graph Minor Theorem is lengthy and relies on the Graph Minor Structure Theorem. For the aforementioned reasons, a corollary is that every class of graphs that is closed under taking of minors has a characterization in terms of finitely many excluded minors. In some sense, the Graph Minor Theorem provides a Kuratowski-type exclusion result for every class that is closed with respect to minors. It should be noted that the finite list of excluded minors is not given by the theorem, which only provides its existence. Together with an algorithm that decides if a fixed graph is a minor of the input graph in polynomial time (also originating from the Graph Minors series [RS95]), the Graph Minor Theorem also implies (again, purely existentially) that decision problems associated to classes that are closed under taking of minors can be solved in polynomial time. The reader is referred to [Lov06, Joh87] for an account on Minor Theory and its algorithmic consequences.

Other natural orderings of graphs are not so generous; they usually do not well-quasi-order all graphs. An illustration of this fact is given in [Figure 1.5](#): no cycle is a subgraph of a larger cycle. This raises the question of identifying the subclasses that are well-quasi-ordered.

In this thesis we provide a partial answer to this question for several orderings on graphs, complementing existing results. Essential parts of our proofs are decomposition theorems, which provide structural information about graphs excluding a certain substructure. Indeed, graphs in these classes often have a specific shape or




Figure 1.5: Cycles form an infinite set of graphs that are pairwise not comparable for the subgraph ordering.

structure, that can be used to dissect them into simpler objects, and eventually order them.

## The Erdős–Pósa property

We now move to a seemingly unrelated topic and show how it is connected to exclusion theorems. Let us start with a concrete example, which is a mathematical puzzle for children (popularized by [Mod], see also [GOB06]). We consider a grid that represents a garden. Imagine that this garden is invaded by rats, which are represented by identical shapes of adjacent cells with a fixed orientation. The goal of the game is to defend the garden, that is, to dispose traps (that cover one cell each) so that any rat in the garden would meet a trap. A rat and a trap are depicted in Figure 1.6.



A rat

A trap

Figure 1.6: Rats and traps.

Naturally, we could place one on every cell, but we aim at using as few traps as possible. Figure 1.7 depicts five rats in the garden (can we have more?) and an arrangement of traps that defend the garden (can we use less?). Observe that this puzzle has a wealth of variants: each choice of a garden (which could be a larger grid, or a more convoluted shape) and of a rat gives rise to a new problem.

This example allows us to define two numbers: the first one is the maximum number  $p$  of rats that fit in the garden without sharing a cell, and the second one is the minimum number  $c$  of traps required to defend the garden. Observe that  $c$  is always at least  $p$ . Indeed, if  $p$  rats are present in the garden, then for every rat, we need to place a trap on some of the cells it occupies. However, in some situations we may need more than  $p$  traps. This raises the following questions.

- (1) What is the relation between  $c$  and  $p$ ? In particular, can we bound  $c$  from above by a function of  $p$ ?




Figure 1.7: Hunting rats.

(2) Can we easily compute the values of  $p$  and  $c$ ?

It is easy to see that, if we consider  $p$  rats in the garden, placing a trap on each cell they occupy is enough to defend the garden. As a rat fills three cells, we get  $c \leq 3p$ . Can we do better? An variant of this puzzle is to protect the garden against several animals species with different shapes, for instance rats and snakes, where a snake is defined as 3 consecutive cells. We assume that our traps are efficient against snake as well. The problem then becomes different: as [Figure 1.8](#) suggests, a trap arrangement protecting against rats may not protect against snakes. Also, the maximum number of rats or snakes that fit in the garden is potentially larger than  $p$ .




Figure 1.8: A solution for rats is not always a solution for snakes.

Coming back to the setting of graphs, we can define similar invariants. Given a graph  $G$  and a class  $\mathcal{H}$ , the *packing number* of  $\mathcal{H}$  in  $G$  is the maximum number of graphs in  $\mathcal{H}$  (with repetitions allowed) that can be found in  $G$  without overlapping. Here,  $G$  plays the role of the garden, and  $\mathcal{H}$  that of the list of potential pests. Similarly, the *covering number* of  $\mathcal{H}$  in  $G$  is the minimum size of set of vertices (corresponding

to traps) that meet every occurrence of a graph of  $\mathcal{H}$  in  $G$ . Packing and covering numbers mirror the numbers  $p$  and  $c$ , respectively. The same questions as above can be asked in this setting. An example of a classic theorem answering question (1) is König's Theorem.

**Theorem 1.4** (König's Theorem, 1931 [Kön31]). *In a bipartite graph, the maximum number of pairwise vertex-disjoint edges is equal to the minimum number of vertices that meet all edges.*

This theorem can be restated as follows: the packing and covering numbers of  $\{K_2\}$  are equal in bipartite graphs. For some class  $\mathcal{H}$  of graphs, it may happen that these numbers are not equal, but that the covering number of  $\mathcal{H}$  is bounded from above by a function of its packing number. In this case,  $\mathcal{H}$  is said to have the *Erdős–Pósa property*. This name originates from the next result, proved in 60's by Erdős and Pósa.

**Theorem 1.5** (Erdős–Pósa Theorem, 1965 [EP65]). *There is a function  $f: \mathbb{N} \rightarrow \mathbb{N}$  such that for every graph  $G$  and every positive integer  $k$ , either  $G$  has  $k$  pairwise vertex-disjoint cycles, or there is a set of  $f(k)$  vertices, the removal of which yields an acyclic graph.*

In other words, this theorem states that the class of cycles have the Erdős–Pósa property. However, there are classes of graphs that do not have the Erdős–Pósa property. One of them is the class of cycles of odd length, as stated below.

**Theorem 1.6** ([DL88, Ree99]). *There is a family  $\{G_i\}_{i \in \mathbb{N}}$  of graphs such that for every  $i \in \mathbb{N}$ , the packing number of odd cycles in  $G_i$  is one and the covering number is at least  $i$ .*

An illustration of a graph in this family is provided by Figure 1.9. It has been proved that this graph does not contain two vertex-disjoint odd cycles. However one can easily check that removing any three vertices keeps some odd cycle intact.




Figure 1.9: An *Escher wall* of height 4.

Erdős–Pósa-type results are interesting as they link together invariants related to apparently orthogonal problems. They invite us to investigate the duality between

packing and covering problems. Let us also briefly mention that they have been used in algorithm design (see e.g. [FLMS12, Corollary 2] for a simple proof using the Erdős–Pósa Theorem) and in bioinformatics [ARS16, ADG04, Ara08]. Therefore, considerable effort has been put in identifying classes that have the Erdős–Pósa property, i.e. answering question (1). It appears that in several situations, a proof of (1) can be obtained by considering the following two cases:

1. the considered graph has a large packing number; or
2. some structural invariant of the graph (most often the treewidth or some alternative “tree-structure” parameter) is bounded.

From the first case we can directly conclude, whereas the bound provided by the second case usually yields structural information on the graph that indicates how to construct a small cover. However, to apply this proof scheme, we first need to ensure that at least one of these two cases is true for the considered graphs. Such a *dichotomy* is provided by exclusion theorems of the type of Turán’s Theorem: in the case where a substructure is absent from the graph, they provide a bound on some parameter.

Question (2) can be asked in the setting of graphs as well. However, it appears that for several graph classes, computing packing and covering numbers is an NP-hard problem [GJ79]. The two natural options to attack them are thus parameterized algorithms and approximation (see e.g. [FLMS12, FLM<sup>+</sup>16]).

In this thesis we continue the long line of research on the Erdős–Pósa property by answering (1) for several classes of graphs. We also consider the problems of (2), for which we design an approximation algorithm on certain classes of graphs. These algorithms rely on exclusion theorems, but also on the fact that the considered classes have the Erdős–Pósa property.

## Overview of the results and organization of the thesis

Most of the definitions and notions used in this thesis are defined on [Chapter 2](#).

**Well-quasi-orderings.** [Chapter 3](#) is devoted to questions related to well-quasi-orderings. In this chapter, we present general tools to prove well-quasi-ordering results ([Section 3.3](#)). We then use these techniques together with antichains introduced in [Section 3.5](#) to show three results on well-quasi-ordering:

1. a characterization of the classes of simple graphs defined by one forbidden induced minor that are well-quasi-ordered with respect to this relation, in [Section 3.6](#);
2. a similar characterization for the relation of contraction, in [Section 3.7](#); and
3. a characterization of the contraction-closed classes of multigraphs that are well-quasi-ordered by the multigraph contraction relation, in [Section 3.8](#).

These results complement the similar characterizations known for most of the usual orderings of graphs that are not well-quasi-orders in general. They are obtained using decomposition theorems for graphs excluding a substructure, most of which are proved in the corresponding sections.

**Exclusion theorems.** Chapter 4 is concerned with exclusion theorems that upper-bounds a parameter of the graphs excluding a substructure. We obtain:

1. bounds on a girth-like parameter of graphs excluding large complete graphs as a minor, in Section 4.1;
2. bounds on the degree of graphs that exclude a large collection of multiedges as a minor, in Section 4.2;
3. (low<sup>5</sup>) bounds on the treewidth of graphs excluding various planar graphs as minors, in Section 4.3; and
4. bounds on the tree-cut width of graphs excluding a planar subcubic graph as an immersion, in Section 4.4.

Most of these results are used in the following chapters.

**The Erdős–Pósa property.** Chapter 5 deals with connections between invariants of packing and covering. It is split into three parts:

1. general techniques for proving Erdős–Pósa-type results, either from tree-like decompositions of graphs, or using invariants as the girth, in Section 5.3;
2. an application of these techniques to several classes of graphs in Section 5.4, most often using exclusion results proved in Chapter 4;
3. a summary of positive and negative results on the Erdős–Pósa property, in Section 5.6.

**Algorithmic applications.** We complement the chapter on the Erdős–Pósa property by presenting in Chapter 6 an approximation algorithm for packing and covering numbers of certain classes of graphs, relying on exclusion theorems proved Section 4.1 and on the properties of the considered classes. The proof is using a notion of equivalence of graphs with respect to the considered problem which allows us to reduce the graph when some structure appears. Using a result of Section 4.1, we can then either reduce further the graph, or use the arguments presented in Section 5.3 to conclude.

**Conclusion.** We summarize in Chapter 7 the contributions of this thesis. We also present directions for future research and list the publications written during the preparation of the thesis.

---

<sup>5</sup>I.e. lower than the bounds provided by the general theorems.

# CHAPTER 2

---

## GENERAL DEFINITIONS

Let us introduce notions that will be used all along this thesis. Specific notions will be defined in the corresponding chapters. The reader may refer to the [index](#) (on page 246) in order to find where a given notion or symbol has been defined.

### 2.1 Basics

In this thesis, logarithms are binary. If  $\mathcal{S}$  is a set of sets, then  $\mathbf{US} = \bigcup_{S \in \mathcal{S}} S$ . For every integers  $i$  and  $j$ , the notation  $\llbracket i, j \rrbracket$  stands for the interval of integers  $\{i, \dots, j\}$ . For every positive integer  $k$ , we denote by  $\mathbb{N}_{\geq k}$  the set  $\mathbb{N} \setminus \llbracket 0, k-1 \rrbracket$ . We denote by  $\mathcal{P}(S)$  the power set of a set  $S$  and by  $\mathcal{P}^{<\omega}(S)$  the set of all its finite subsets. Given a function  $\phi : A \rightarrow B$  and a subset  $C \subseteq A$ , we define  $\phi(C) = \{\phi(x) \mid x \in C\}$ . Let  $\mathbf{t} = (x_1, \dots, x_l) \in \mathbb{N}^l$  and  $\chi, \psi : \mathbb{N} \rightarrow \mathbb{N}$ . We say that  $\chi(n) = O_{\mathbf{t}}(\psi(n))$  if there exists a computable function  $\phi : \mathbb{N}^l \rightarrow \mathbb{N}$  such that  $\chi(n) = O(\phi(\mathbf{t}) \cdot \psi(n))$ . We set  $\mathbb{N}^+ = \mathbb{N} \setminus \{0\}$ .

By  $\text{polylog}(t)$  we denote some function that is a polynomial in the logarithm of  $t$ . More formally, we write  $f(t) = O(\text{polylog } t)$  if there are constants  $t_0, A, \alpha$  such that  $\forall t > t_0, f(t) \leq A \log^\alpha(t)$ .

### 2.2 Orders

In this section we introduce basic definitions that are related to order theory.

**Sequences.** A *sequence* of elements of a set  $A$  is an ordered countable collection of elements of  $A$ . Unless otherwise stated, sequences are finite. The sequence of elements  $s_1, \dots, s_k \in A$  in this order is denoted by  $\langle s_1, \dots, s_k \rangle$ . We use the notation  $A^*$  for the class of all finite sequences over  $A$  (including the empty sequence). The length of a finite sequence  $s \in A^*$  is denoted by  $|s|$ .

**Ordered sets.** A *quasi-order* (qoset for short) is a pair  $(A, \preceq)$  where  $A$  is a set and  $\preceq$  is a binary relation on  $A$  that is reflexive and transitive.

Two elements  $x$  and  $y$  of a qoset  $(A, \preceq)$  are said to be *non-comparable* if none of  $x \preceq y$  and  $y \preceq x$  holds. In a qoset, an *antichain* is a sequence of pairwise non-comparable elements. In a sequence  $\langle x_i \rangle_{i \in I \subseteq \mathbb{N}}$  of a qoset  $(A, \preceq)$ , a pair  $(x_i, x_j)$ ,  $i, j \in I$  is a *good pair* if  $x_i \preceq x_j$  and  $i < j$ . A qoset  $(A, \preceq)$  is a *well-quasi-order* (*wqo* for short), and  $A$  is said to be *well-quasi-ordered* by  $\preceq$ , if every infinite sequence has a good pair, or equivalently, if  $(A, \preceq)$  has neither an infinite decreasing sequence, nor an infinite antichain. An infinite sequence containing no good pair is called a *bad sequence*.

Other notions related to orders are defined either in [Section 2.4](#) (graph orderings) or in [Section 3.2](#) (notions specific to the chapter on well-quasi-ordering).

## 2.3 Graphs

### 2.3.1 Basics

A *graph*  $G$  is a pair  $(V, E)$ , where  $V$  is a set referred to as the *set of vertices* of  $G$  and  $E$  is a multiset, the underlying set of which is a subset of  $\binom{V}{2}$ , that we call the *set of edges* of  $G$  (even if it is a multiset). We denote by  $V(G)$  the vertex set of the graph  $G$  and by  $E(G)$  its edge set. In this thesis, we do not consider graphs with loops, i.e. edges connecting a vertex to itself. This definition corresponds to what is sometimes called a “loopless multigraph”. The *order* of a graph  $G$  is its number of vertices, that we denote by  $n(G)$ , whereas we use  $m(G)$  for its number of edges (counting multiplicities). We sometimes use the notation  $|G| = |V(G)|$  and  $\|G\| = |E(G)|$  (counting multiplicities). We denote by  $\text{mult}_G(\{u, v\})$  the function that gives the multiplicity in  $G$  of a given edge (which might be zero if  $\{u, v\}$  does not belong to  $E$ ). We drop the subscript when it is clear from the context. To refer to a particular edge of a multiedge  $\{u, v\}$ , we use the subscript  $\{u, v\}_i$ , where  $i \in \llbracket 1, \text{mult}_G(\{u, v\}) \rrbracket$ .

We also deal with *simple graphs*, which are graphs where every edge has multiplicity one. We use the notations  $V(G)$  and  $E(G)$  for the vertex and edge sets of a graph, respectively. The *underlying simple graph* of a graph  $G$  is the graph  $G'$  such that  $V(G') = V(G)$  and  $E(G')$  is the underlying set of  $E(G)$ . All the graphs we consider in this thesis are finite, i.e. both  $V(G)$  and the values taken by  $\text{mult}_G$  are finite. For  $H$  and  $G$  graphs, we write  $H + G$  for the disjoint union of  $H$  and  $G$ . Also, for every  $k \in \mathbb{N}$ ,  $k \cdot G$  is the disjoint union of  $k$  copies of  $G$ . The *Cartesian product*  $H \times G$  of  $H$  and  $G$  is the graph on vertex set  $V(H) \times V(G)$  and where, given two vertices  $(u, v), (u', v') \in V(H) \times V(G)$ , either  $u = u'$  and  $\text{mult}_{H \times G}(\{(u, v), (u', v')\}) = \text{mult}_G(\{v, v'\})$ , or  $v = v'$  and  $\text{mult}_{H \times G}(\{(u, v), (u', v')\}) = \text{mult}_H(\{u, u'\})$ .

For every  $X \subseteq V(G)$ , the subgraph *induced* by  $X$ , that we write  $G[X]$ , is the subgraph with vertex set  $X$  and edge set  $\{e \in E(G), e \subseteq X\}$ . We denote by  $G \setminus X$  the subgraph of  $G$  induced by  $V(G) \setminus X$ . If  $X \subseteq E(G)$ , then  $G \setminus X$  is the subgraph  $(V(G), E(G) \setminus X)$ . The complement of a simple graph  $G$ , denoted  $\overline{G}$  is the graph obtained by replacing every edge by a non-edge and vice-versa in  $G$ . Given a graph class  $\mathcal{C}$  and a graph  $G$ , we call  $\mathcal{C}$ -*subgraph* of  $G$  any subgraph of  $G$  that is isomorphic to some graph in  $\mathcal{C}$ . If the class of  $\mathcal{C}$ -subgraphs of  $G$  is empty, then we say that  $G$  is  $\mathcal{C}$ -*free*.

**Neighbors and degree.** Two vertices  $u, v \in V(G)$  are said to be *adjacent* if  $\text{mult}(\{u, v\}) \geq 1$ . An edge  $e \in E(G)$  is *incident* to a vertex  $v \in V(G)$  if  $v \in e$ . Two edges are *incident* if they share some endpoint. The *neighborhood* of a vertex  $v \in V(G)$ , denoted  $N_G(v)$ , is the set of all vertices of  $G$  that are adjacent to  $v$ . For every subset  $S \subseteq V(G)$ , we set  $N_G(S) = \bigcup_{v \in S} N_G(v) \setminus S$  (all vertices of  $V(G) \setminus S$  that have a neighbor in  $S$ ). We extend the definition of  $\text{mult}_G$  as follows, for every subsets  $X, Y \subseteq V(G)$ :

$$\text{mult}_G(X, Y) = \sum_{(x,y) \in X \times Y} \text{mult}_G(\{x, y\}).$$

If  $\text{mult}_G(X, Y) \geq 1$ , we say that  $X$  and  $Y$  are *adjacent* in  $G$ .

The *degree*  $\deg_G(v)$  of a vertex  $v \in V(G)$  is the cardinality of  $N_G(v)$ , i.e. the number of vertices of  $G$  that are adjacent to  $v$ . On the other hand, the *multidegree*  $\text{mdeg}_G(v)$  of  $v$  is defined as the cardinality of  $\text{mult}_G(\{v\}, N_G(v))$ , i.e. the number of edges incident with  $v$ . Observe that these values are equal in simple graphs. We drop the subscript when it is obvious from the context. The minimum degree over all vertices of a graph  $G$  is denoted by  $\delta(G)$ , and the maximum degree by  $\Delta(G)$ .

**Distances.** For a given graph  $G$  and two vertices  $u, v \in V(G)$ ,  $\text{dist}_G(u, v)$  denotes the *distance between  $u$  and  $v$* , which is the number of edges on a shortest path between  $u$  and  $v$ , and  $\text{diam}(G)$  denotes  $\max\{\text{dist}_G(u, v) \mid u, v \in V(G)\}$ . For a set  $S \subseteq V(G)$  and a vertex  $w \in V$ ,  $\text{dist}_G(S, w)$  denotes  $\min\{\text{dist}_G(v, w) \mid v \in S\}$ . Also, for a given vertex  $u \in V(G)$ ,  $\text{ecc}_G(u)$  denotes the *eccentricity* of vertex  $v$ , that is,  $\max\{\text{dist}_G(u, v) \mid v \in V(G)\}$ . The *girth* of a graph  $G$ , denoted  $\text{girth}(G)$ , is the length of a shortest cycle in  $G$  if  $G$  is not a forest, and  $\infty$  otherwise.

**Connectivity and separations.** A graph  $G$  is *connected* if, for every  $x, y \in V(G)$ , there is a path starting in  $x$  and ending in  $y$ . A pair  $(A, B)$  of subsets of  $V(G)$  is called a *separation of order  $k$  in  $G$*  if  $A \cup B \subseteq V(G)$ ,  $k = |A \cap B|$ , none of  $A, B$  is a subset of the other, and there is no edge of  $G$  between  $A \setminus B$  and  $B \setminus A$ .

For every  $k \in \mathbb{N}$ ,  $k \geq 2$ , we say that  $G$  is  *$k$ -connected* if there is no separation of order less than  $k$  in  $G$ . A *connected component* of a graph is a maximal connected subgraph. We denote by  $\text{cc}(G)$  the number of connected component of a graph  $G$ . We say that a subgraph is a *2-connected component* is a maximal 2-connected subgraph. A maximal subgraph with no cutvertex is called a *block*. A *cut*  $C = (X, Y)$  of  $G$  is a partition of  $V(G)$  into two subsets  $X$  and  $Y$ . The *cut-set* of  $C$  is  $E(X, Y)$  and the *width* of the cut is  $|E(X, Y)|$ .

**Trees, paths, and cycles.** A *tree* is a connected acyclic simple graph. A *leaf* in a tree is a vertex of degree one.

For every two vertices  $u, v \in V(T)$ , there is exactly one path in  $T$  between  $u$  and  $v$ , that we denote by  $uTv$ . Also, given that  $uTv$  has at least 2 vertices, we denote by  $\overset{!}{u}Tv$  (respectively  $u\overset{!}{v}$ ) the path  $uTv$  with the vertex  $u$  (respectively  $v$ ) deleted. Let  $C$  be a cycle on which we fixed some orientation. Then, there is exactly one path following

this orientation between any two vertices  $u, v \in V(C)$ . Similarly, we denote this path by  $uCv$  and we define  $\overset{\circ}{u}Cv$  and  $uC\overset{\circ}{v}$  as we did for the tree. Two paths in a graph are *internally disjoint* if they do not share any internal vertex.

**Rooted trees.** A *rooted tree* is a pair  $(T, r)$ , where  $T$  is a tree and  $v \in V(T)$  is a distinguished vertex called *root*. Let  $(T, s)$  be a rooted tree. Given a vertex  $x \in V(T)$ , the *descendants* of  $x$  in  $(T, s)$ , denoted by  $\text{desc}_{(T,s)}(x)$ , is the set containing each vertex  $w$  such that the unique path from  $w$  to  $s$  in  $T$  contains  $x$ . Given a rooted tree  $(T, s)$  and a vertex  $x \in V(T)$ , the *height* of  $x$  in  $(T, s)$  is the maximum distance between  $x$  and a vertex in  $\text{desc}_{(T,s)}(x)$ . The *height* of  $(T, s)$  is the height of  $s$  in  $(T, s)$ . The *children* of a vertex  $x \in V(T)$  are the vertices in  $\text{desc}_{(T,s)}(x)$  that are adjacent to  $x$ . A *leaf* of  $(T, s)$  is a vertex of  $T$  without children. The *parent* of a vertex  $x \in V(T) \setminus \{s\}$ , denoted by  $\text{p}(x)$ , is the unique vertex of  $T$  that has  $x$  as a child. In a rooted tree  $T$  with root  $r$ , the *least common ancestor* of two vertices  $u$  and  $v$ , written  $\text{lca}^T(u, v)$ , is the first common vertex of the paths  $uTr$  and  $vTr$ .

### 2.3.2 Special graphs and graph classes

In this section we define specific graphs that will appear all along this thesis. Let  $n$  be a positive integer. We denote by:

- $K_n$  the complete graph on  $n$  vertices;
- $P_n$  the path on  $n$  vertices;
- $C_n$  the cycle on  $n$  vertices (when  $n \geq 2$ );
- $\theta_n$  the graph obtained by connecting two vertices with an edge of multiplicity  $n$ , sometime called the *n-pumpkin* (see [Figure 2.1](#) for an example);
- $D_n$  the graph  $\overline{2 \cdot K_1 \cup K_n}$  (idem), and  $D_2$  is the graph sometimes called *diamond*.



Figure 2.1: The graphs  $\theta_5$  (left) and  $D_4$  (right).

We call *prism* the Cartesian product of  $K_3$  and  $K_2$ . A *cograph* is a graph not containing the path on four vertices as induced subgraph. A *linear forest* is a disjoint union of paths. A graph is *subcubic* its maximum degree is upper-bounded by 3.

**Grids and Walls.** Let  $k$  and  $r$  be positive integers where  $k, r \geq 2$ . The  $(k \times r)$ -grid  $\Gamma_{k,r}$  is the Cartesian product of two paths of lengths  $k-1$  and  $r-1$  respectively. We denote by  $\Gamma_k$  the  $(k \times k)$ -grid. The  $k$ -wall  $W_k$  is the graph obtained from a  $((k+1) \times (2 \cdot k+2))$ -grid with vertices  $(x, y), x \in \{1, \dots, k+1\}, y \in \{1, \dots, 2k+2\}$ , after the removal of the “vertical” edges  $\{(x, y), (x, y+1)\}$  for odd  $x+y$ , and then the removal of all vertices of degree 1. The graphs  $\Gamma_4$  and  $W_4$  are depicted on [Figure 2.2](#).

Let  $W_k$  be a wall. We denote by  $P_j^{(v)}$  the shortest path connecting vertices  $(1, 2j)$  and  $(k+1, 2j), j \in [k]$  and call these paths the *vertical paths* of  $W_k$ , with the assumption that  $P_j^{(v)}$  contains only vertices  $(x, y)$  with  $y = 2j, 2j-1$ . Note that these paths are vertex-disjoint. Similarly, for every  $i \in [k+1]$  we denote by  $P_i^{(h)}$  the shortest path connecting vertices  $(i, 1)$  and  $(i, 2k+2)$  (or  $(i, 2k+1)$  if  $(i, 2k+2)$  has been removed) and call these paths the *horizontal paths* of  $W_k$ .




Figure 2.2: The  $(4 \times 4)$ -grid (left) and the 4-wall (right).

**Wheels.** For every positive integer  $n$ , a  $n$ -wheel, also called *wheel of order  $n$* , is a simple graph obtained by connecting a (new) vertex to  $n$  distinct vertices of an induced cycle  $C$ . This cycle is said to be the *cycle* of the  $n$ -wheel, whereas the new vertex is its *center*. A *double wheel of order  $n$*  is obtained from a cycle of order  $n$  by adding two non-adjacent vertices, each connected to the same  $n$  vertices of the cycle. See [Figure 2.3](#) for examples.




Figure 2.3: A wheel of order six (left) and a double wheel of order 6 (right).

**Yurts.** For every integer  $n > 0$ , the *yurt graph* of order  $n$  the graph  $Y_n$  of the form

$$\begin{aligned} V(Y_n) &= \{x_1, \dots, x_n, y_1, \dots, y_n, o\} \\ E(Y_n) &= \left\{ \{x_i, x_{i+1}\}_{i \in \llbracket 1, n-1 \rrbracket} \right\} \\ &\quad \cup \left\{ \{y_i, y_{i+1}\}_{i \in \llbracket 1, n-1 \rrbracket} \right\} \\ &\quad \cup \left\{ \{x_i, y_i\}_{i \in \llbracket 1, n \rrbracket} \right\} \\ &\quad \cup \left\{ \{y_i, o\}_{i \in \llbracket 1, n \rrbracket} \right\} \end{aligned}$$

(see [Figure 2.4](#) for an example).




Figure 2.4: The yurt graph of order 5.

## 2.4 Graph operations and orderings

Most of the common order relations on graphs can be defined in two equivalent ways: either in terms of *graph operations*, or by using *models*. Let us have a closer look at them.

### 2.4.1 Local operations

Let  $G$  be a graph. We here describe the effects of the following local operations when applied to  $G$ : the vertex deletion, the edge deletion, the vertex dissolution, the edge contraction, the vertex identification and the lift.

*Deleting a vertex  $v$*  (respectively an edge  $e$ ) yields the graph  $G \setminus \{v\}$  (respectively  $G \setminus \{e\}$ ). For every  $\{u, v\} \in E(G)$ , the *contraction* of the edge  $\{u, v\}$  adds a new vertex  $w$ , sets  $\text{mult}(\{w, w'\}) = \text{mult}(\{u, w'\}) + \text{mult}(\{v, w'\})$  for every  $w \in N(u) \cup N(v)$  and then deletes  $u$  and  $v$  (see [Figure 2.5](#)). If  $G$  is a simple graph, we do not keep the multiple edges that might be created during this process (i.e. we set their multiplicity to one). In the case where  $G$  is labeled, we label the new vertex  $w$  in the obtained graph with  $\lambda_G(u) \cup \lambda_G(v)$ .




Figure 2.5: The contraction of the edge  $e$  creates a double edge.

If  $u$  is a vertex of degree two, the *dissolution* of  $u$  is the contraction of one edge incident with  $u$ . On the other hand, a *subdivision* of the edge  $\{u, v\}$  adds a new vertex adjacent to  $u$  and  $v$  and decreases the multiplicity of the edge  $\{u, v\}$  by one (i.e., removes this edge in the case where  $G$  is a simple graph). We delete edges of multiplicity 0. The vertex added during this process is called a *subdivision vertex*. These two operations are depicted on [Figure 2.6](#).




Figure 2.6: Dissolution and subdivision as complementary operations.

The *identification* of two vertices  $u$  and  $v$  adds the edge  $\{u, v\}$  if it was not already existing, and contracts it. If  $G$  is  $(\Sigma, \preceq)$ -labeled (for some qoset  $(\Sigma, \preceq)$ ), a *label contraction* is the operation of relabeling a vertex  $v \in V(G)$  with a label  $l$  such that  $l \preceq^{\mathcal{P}} \lambda_G(v)$ . This operation will be used when dealing with well-quasi-orders in [Chapter 3](#). The *lift* of two incident edges  $\{u, v\}$  and  $\{v, w\}$  decreases by one the multiplicities of these edges and increases by one the multiplicity of  $\{u, w\}$  (or create the edge if it was not existing).

The *closure* of a class  $\mathcal{G}$  by a given operation is the class obtained from graphs of  $\mathcal{G}$  by a finite application of this operation.

## 2.4.2 Containment models

In this section we define models, which are functions witnessing the presence of a substructure in a graph. They come in different flavours, depending on the type of substructure considered.

**Containment models.** Let  $G$  and  $H$  be two graphs and let us consider a function  $\mu: V(H) \rightarrow \mathcal{P}^{<\omega}(V(G))$  and the following properties:

- (M1): for every two distinct  $u, v \in V(H)$ , the sets  $\mu(u)$  and  $\mu(v)$  are vertex-disjoint;
- (M2): for every  $u \in V(H)$ , the subgraph of  $G$  induced by  $\mu(u)$  is connected;
- (M3):  $\lambda_H(u) \preceq^{\mathcal{P}} \bigcup_{v \in \mu(u)} \lambda_G(v)$ ;
- (M4):  $\text{root}(G) \in \mu(\text{root}(H))$ ;
- (M5):  $\forall u, v \in V(H)$ ,  $\text{mult}_G(\mu(u), \mu(v)) \geq \text{mult}_H(\{u, v\})$ ;
- (M6):  $\forall u, v \in V(H)$ ,  $\text{mult}_G(\mu(u), \mu(v)) = \text{mult}_H(\{u, v\})$ ;
- (M7):  $\bigcup_{v \in V(H)} \mu(v) = V(G)$ ;
- (M8):  $\forall v \in V(H)$ ,  $|\mu(v)| = 1$ .

...-model	(M1)	(M2)	(M3)	(M4)	(M5)	(M6)	(M7)	(M8)
minor	✓	✓	(✓)	(✓)	✓			
induced minor	✓	✓	(✓)	(✓)	✓	(✓)		
contraction	✓	✓	(✓)	(✓)	✓	(✓)	✓	
subgraph	✓	✓	(✓)	(✓)	✓		✓	✓
induced subgraph	✓	✓	(✓)	(✓)	✓	✓	✓	✓

Table 2.1: Requirements for containment models

If  $\mu$  satisfies (M1) up to (M5), then we call it an  $H$ -minor model in  $G$ , or a *minor model of  $H$* . A minor model that also satisfies (M6) is an *induced minor model*, and an induced minor model where (M7) holds is a *contraction model*. A minor model that additionally satisfies (M7) and (M8) is a *subgraph model*; it is an *induced subgraph model* if it satisfies (M1) up to (M8). These definitions are summarized by Table 2.1. Items (M3) and (M4) are required only when  $H$  and  $G$  are labeled or rooted, respectively. We do not require (M6) for simple graphs. An example of a minor model is given in Figure 2.7. When (M3) holds,  $\mu$  is said to be *label-preserving*.




Figure 2.7: A  $K_4$ -minor model (dashed arrows) in the  $3 \times 3$  grid, that is also a topological minor model and an induced minor model.

**Models for immersions and topological minors.** In order to deal with the orderings of immersion and topological minor, to be defined in Subsection 2.4.3, we need a different kind of model. An  $H$ -immersion model is a pair of functions  $(\phi, \psi)$  satisfying the following properties:

1.  $\phi: V(H) \rightarrow V(G)$  is an injection;
2.  $\psi$  sends  $\{u, v\}_i$  to a path of  $G$  between  $\phi(u)$  and  $\phi(v)$ , for every  $\{u, v\} \in E(H)$  and every  $i \in \llbracket 1, \text{mult}_H(\{u, v\}) \rrbracket$ , in a way such that distinct edges are sent to edge-disjoint paths.

Every vertex in the image of  $\phi$  is called a *branch vertex* and every path in  $\psi(E(H))$  a *certifying path*. If it also holds that no branch vertex is an internal vertex of any certifying path, then the function  $(\phi, \psi)$  is an  $H$ -strong-immersion model. If moreover,

the paths in the image of  $\psi$  are internally disjoint, then  $(\phi, \psi)$  is an  $H$ -topological minor model.

With these definitions the following observation is straightforward.

*Observation 2.1.* Let  $H$  and  $G$  be graphs. If  $(\phi, \psi)$  is an  $H$ -topological-minor model in  $G$  then  $(\phi, \psi)$  is also an  $H$ -strong-immersion model in  $G$ .




Figure 2.8: A  $K_4$ -immersion model in a graph that has no  $K_4$ -minor model: vertices are sent on vertices with the same name, and edges are sent to paths of the same color.

The next section links together models and local operations in the definition of graph orderings.

### 2.4.3 Graph orderings

Local operations and models can be used to express that a graph is *contained* as a substructure of an other one. The graph ordering where this notion of *containment* is the most obvious is perhaps the subgraph relation which can be defined as follows:  $H$  is a *subgraph* of  $G$  if  $V(H) \subseteq V(G)$  and  $E(H) \subseteq E(G)$ .

The orderings we consider here can be defined in two equivalent ways: either using models, of using local operations. Let us start with the definition involving models.

- $H$  is a *subgraph* of  $G$ , what we note  $H \leq_{\text{sg.}} G$ , if there is an  $H$ -subgraph model in  $G$ ;
- $H$  is an *induced subgraph* of  $G$ , what we note  $H \leq_{\text{i.sg.}} G$ , if there is an  $H$ -induced subgraph model in  $G$ ;
- $H$  is a *contraction* of  $G$ , what we note  $H \leq_{\text{c.}} G$  when these graphs are simple and  $H \leq_{\text{m.c.}} G$  when they may have multiple edges<sup>1</sup>, if there is an  $H$ -contraction model in  $G$ ;
- $H$  is a *minor* of  $G$ , what we note  $H \leq_{\text{m.}} G$ , if there is an  $H$ -minor model in  $G$ ;
- $H$  is an *induced minor* of  $G$ , what we note  $H \leq_{\text{i.m.}} G$ , if there is an  $H$ -induced minor model in  $G$ ;

<sup>1</sup>As we will consider these two settings, we choose to use two different symbols to stress that the corresponding order relations are different.

- $H$  is a *topological minor* of  $G$ , what we note  $H \leq_{t.m.} G$ , if there is an  $H$ -topological minor model in  $G$ ;
- $H$  is an *immersion* of  $G$ , what we note  $H \leq_{imm.} G$ , if there is an  $H$ -immersion model in  $G$ ;
- $H$  is a *strong immersion* of  $G$ , what we note  $H \leq_{s.im.} G$ , if there is an  $H$ -strong immersion model in  $G$ ;

We add  $\mu$  as superscript of the aforementioned order symbols when we want to specify that  $\mu$  is a model witnessing the relation (like in  $H \leq_{c.}^{\mu} G$ ). Observe that each of the aforementioned relation defines a qoset on the class of graphs. For every  $\preceq \in \{\leq_{sg.}, \leq_{i.sg.}, \leq_{c.}, \leq_{m.}, \leq_{i.m.}\}$ , the graph  $G$  is said to be  $H$ - $\preceq$ -free, or to *exclude*  $H$  with respect to  $\preceq$ , if  $H \preceq G$  does not hold.

We now give the equivalent definition of these containment relations in terms of local operations. The list below also contains the definitions of dissolution, induced immersions, and induced topological minors, that we did not defined in terms of models as we will not consider them (see e.g. [KO04b] for a definition).

- $H \leq_{i.sg.} G$  iff there is a sequence of vertex deletions transforming  $G$  into  $H$ ;
- $H \leq_{sg.} G$  iff there is a sequence of vertex deletions and edge deletions transforming  $G$  into  $H$ ;
- $H \leq_{m.c.} G$  (or  $H \leq_{c.} G$  if  $H$  and  $G$  are simple graphs) iff there is a sequence of edge contractions transforming  $G$  into  $H$ ;
- $H \leq_{i.m.} G$  iff there is a sequence of vertex deletions and edge contractions transforming  $G$  into  $H$ ;
- $H \leq_{m.} G$  iff there is a sequence of vertex deletions, edge deletions and edge contractions transforming  $G$  into  $H$ ;
- $H \leq_{imm.} G$  iff there is a sequence of vertex deletions, edge deletions and lifts transforming  $G$  into  $H$ ;
- $H \leq_{t.m.} G$  iff there is a sequence of vertex deletions, edge deletions, and vertex dissolutions transforming  $G$  into  $H$ ;
- $H$  is an induced topological minor of  $G$  iff there is a sequence of vertex deletions and vertex dissolutions transforming  $G$  into  $H$ ;
- $H$  is an induced immersion of  $G$  iff there is a sequence of vertex deletions and lifts transforming  $G$  into  $H$ .

Note that the aforementioned sequences of operations are allowed to be empty. **Table 2.2** summarizes these definitions. In this table,  $VD$  stands for “vertex deletion”,  $ED$  for “edge deletion”,  $C^-$  for vertex dissolution,  $C$  for edge contraction and  $L$  for lift. The relations between the different containment relations are depicted on **Figure 2.9** where the same abbreviations are used.

The aforementioned orderings allow us to define classes of graphs.

Relation	$VD$	$ED$	$C^-$	$C$	$L$
isomorphism					
spanning subgraph		✓			
induced subgraph	✓				
subgraph	✓	✓			
dissolution			✓		
contraction			✓	✓	
induced minor	✓		✓	✓	
induced topological minor	✓		✓		
topological minor	✓	✓	✓		
minor	✓	✓	✓	✓	
induced immersion	✓				✓
immersion	✓	✓			✓

Table 2.2: Containment relations defined in terms of local operations.

**Classes defined by graph orderings.** Let  $H$  be a graph. An *major of  $H$*  ( $H$ -major for short) is a subgraph-minimal graph that contains  $H$  as a minor. We denote the class of  $H$ -majors by  $\mathcal{M}(H)$ .

A *subdivision of  $H$*  ( $H$ -subdivision for short) is a graph obtained from  $H$  by subdividing edges. Observe that a graph  $G$  has a subgraph isomorphic to subdivision of  $H$  iff it contains  $H$  as a topological minor. We denote by  $\mathcal{T}(H)$  the class of all subdivisions of  $H$ .

Also, an *immersion expansion of  $H$*  ( $H$ -immersion expansion for short) is a subgraph-minimal graph that contain  $H$  as an immersion. Again, a graph  $G$  has a subgraph isomorphic to an immersion expansion of  $H$  iff it contains  $H$  as an immersion. We write  $\mathcal{I}(H)$  for the class of immersions expansions of  $H$ . These definitions are extended to classes:  $\mathcal{M}(\mathcal{H})$  is the class of all subgraph-minimal graphs that contain some member of a class  $\mathcal{H}$  as a minor (and similarly for  $\mathcal{T}(\mathcal{H})$  and  $\mathcal{I}(\mathcal{H})$ ).

## 2.5 Tree-like decompositions and width parameters

A considerable amount of the recent algorithmic advances relies on tree-like decompositions, that are a way to decompose the graph into subsets organized in a tree-like fashion. These decompositions give rise to graph parameters usually called *width parameters*. Among the existing decompositions, tree-decompositions are certainly those that received the most considerable attention.

**Definition 2.2.** A *tree decomposition* of a graph  $G$  is a pair  $(T, \mathcal{X})$  where  $T$  is a tree and  $\mathcal{X}$  a family  $\{X_t\}_{t \in V(T)}$  of subsets of  $V(G)$  (called *bags*) indexed by elements of  $V(T)$  and such that the following holds

- (i)  $\bigcup_{t \in V(T)} X_t = V(G)$ ;
- (ii) for every edge  $e$  of  $G$  there is an element of  $\mathcal{X}$  containing both ends of  $e$ ;




Figure 2.9: Connections between common orderings of graphs. An arrow from an order relation  $\preceq$  to another  $\preceq'$  means that  $H \preceq' G$  holds whenever  $H \preceq G$  does. Colored areas represent allowed local operations for each relation: vertex deletion (red), edge deletion (blue), edge contraction or vertex dissolution (green), and lift (orange). The allowed operations are also specified by letters with the same convention as in Table 2.2.

- (iii) for every  $v \in V(G)$ , the subgraph of  $T$  induced by  $\{t \in V(T), v \in X_t\}$  is connected.

The *width* of a tree decomposition  $T$  is defined as equal to  $\max_{t \in V(T)} |X_t| - 1$ . The *treewidth* of  $G$ , written  $\text{tw}(G)$ , is the minimum width of any of its tree decompositions.

The *torso* of a bag  $X_t$  of a tree decomposition  $(T, \{X_t\}_{t \in V(T)})$  is the underlying simple graph of the graph obtained from  $G[X_t]$  by adding all the edges  $\{x, y\}$  such that  $x, y \in X_t \cap X_{t'}$  for some neighbor  $t'$  of  $t$  in  $T$ .

Treewidth has been extensively used in Algorithmics and Combinatorics. The Graph Minor series of Robertson and Seymour provides several results and tools related to this parameter. In particular, Robertson and Seymour proved that every graph of *big enough* treewidth contains as a minor a *big* grid. This result will be discussed in Chapter 4. Also, it appears that several algorithmic problems, that are hard in general, become tractable on graphs of bounded treewidth. In this direction, Courcelle proved [Cou90] that a large family of problems (those that can be expressed

in monadic second-order logic<sup>2</sup>) can be solved in linear time on graphs of bounded treewidth.

In order to deal with tree decompositions, we sometimes consider nice tree decompositions, which are defined as follows.

**Definition 2.3** ([Klo94a]). A triple  $(T, r, \{X_t\}_{t \in V(T)})$  is said to be a *nice* tree decomposition of a graph  $G$  if  $(T, \{X_t\}_{t \in V(T)})$  is a tree-decomposition where the following holds:

1. every vertex of  $T$  has degree at most 3;
2.  $(T, r)$  is a rooted tree and the bag of the root  $r$  is empty ( $X_r = \emptyset$ );
3. every vertex  $t$  of  $T$  is
  - either a *base node*, i.e. a leaf of  $T$  whose bag is empty ( $X_t = \emptyset$ ) and different from the root;
  - or an *introduce node*, i.e. a vertex with only one child  $t'$  such that  $X_{t'} = X_t \cup \{u\}$  for some  $u \in V(G) \setminus X_t$ ;
  - or a *forget node*, i.e. a vertex with only one child  $t'$  such that  $X_t = X_{t'} \cup \{u\}$  for some  $u \in V(G) \setminus X_t$ ;
  - or a *join node*, i.e. a vertex with two child  $t_1$  and  $t_2$  such that  $X_t = X_{t_1} = X_{t_2}$ .

It is known that every graph  $G$  has an nice tree decomposition with width  $\text{tw}(G)$  and at most  $4n$  nodes [Klo94a]. A more restrictive notion is the one of path decomposition, defined as follows.

**Definition 2.4** ([Klo94b]). A tree decomposition  $(T, \{X_t\}_{t \in V(T)})$  is a *path decomposition* if  $T$  is a path, and the *pathwidth* of  $G$ , that we write  $\text{pw}(G)$ , is the minimum width of a path decomposition of  $G$ .

A path decomposition  $(p_1 p_2 \dots p_k, \{X_{p_i}\}_{i \in \llbracket 1, k \rrbracket})$  of a graph  $G$  is said to be *nice* if  $|X_{p_1}| = 1$  and

$$\forall i \in \llbracket 2, k \rrbracket, |(X_{p_i} \setminus X_{p_{i-1}}) \cup (X_{p_{i-1}} \setminus X_{p_i})| = 1$$

It is known [BT04] that every graph have an optimal path decomposition which is nice and that in such decomposition, every node  $p_i$  is either an *introduce node* (i.e. either  $i = 1$  or  $|X_{p_i} \setminus X_{p_{i-1}}| = 1$ ) or a *forget node* (i.e.  $|X_{p_{i-1}} \setminus X_{p_i}| = 1$ ).

*Remark 2.5.* For every graph  $G$  on  $n$  vertices, there is an optimal path decomposition with  $n$  introduce nodes and  $n$  forget nodes (one of each for each vertex of  $G$ ), thus of length  $2n$ .

Tree partitions have proven useful when dealing with problems related to vertices. For instance, deciding if the deletion of a certain number of vertices yields a given property is such a problem. However, these decompositions are not well-suited for the study of problems related to edges. Therefore, several attempts have been done to design an *edge-analogues* of treewidth. Let us present two of them.

---

<sup>2</sup>We do not define this logic here as we will not use it.

**Definition 2.6.** A *tree partition* of a graph  $G$  is a pair  $\mathcal{D} = (T, \mathcal{X})$  where  $T$  is a tree and  $\mathcal{X} = \{X_t\}_{t \in V(T)}$  is a partition of  $V(G)$  (the elements of which are called *bags*) such that either  $n(T) = 1$  or for every  $\{x, y\} \in E(G)$ , there exists an edge  $\{t, t'\} \in E(T)$  where  $\{x, y\} \subseteq X_t \cup X_{t'}$ . In other words, the endpoints of every edge of  $G$  either belong to the same bag or they belong to bags of adjacent vertices of  $T$ . Given an edge  $f = \{t, t'\} \in E(T)$ , we define  $E_f$  as the set of edges with one endpoint in  $X_t$  and the other in  $X_{t'}$ . The *width* of  $\mathcal{D}$  is defined as  $\max\{\max_{t \in V(T)} |X_t|, \max_{f \in E(T)} |E_f|\}$ . The *tree partition width* of  $G$  is the minimum width over all tree partitions of  $G$  and is denoted by  $\text{tpw}(G)$ .

An example of a tree-partition of a graph is given in [Figure 2.10](#).




Figure 2.10: A depiction of a tree-partition of a graph, where the tree-structured bags of the tree-partition are represented in gray and the graph in black. Notice that edges of the graph are not allowed to connect vertices of bags that are not adjacent.

A *rooted tree partition* of a graph  $G$  is a triple  $\mathcal{D} = (T, s, \mathcal{X})$  where  $(T, s)$  is a rooted tree and  $(T, \mathcal{X})$  is a tree partition of  $G$ .

Tree partitions have been introduced in [\[See85\]](#) (see also [\[Hal91\]](#)) and tree partition width has been defined for simple graphs in [\[DO96\]](#). The above definition is an extension of the original definition to the setting of (multi)graphs. Tree partitions will in particular be used in [Section 4.1](#) and [Subsection 5.3.1](#). However, a drawback of tree partition width compared to treewidth is that a graph with large tree partition width does not necessarily contain a large substructure, whereas a graph of large treewidth is known to contain a large grid, as mentioned above (see [Theorem 4.1](#) for a formal statement). For instance, the graph obtained by setting to  $k$  the multiplicity of every edge of a path on  $k$  edges (for some positive integer  $k$ ) has tree partition width  $k$ , however its subgraphs are quite *poor* compared to those of a graph containing a large grid as a minor. Ding described in [\[DO96, Theorem 1.2\]](#) the subgraphs that one can expect in graphs of large tree partition width, as we mention in [Subsection 5.4.3](#).

A decomposition that avoids this pitfall has been recently introduced by Wollan in [\[Wol15\]](#) under the name *tree-cut decomposition*. Let us give a formal definition of such a decomposition. A *near-partition* of a set  $S$  is a collection of pairwise disjoint subsets  $S_1, \dots, S_k \subseteq S$  (for some  $k \in \mathbb{N}$ ) such that  $\bigcup_{i=1}^k S_i = S$ . Observe that this definition allows a set of the family to be empty.

**Definition 2.7** ([\[Wol15\]](#)). A *tree-cut decomposition* of a graph  $G$  is a pair  $\mathcal{D} = (T, \mathcal{X})$  where  $T$  is a tree and  $\mathcal{X} = \{X_t\}_{t \in V(T)}$  is a near-partition of  $V(G)$ .

In particular, a tree-partition is a special type of tree-cut decomposition. However, tree-cut decompositions are much more general as edges are allowed between vertices of non-adjacent bags. An example of a tree-cut decomposition is given in [Figure 2.11](#).




Figure 2.11: A depiction of a tree-cut decomposition a graph, where the tree-structured bags of the tree-cut decomposition are represented in gray and the graph in black. Notice that bags may be empty and edges are allowed to connect vertices of bags that are not adjacent. Edges of the graph are drawn along the edges of the tree to stress that they belong to the adhesions of these edges.

Let  $\mathcal{D} = (T, \{X_t\}_{t \in V(T)})$  be a tree-cut decomposition of a graph  $G$ . If, for an edge  $\{u, v\}$  of  $T$ , we define  $T_u$  (respectively  $T_v$ ) as the connected component of  $T \setminus \{\{u, v\}\}$  that contains  $u$  (respectively  $v$ ), then the *adhesion* of the edge  $\{u, v\}$  in  $\mathcal{D}$ , that we write  $\text{adh}_{\mathcal{D}}(\{u, v\})$  (or  $\text{adh}(\{u, v\})$  when the decomposition is given by the context), is the set of edges of  $G$  that have the one endpoint in  $\bigcup_{t \in V(T_u)} X_t$  and the other one in  $\bigcup_{t \in V(T_v)} X_t$ . Intuitively, the adhesion of  $e$  is the set of edges of the graph that “fly over”  $e$  in the tree-cut decomposition. On [Figure 2.11](#), the adhesion of an edge of the tree (depicted by a thick grey line) is the set of edges of the graph (depicted by black lines) that are drawn inside this edge. The *width* of the decomposition  $(T, \mathcal{X})$ , is

$$\max\left\{\max_{e \in E(T)} |\text{adh}(e)|, \max_{t \in V(T)} |V(H_t)|\right\},$$

where  $H_t$  is the graph obtained from  $G$  by first consolidating the vertices of the bags of each subtree of  $T \setminus t$  into one vertex and then removing the possibly created loops and vertices of degree two. The *tree-cut width* of  $G$ , denoted by  $\text{tcw}(G)$ , is the minimum width of a tree-cut decomposition of  $G$ .

The motivation for the introduction of tree-cut decompositions is that, as proved by Wollan in [[Wol15](#)], every graph of *large* tree-cut width contains a *large* wall as an immersion (cf. [Theorem 4.67](#)), which was not true for tree-partitions, as mentioned above. Also, an immersion of a graph of tree-cut width  $k$  (for some  $k \in \mathbb{N}$ ) has tree-cut width at most  $k$ . In that sense, tree-cut decompositions seems to play the same role for immersions as tree decompositions do for minors.

Let us define a concept similar to nice tree decompositions for tree-cut decompositions. A *rooted tree-cut decomposition* of a graph  $G$  is a triple  $\mathcal{D} = (T, s, \mathcal{X})$  where

$(T, s)$  is a rooted tree and  $(T, \mathcal{X})$  is a tree-cut decomposition of  $G$ . Let  $\mathcal{D} = (T, s, \mathcal{X})$  be a rooted tree-cut decomposition of  $G$ . We set

$$G_t = G \left[ \bigcup_{u \in \text{desc}_{(T,s)}(t)} X_u \right].$$

In a rooted tree-cut decomposition, the adhesion of a vertex  $t \in V(T)$  (noted  $\text{adh}(t)$ ) is the adhesion of the first edge of a path from  $t$  to the root, or the empty set if  $t = s$ . A vertex  $t \in V(T)$  is *thin* if  $\text{adh}_{\mathcal{D}}(t) \leq 2$ , *bold* otherwise. We also say that  $\mathcal{D}$  is *nice* if for every thin vertex  $t \in V(T)$  we have

$$N(V(G_t)) \cap \bigcup_{b \text{ is a sibling of } t} V(G_b) = \emptyset.$$

In other words, there is no edge from a vertex of  $G_t$  to a vertex of  $G_b$ , for any sibling  $b$  of  $t$ , whenever  $t$  is thin. This notion can be seen as an analogue of nice tree-decompositions.

In [GKS15], Ganian et al. proved that every graph has a tree-cut decomposition of minimal width, that, moreover, is nice.

**Proposition 2.8** ([GKS15]). *Every rooted tree-cut decomposition can be transformed into a nice tree-cut decomposition without increasing the width.*

# CHAPTER 3

---

## GRAPHS AND WELL-QUASI-ORDERING

In this chapter, we consider quasi-orders defined by the usual order relations on graphs. After presenting some results and techniques on well-quasi-ordering, we focus on the contraction and induced minor relations. For these relations, we characterize the well-quasi-ordered subclasses, among those that are defined by excluding a unique graph. This chapter contains material that previously appeared in the following articles:

- [BKRT15a, BKRT15b] *Induced minors and well-quasi-ordering*, co-authored with Jarosław Błasiok, Marcin Kamiński, and Théophile Trunck, EuroComb 2015;
- [KRT14] *Multigraphs without large bonds are well-quasi-ordered by contraction*, co-authored with Marcin Kamiński and Théophile Trunck, submitted (2014); and
- [KRT16] *Well-quasi-ordering  $H$ -contraction-free graphs*, co-authored with Marcin Kamiński and Théophile Trunck, submitted (2015).

### 3.1 Preliminaries on well-quasi-orders

A *well-quasi-order* is a quasi-order which contains neither an infinite decreasing sequence nor an infinite collection of pairwise incomparable elements. This strengthening of the concept of well-order has been introduced in the 50's. Since then, a whole theory of well-quasi-orders has been developed and has led to surprising results and unsuspected developments.

Before going more into detail, let us present one of the most remarkable aspects of well-quasi-orders. Several objects of interest in graph theory are classes of graphs that are *downward closed*. That is, every graph that is smaller (wrt. a given order) than a graph in the class also belongs to the class. For instance, forests, planar graphs, more generally, graphs of genus at most  $g$  (for every fixed  $g \in \mathbb{N}$ ), and graphs of treewidth at most  $k$  (for every fixed  $k \in \mathbb{N}$ ) are downward closed wrt. the minor relation. On

the opposite, the class of 3-colorable graphs is not downward closed wrt. the minor relation, as witnessed by the graph of [Figure 3.1](#), which is 3-colorable but contains the non-3-colorable graph  $K_4$  as a minor.




Figure 3.1: A 3-colorable graph that contains  $K_4$  as minor.

For every class  $C$  of a quasi-order  $(S, \preceq)$ , we can ask the question : “can we *easily* characterize the elements of  $S$  that belong to  $C$ ?”. An approach on this question when  $C$  is downward closed is to consider the minimal elements of the complementary of  $C$ , when they exist. These elements are called *obstructions*, because every element  $x$  of  $S$  belongs to  $C$  if and only if there is no obstruction  $y$  such that  $y \preceq x$ . That way, obstructions provide a precise characterization of  $C$ . Here the well-quasi-orders come into play: if  $(S, \preceq)$  is a well-quasi-order, then  $C$  has finitely many obstructions. These obstructions exist because  $S \setminus C$  does not contain infinite decreasing sequences and they are finitely many as the set of obstructions is an antichain. In other words, there are elements  $x_0, \dots, x_c$  (for some  $c \in \mathbb{N}$  depending on  $C$ ) such that the following holds:

$$\forall x \in S, x \in C \iff \forall i \in \llbracket 0, c \rrbracket, x_i \not\preceq x.$$

This property, that Pouzet refers to as a *finite basis property* in [\[Pou85\]](#), has the following algorithmic implication. If, for every  $i \in \llbracket 0, c \rrbracket$ , there is a algorithm that can decide if  $x_i \preceq y$  in a time that is polynomial in the size of  $y$  for every  $y \in S$ , then the membership of  $C$  can be decided in a time that is polynomial in the input size. As an illustration, let us consider the minor relation. Robertson and Seymour proved in their Graph Minors series [\[RS04\]](#) that this relation well-quasi-orders all (finite) graphs and they also gave an algorithm that, for every fixed graph  $H$ , decides if  $H$  is a minor of the input graph  $G$  in a time that is polynomial in the size of  $G$  [\[RS95\]](#). As mentioned above, a consequence of these results is that, for every graph class that is downwards closed wrt. minors, there is an algorithm that answers in polynomial time whether the input graph belongs to the considered class. It is worth noting that these results are purely existential: they do not provide any way to construct the algorithm. Even the upper-bound on the complexity of these algorithms is existential, as it depends on the number of obstructions for the considered class.

Recall that a qoset (for quasi-ordered set) is a pair of a set and a relation that is reflexive and transitive. Also, for every qoset  $(A, \preceq)$ , we denote by  $(A^*, \preceq^*)$  the qoset of finite sequences over  $A$  ordered by the subsequence relation (cf. [Section 2.2](#)). A cornerstone of the theory of well-quasi-orders is the following theorem, usually called “Higman’s Lemma”.

**Theorem 3.1** (Higman’s Lemma, [\[Hig52\]](#)). *If  $(A, \preceq)$  is a wqo, then so is  $(A^*, \preceq^*)$ .*

It is noteworthy that Higman’s Lemma appears in the proofs of several later results on well-quasi-ordering. In fact the main result of [\[Hig52\]](#) is more general than

**Theorem 3.1** but we will only use this form in this chapter. Given a qoset  $(S, \preceq)$ , the main question we are here concerned with is the following.

*Question 3.2.* Is  $(S, \preceq)$  a well-quasi-order?

Let us now present some known results answering this question for various choices of  $S$  and  $\preceq$  among graph classes and graph orderings. Unless otherwise specified, the word *graph* will in this chapter refer to simple graphs, i.e. graphs without loops or multiple edges. One of the most important well-quasi-ordering result on graphs is certainly the aforementioned Graph Minor Theorem by Robertson and Seymour.

**Theorem 3.3** (Graph Minor Theorem, [RS04]). *The class of all graphs is well-quasi-ordered by the minor relation.*

The Graph Minor Theorem extends the earlier Kruskal Tree Theorem, which was concerned with trees.

**Theorem 3.4** (Kruskal Tree Theorem, [Kru60], see also [NW63]). *The class of all trees is well-quasi-ordered by the topological minor relation.*

Robertson and Seymour later proved that the same also holds for the immersion relation.

**Theorem 3.5** ([RS10]). *The class of all graphs is well-quasi-ordered by the immersion relation.*

However, among the usual containment relations on graphs (listed in Section 2.4), the minor relation and the immersion relation are the only relations known to be well-quasi-order in general (i.e. on the class of all graphs). Regarding other relations, either the problem is still open, or infinite antichains are known. Table 3.1 summarizes the status of Question 3.2 when  $S$  is the class of all graphs.

well-quasi-order	open	not a well-quasi-order
minors	strong immersions	subgraphs
immersions	induced immersions	induced subgraphs
		topological minors
		induced topological minors
		contractions

Table 3.1: Status of Question 3.2 for the common orderings of graphs.

For all the qosets that are not well-quasi-orders, a natural question is to identify the well-quasi-ordered subclasses.

*Question 3.6.* For which  $C \subseteq S$  is  $(C, \preceq)$  a well-quasi-order?

Much attention has been brought to this question in the last decades. For instance, Fellows et al. proved in [FHR12] that graphs with bounded feedback-vertex-set are well-quasi-ordered by topological minors. Another result is that of Oum [Oum08]

who proved that graphs of bounded rank-width are wqo by vertex-minors<sup>1</sup>. Other papers considering this question include [Tho85, Dam90, Din92, Din98, Pet02, Din09, DRT10, HL14, AL14]. In order to approach Question 3.6, which is quite general, one may consider a fixed family of subclasses only. One way to do this is to look at graph classes defined by forbidden substructures.

*Question 3.7.* For which  $x \in S$  is the set  $\{y \in S, x \not\preceq y\}$  well-quasi-ordered by  $\preceq$ ?

This line of research has been fruitful. Let us present some results in this direction, starting from the following theorem by Damaschke, that initiated the quest for answers to Question 3.7.

**Theorem 3.8** (Dichotomy for induced subgraphs, [Dam90, Theorem 4]). *Let  $H$  be a graph. The class of  $H$ -induced subgraph-free graphs is well-quasi-ordered by the induced subgraph relation iff  $H \leq_{i.sg.} P_4$ .*

Notice that that the above result is a full characterization of subclasses that are well-quasi-ordered by the induced subgraph relation, among those that are uniquely defined. Such a dichotomy has also been proved for other orderings. Damaschke has been followed by Ding who proved two years later the following non-induced counterpart of Theorem 3.8.

**Theorem 3.9** (Dichotomy for subgraphs, [Din92]). *Let  $H$  be a graph. The class of  $H$ -subgraph-free graphs is well-quasi-ordered by the subgraph relation iff  $H \leq_{sg.} P_n$  for some  $n \in \mathbb{N}$ .*

More recently, Liu obtained a result of the same type for topological minors.

**Theorem 3.10** (Dichotomy for topological minors, [HL14]). *Let  $H$  be a graph. The class of  $H$ -topological minor-free graphs is well-quasi-ordered by topological minors iff  $H \leq_{t.m.} R_n$  for some  $n \in \mathbb{N}$ , where  $R_n$  is the multigraph obtained by doubling every edge of a path on  $n$  edges.*

About the induced minor relation, a well-quasi-ordered class defined by excluding an induced minor has been identified by Thomas.

**Theorem 3.11** ([Tho85]). *The class of  $K_4$ -(induced) minor-free graphs is well-quasi-ordered by the induced minor relation.*

Question 3.7 has also been considered in [KL11a, KL11b, Che11]. In this part, we provide an answer to this question for the following graph containment relations:

- the induced minor relation  $\leq_{i.m.}$  on simple graphs, in Section 3.6;
- the contraction relation  $\leq_c$  on simple graphs, in Section 3.7;
- the contraction relation  $\leq_{m.c.}$  on multigraphs<sup>2</sup>, in Section 3.8.

<sup>1</sup>Vertex-minors is another ordering of graphs, which we will not consider in this thesis.

<sup>2</sup>Recall that we use the symbol  $\leq_{m.c.}$  to stress that we are dealing with the contraction relation on multigraphs, i.e. that multiple edges that may appear after a contraction are not deleted.

As mentioned above, none of these relations is a well-quasi order in general. Infinite antichains witnessing this fact will be given in [Subsection 3.5.1](#). Let us introduce three graphs that play a major role in this chapter. They are depicted on [Figure 3.2](#). The first one,  $\hat{K}_4$ , is obtained by adding a vertex of degree two to  $K_4$ . The second one, called the *gem*, is constructed by adding a dominating vertex to  $P_4$ . The last one is the *diamond* (or  $D_2$ ), that one can obtain by removing an edge in  $K_4$ . In the forthcoming




Figure 3.2: The graphs  $\hat{K}_4$ , *gem*, and  $D_2$  (from left to right).

section we prove the following theorems.

**Theorem 3.12** (Dichotomy for induced minors). *Let  $H$  be a graph. The class of  $H$ -induced minor-free simple graphs is wqo by  $\leq_{i.m.}$  iff  $H \leq_{i.m.} \hat{K}_4$  or  $H \leq_{i.m.} \text{Gem}$ .*

**Theorem 3.13** (Dichotomy for contractions). *Let  $H$  be a graph. The class of  $H$ -contraction-free simple graphs is wqo by  $\leq_c$  iff  $H \leq_c D_2$ .*

**Theorem 3.14** (Dichotomy for multigraph contractions). *Let  $\mathcal{H}$  be a class of graphs. The class of  $\mathcal{H}$ -contraction-free multigraphs is wqo by  $\leq_{m.c.}$  iff there is an integer  $n$  such that  $\theta_{n'} \in \mathcal{H}$  and  $\overline{K_{n'}} \in \mathcal{H}$  for every integer  $n' \geq n$ .*

[Theorem 3.12](#) and [Theorem 3.13](#) are the induced minor and contraction counterparts of [Theorem 3.8](#). Besides, [Theorem 3.12](#) completes [Theorem 3.11](#) into a full dichotomy. We shall stress that [Theorem 3.14](#) is more general, in the sense that it characterizes *all* the well-quasi-ordered classes defined by forbidding contractions, whereas [Theorem 3.12](#) and [Theorem 3.13](#) only deal with uniquely defined such classes. It could be interesting to extend [Theorem 3.14](#) to induced minors and contractions in simple graphs, that is, to answer the following question.

*Question 3.15.* For which classes  $\mathcal{H}$  is the class of  $\mathcal{H}$ -induced minor-free (respectively  $\mathcal{H}$ -contraction-free) graphs well-quasi-ordered by  $\leq_{i.m.}$  (respectively  $\leq_c$ )?

A first step towards this goal could be to look at classes defined by excluding two graphs.

*Question 3.16.* For which graphs  $H_1, H_2$  is the class of  $\{H_1, H_2\}$ -induced minor-free (respectively  $\{H_1, H_2\}$ -contraction-free) graphs well-quasi-ordered by  $\leq_{i.m.}$  (respectively  $\leq_c$ )?

This question already received partial answers on the induced subgraph ordering, see e.g. [KL11a].

Another line of research when considering qosets that are not wqos is to consider infinite antichains. In his study of infinite antichains for the (induced) subgraph relation, Ding [Din09] introduced the concept of *canonical antichain*. Let us announce the definition we will give in Section 3.2. Let us consider a quasi-order  $(S, \preceq)$ . An antichain  $\mathcal{A}$  of this quasi-order is said to be *canonical* if, for every finite subset  $\mathcal{A}'$  of  $\mathcal{A}$ , the class  $\text{Excl}_{\preceq}(\mathcal{A} \setminus \mathcal{A}')$  (elements that are not larger than any element of  $\mathcal{A} \setminus \mathcal{A}'$ ) has no infinite antichain. Informally, a canonical antichain  $\mathcal{A}$  represents all infinite antichains of a quasi-order. Indeed, as excluding all but finitely many elements of  $\mathcal{A}$  also excludes all infinite antichains (according to the definition), an infinite antichain has only finitely many elements that are not larger than some element of  $\mathcal{A}$ . In that sense, a canonical antichain can be seen as a common denominator of all infinite antichains. Also, the presence of a canonical antichains discriminates between the quasi-orders where the infinite antichains are “similar enough” so that they can be represented by one of them, and the quasi-orders where infinite antichains are “too diverse” for this to happen.

Another motivation for introduction by Ding [Din09] of the concept of canonical antichains is the following: from the definition given above, a quasi-order has a finite canonical antichain iff it has no infinite antichain. This allows us to partition quasi-orders in three classes: those that have no infinite antichains, those that have infinite antichains and a canonical antichain, and those that have infinite antichains and no canonical antichain. The study of canonical antichains can then be seen as an extension of well-quasi-ordering theory, which mostly focuses on the presence of infinite antichains. The most natural question about canonical antichains is the following.

*Question 3.17.* If  $(S, \preceq)$  is not a wqo, does it have a canonical antichain?

As shown by the results below, the question of the presence or absence of a canonical antichain has been studied for several containment relations and graph classes.

**Theorem 3.18** ([Din09]). *Under the subgraph relation, the class of finite graphs has a canonical antichain.*

The canonical antichain for Theorem 3.18 given in [Din09] can be written  $\{B_i^j, i \in \mathbb{N}, j \in \llbracket 1, 4 \rrbracket\}$ , where the graphs  $B_i^j$  are depicted in Figure 3.3. Observe that from the definition of an antichain, we can immediately deduce that every infinite antichain for the subgraph relation contains infinitely many graphs of the same “type” among those presented on Theorem 3.18 (formally, infinitely many graphs of  $\{B_i^j\}_{i \in \mathbb{N}}$ , for some  $j \in \llbracket 1, 4 \rrbracket$ ).

**Theorem 3.19** ([Din09]). *Under the induced subgraph relation, the class of finite graphs does not have a canonical antichain.*

**Theorem 3.20** ([Din09]). *Under the induced subgraph relation, both the class of interval graphs and the class of bipartite permutation graphs have a canonical antichain.*

We give an answer to Question 3.17 for the contraction relation with the following result.




Figure 3.3: The elements of a canonical antichain for the subgraph relation.

**Theorem 3.21.** *Under the contraction relation, the class of all graphs does not have a canonical antichain.*

The proof of [Theorem 3.21](#) relies on the tools introduced in [\[Din09\]](#) that can be used to prove that a quasi-order does not have a canonical antichain. We also obtained a complete characterization of the canonical antichains of the multigraph contraction ordering. Let  $\mathcal{A}_\theta$  be the class of all connected multigraphs on two vertices, and  $\mathcal{A}_{\overline{K}}$  be the class of all edgeless graphs (see [Figure 3.4](#)).




Figure 3.4: Two infinite antichains for multigraph contractions: multiedges and edgeless graphs.

**Theorem 3.22.** *Every antichain  $\mathcal{A}$  of  $\leq_{\text{m.c.}}$  is canonical iff each of the following sets are finite:*

$$\mathcal{A}_\theta \setminus \mathcal{A}; \quad \mathcal{A}_{\overline{K}} \setminus \mathcal{A}; \quad \text{and} \quad \mathcal{A} \setminus \{\mathcal{A}_\theta \cup \mathcal{A}_{\overline{K}}\}.$$

In other words, an antichain  $\mathcal{A}$  is canonical iff it contains all but finitely many graphs from  $\mathcal{A}_\theta$ , all but finitely many graphs from  $\mathcal{A}_{\overline{K}}$ , and a finite number of graphs that do not belong to  $\mathcal{A}_\theta \cup \mathcal{A}_{\overline{K}}$ . Two easy consequences are that  $\leq_{\text{m.c.}}$  has infinite canonical antichains and the following result.

**Corollary 3.23.** *Every canonical antichain of  $\leq_{\text{m.c.}}$  is fundamental<sup>3</sup>.*

<sup>3</sup>The definition of the concept of *fundamental antichain* is given in [Section 3.2](#).

## 3.2 Notions specific to this chapter

Let us introduce notions that will be used in this chapter.

**Union and product.** If  $(A, \preceq_A)$  and  $(B, \preceq_B)$  are two qosets, then

- their union  $(A \cup B, \preceq_A \cup \preceq_B)$  is the qoset defined as follows: for every  $x, y \in A \cup B$ , we have  $x \preceq_{A \cup B} y$  if

$$(x, y \in A \text{ and } x \preceq_A y), \text{ or } (x, y \in B \text{ and } x \preceq_B y);$$

- their Cartesian product  $(A \times B, \preceq_A \times \preceq_B)$  is the qoset defined by:

$$\forall (a, b), (a', b') \in A \times B, (a, b) \preceq_{A \times B} (a', b') \text{ if } a \preceq_A a' \text{ and } b \preceq_B b'.$$

**Ordering sequences.** For any qoset  $(A, \preceq)$ , we define the relation  $\preceq^*$  on  $A^*$  as follows: for every  $r = \langle r_1, \dots, r_p \rangle$  and  $s = \langle s_1, \dots, s_q \rangle$  of  $A^*$ , we have  $r \preceq^* s$  if there is an increasing function  $\varphi: \llbracket 1, p \rrbracket \rightarrow \llbracket 1, q \rrbracket$  such that for every  $i \in \llbracket 1, p \rrbracket$  we have  $r_i \preceq_{s_{\varphi(i)}}$ . Observe that  $=^*$  is then the subsequence relation. This order relation is extended to the class  $\mathcal{P}^{<\omega}(A)$  of finite subsets of  $A$  as follows, generalizing the subset relation: for every  $B, C \in \mathcal{P}^{<\omega}(A)$ , we write  $B \preceq^P C$  if there is an injection  $\varphi: B \rightarrow C$  such that  $\forall x \in B, x \preceq \varphi(x)$ . Observe that  $=^P$  is the subset relation.

In order to stress that domain and codomain of a function are qosets, we sometimes use, in order to denote a function  $\varphi$  from a qoset  $(A, \preceq_A)$  to a qoset  $(B, \preceq_B)$ , the following notation:  $\varphi: (A, \preceq_A) \rightarrow (B, \preceq_B)$ .

**Monotonicity.** A function  $\varphi: (A, \preceq_A) \rightarrow (B, \preceq_B)$  is said to be *monotone* if it satisfies the following property:

$$\forall x, y \in A, x \preceq_A y \Rightarrow \varphi(x) \preceq_B \varphi(y).$$

Informally, a monotone function preserves the order. A function  $\varphi: (A, \preceq_A) \rightarrow (B, \preceq_B)$  is a *qoset epimorphism* (*epi* for short) if it is surjective and monotone.

**Closed sets.** Let  $(A, \preceq)$  be a qoset. A subset  $B \subseteq A$  is said to be *downward  $\preceq$ -closed* (sometimes shortened as  *$\preceq$ -closed*) if for every  $x, y \in A$  such that  $x \preceq y$  and  $y \in B$  we have  $x \in B$ . Symmetrically, we say that  $B$  is *upward  $\preceq$ -closed* if for every  $x, y \in A$  such that  $x \preceq y$  and  $x \in B$  we have  $y \in B$ .

**Incl and Excl.** Let  $(A, \preceq)$  be a qoset. For every  $x \in A$ , we define

$$\begin{aligned} \text{excl}_{\preceq}(x) &= \{y \in A, x \not\preceq y\} \quad \text{and} \\ \text{Incl}_{\preceq}(x) &= \{y \in A, y \preceq x\}. \end{aligned}$$

Informally,  $\text{Excl}_{\preceq}(x)$  is the class of elements *excluding*  $x$  (with respect to  $\preceq$ ), and  $\text{Incl}_{\preceq}(x)$  the class of elements *included* in  $x$ . We extend these definitions to any subset  $B \subseteq A$  as follows:

$$\begin{aligned} \text{Excl}_{\preceq}(B) &= \{y \in A, \forall x \in B, x \not\preceq y\} \quad \text{and} \\ \text{Incl}_{\preceq}(B) &= \{y \in A, \exists x \in B, y \preceq x\}. \end{aligned}$$

The set  $\text{Incl}_{\preceq}(B)$  contains all the elements of  $B$  and all the elements that are smaller than some element of  $B$ . It is sometimes referred to as the *closure* of  $B$  with respect to the relation  $\preceq$ . Observe both  $\text{Incl}_{\preceq}(B)$  and  $\text{Excl}_{\preceq}(B)$  are downward  $\preceq$ -closed sets. A subset  $B \subseteq A$  is said to be *uniquely defined* if  $B = \text{Excl}_{\preceq}(x)$  for some  $x \in A$ .

**Canonical antichains.** An antichain  $A$  of a qoset  $(S, \preceq)$  is said to be *canonical* if it is such that every  $\preceq$ -downwards-closed subclass  $J$  of  $S$  has an infinite antichain iff  $J \cap A$  is infinite. This definition differs from that given in the previous section, however they are equivalent [Din09, Proposition 1.2]. More intuition about canonical antichains is given in the previous section. If  $\text{Incl}(A)$  has no infinite antichains, then  $A$  is a *fundamental antichain*. This can be seen as a notion of minimality, as the elements that are below (with respect to the considered order) do not have infinite antichains.

**Labeled graphs.** As detailed in this chapter, labeling the vertices of graphs may simplify well-quasi-ordering proofs. Let us introduce some definitions related to graph labeling. For this, we consider a qoset  $(S, \preceq)$ . A  $(S, \preceq)$ -*labeled graph* is a pair  $(G, \lambda)$  such that  $G$  is a graph, and  $\lambda: V(G) \rightarrow \mathcal{P}^{<\omega}(S)$  is a function referred as the *labeling of the graph*. For the sake of simplicity, we will refer to the labeled graph of a pair  $(G, \lambda)$  by  $G$  and to  $\lambda$  by  $\lambda_G$ . If  $\mathcal{G}$  is a class of (unlabeled) graphs,  $\text{lab}_{(S, \preceq)}(\mathcal{G})$  denotes the class of  $(S, \preceq)$ -labeled graphs of  $\mathcal{G}$ . The notion of labeled graph generalizes the one of graph as any (unlabeled) graph can be seen as a  $\emptyset$ -labeled graph.

This definition of a labeled graph differs from the commonly used one where vertices are assigned single elements of  $S$  instead of finite subsets. Whereas the usual definition can be easily used to prove well-quasi-ordering results with the induced subgraph ordering, the definition we introduced is suited for proofs on graph orderings involving contractions. The extensions of these orderings to labelled graphs will be given in [Section 2.4](#).

**Rooted graphs.** In several proofs, the labels that will be assigned to vertices of graphs will be sets of rooted graphs. This will be used in order to *encode* a class of connected graphs as labeled 2-connected graphs (cf. [Lemma 3.31](#) and [Lemma 3.32](#)). Roots will also be used on trees to defined a partial order on the vertices.

A *rooted graph* is a couple  $(G, r)$  where  $G$  is a graph and  $r$  is a vertex of  $G$ , called *root* of  $G$ . For the sake of simplicity, we sometimes denote by  $G$  the rooted graph  $(G, r)$  and refer to its root by  $\text{root}(G)$ . If  $\mathcal{H}$  is a class of graphs, we define its *rooted closure*, denoted  $\mathcal{H}_r$  as the class of rooted graphs  $\mathcal{H}_r = \{(G, v) : G \in \mathcal{H}, v \in G\}$ .

### 3.3 Raising well-quasi-orders

As we explain in the forthcoming [Section 3.4](#), a way to show that a given qoset  $(A, \preceq_A)$  is a wqo is to build a wqo  $(B, \preceq_B)$  from smaller qosets that are known to be well-quasi-ordered and then find a correspondence between the elements of  $A$  and  $B$  that *preserves* the well-quasi-orderness. In this section, we present different constructions to obtain well-quasi-orders. In particular, we recall that being well-quasi-ordered is preserved by several operations including union, Cartesian product, and application of a monotone function.

Let us start with an easy remark.

*Remark 3.24.* If  $B \subseteq A$  and  $(A, \preceq)$  is a wqo, then so is  $(B, \preceq)$ .

Indeed, any infinite antichain of  $(B, \preceq)$  is an antichain of  $(A, \preceq)$ .

**Union and product.** Recall that, in the union of two qosets  $(A, \preceq_A)$  and  $(B, \preceq_B)$ , two elements are comparable only if both belong to the same qoset, in which case they are ordered as in this qoset. In the Cartesian product of  $(A, \preceq_A)$  and  $(B, \preceq_B)$ , the elements are pairs over  $A \times B$  and are compared coordinate-wise. The formal definitions can be found in [Section 2.2](#).

*Remark 3.25* (union of wqos). If  $(A, \preceq_A)$  and  $(B, \preceq_B)$ , are two wqos, then so is  $(A \cup B, \preceq_A \cup \preceq_B)$ .

In fact, for every infinite antichain  $S$  of  $(A \cup B, \preceq_A \cup \preceq_B)$ , there is an infinite subsequence of  $S$  whose all elements belong to one of  $A$  and  $B$  (otherwise  $S$  is finite). But then one of  $(A, \preceq_A)$  and  $(B, \preceq_B)$  has an infinite antichain, a contradiction to our initial assumption. Similarly, every finite union of wqos is a wqo.

**Lemma 3.26** (Higman [[Hig52](#)]). *If  $(A, \preceq_A)$  and  $(B, \preceq_B)$  are wqo, then so is  $(A \times B, \preceq_A \times \preceq_B)$ .*

**Sequences and subsets.** Higman's Lemma ([Theorem 3.1](#)), which we mentioned in the previous section, states that finite sequences over a well-quasi-order  $(A, \preceq)$  are well-quasi-ordered by  $\preceq^*$ . It has the following easy corollary.

**Corollary 3.27.** *If  $(A, \preceq)$  is a wqo, then so is  $(\mathcal{P}^{<\omega}(A), \preceq^{\mathcal{P}})$ .*

**Surjective images.** Recall that an epi is a function from one qoset to an other that is surjective and monotone. Epis have the following interesting property, which we will extensively use to show that some qosets are well-quasi-ordered.

*Remark 3.28* (epi from a wqo). If the domain of an epi is wqo, then its codomain is also wqo.

Indeed, for any pair  $x, y$  of elements of the domain of an epi  $\varphi$  such that  $\varphi(x)$  and  $\varphi(y)$  are incomparable,  $x$  and  $y$  are incomparable as well (by monotonicity of  $\varphi$ ). Therefore, and as  $\varphi$  is surjective, any infinite antichain of the codomain of  $\varphi$  can be translated into an infinite antichain of its domain.

In order to apply [Remark 3.28](#) one needs to show that the considered function is monotone. The aim of the next remark is to make this task easier when dealing with multivariate functions.

*Remark 3.29* (componentwise monotonicity). Let  $(A, \preceq_A)$ ,  $(B, \preceq_B)$ , and  $(C, \preceq_C)$  be three qosets and let  $f: (A \times B, \preceq_A \times \preceq_B) \rightarrow (C, \preceq_C)$  be a function. If we have both

$$\forall a \in A, \forall b, b' \in B, b \preceq_B b' \Rightarrow f(a, b) \preceq_C f(a, b') \quad (3.1)$$

$$\text{and } \forall a, a' \in A, \forall b \in B, a \preceq_A a' \Rightarrow f(a, b) \preceq_C f(a', b), \quad (3.2)$$

then  $f$  is monotone.

Indeed, let  $(a, b), (a', b') \in A \times B$  be such that  $(a, b) \preceq_A \times \preceq_B (a', b')$ . By definition of the relation  $\preceq_A \times \preceq_B$ , we have both  $a \preceq_A a'$  and  $b \preceq_B b'$ . From line (3.1) we get that  $f(a, b) \preceq_C f(a, b')$  and from line (3.2) that  $f(a, b') \preceq_C f(a', b')$ , hence  $f(a, b) \preceq_C f(a', b')$  by transitivity of  $\preceq_C$ . Thus  $f$  is monotone. Observe that this remark can be generalized to functions with more than two arguments.

**Excluding elements.** We will often deal with classes defined by forbidding an element, that is, subclasses of a qoset  $(S, \preceq)$  of the form  $\{y \in S, x \not\preceq y\}$  for some  $x \in S$ .

*Remark 3.30.* Let  $(S, \preceq)$  be a qoset. For every  $x, x' \in S$  such that  $x' \preceq x$ , we have  $\{y \in S, x' \not\preceq y\} \subseteq \{y \in S, x \not\preceq y\}$ . As a consequence,  $(\{y \in S, x' \not\preceq y\}, \preceq)$  is a wqo whenever  $(\{y \in S, x \not\preceq y\}, \preceq)$  is.

**Labeled graphs.** As defined in [Section 2.3](#), a graph labeled by a qoset is a graph, the vertices of which are assigned subsets of this qoset. Informally speaking, labels will be used to encode connected graphs into labeled 2-connected graphs. Given a connected graph which is not 2-connected, we can pick an arbitrary block  $B$ , delete the rest of the graph, and label each vertex  $v$  of  $B$  by the subgraph it was attached to in the original graph if  $v$  was a cutvertex, and by  $\emptyset$  otherwise. That way, contracting the label of a vertex  $v$  corresponds to contracting a subgraph. This intuition is formalized by the next result.

**Lemma 3.31** ([\[FHR12\]](#)). *Let  $\mathcal{G}$  be a class of graphs. If for any wqo  $(S, \preceq)$  the class of  $(S, \preceq)$ -labeled 2-connected graphs of  $\mathcal{G}$  is wqo by  $\leq_{i.m.}$ , then  $(\mathcal{G}, \leq_{i.m.})$  is a wqo.*

This result can also be proved for multigraph contractions.

**Lemma 3.32.** *Let  $\mathcal{G}$  be a class of connected multigraphs. If for any wqo  $(S, \preceq)$  the class of  $(S, \preceq)$ -labeled 2-connected graphs of  $\mathcal{G}$  is wqo by  $\leq_{m.c.}$ , then  $(\mathcal{G}, \leq_{m.c.})$  is a wqo.*

*Proof.* This proof is very similar to the proof of [Lemma 3.31](#) given in [\[FHR12\]](#). Assuming that  $(\mathcal{H}, \leq_{\text{m.c.}})$  is not a wqo, we will reach a contradiction by showing that its rooted closure  $(\mathcal{H}_r, \leq_{\text{m.c.}})$  is a wqo.

Let  $\langle G_i \rangle_{i \in \mathbb{N}}$  be a bad sequence in  $\mathcal{H}_r$  such that, for every  $i \in \mathbb{N}$ , there is no  $G \leq_{\text{m.c.}} G_i$  such that a bad sequence starts with  $G_0, \dots, G_{i-1}, G$  (a so-called *minimal bad sequence*). For every  $i \in \mathbb{N}$ , let  $A_i$  be the block of  $G_i$  which contains  $\text{root}(G_i)$ . Let  $C_i$  be the set of cutvertices of  $G_i$  that are included in  $A_i$ . For each cutvertex  $c \in C_i$ , let  $B_c^i$  be the connected component in  $G_i \setminus (V(A_i) \setminus C_i)$ , and rooted at  $c$  (i.e., we set  $\text{root}(B_c^i) = c$ ). Note that we have  $B_c^i \leq_{\text{m.c.}} G_i$ .

Let us denote by  $\mathcal{B}$  the family of rooted graphs  $\mathcal{B} = \{B_c^i : c \in C_i, i \in \mathbb{N}\}$ . We will show that  $(\mathcal{B}, \leq_{\text{m.c.}})$  is a wqo. Let  $\langle H_j \rangle_{j \in \mathbb{N}}$  be an infinite sequence in  $\mathcal{B}$  and for every  $j \in \mathbb{N}$  choose an  $i = \varphi(j) \in \mathbb{N}$  for which  $H_j \leq_{\text{m.c.}} G_i$ . Pick a  $j$  with smallest  $\varphi(j)$  and consider the sequence  $G_1, \dots, G_{\varphi(j)-1}, H_j, H_{j+1}, \dots$ . By minimality of  $\langle G_i \rangle_{i \in \mathbb{N}}$  and by our choice of  $j$ , since  $H_j \leq_{\text{m.c.}} G_{\varphi(j)}$  and  $H_j \neq G_{\varphi(j)}$ , this sequence is good so contains a good pair  $(G, G')$ . Now, if  $G$  is among the first  $\varphi(j) - 1$  elements, then as  $\langle G_i \rangle_{i \in \mathbb{N}}$  is bad we must have  $G' = H_{j'}$  for some  $j' \geq j$  and so we have  $G_{i'} = G \leq_{\text{m.c.}} G' = H_{j'} \leq_{\text{m.c.}} G_{\varphi(j')}$ , a contradiction. So there is a good pair in  $\langle H_i \rangle_{i \geq j}$  and hence the infinite sequence  $\langle H_j \rangle_{j \in \mathbb{N}}$  has a good pair, so  $(\mathcal{B}, \leq_{\text{m.c.}})$  is a wqo.

We will now find a good pair in  $\langle G_i \rangle_{i \in \mathbb{N}}$  to show a contradiction. The idea is to label the graph family  $\mathcal{A} = \{A_i\}_{i \in \mathbb{N}}$  so that each cutvertex  $c$  of a graph  $A_i$  gets labeled by their corresponding connected component  $B_c^i$ , and the roots are preserved under this labeling. More precisely, for each  $A_i$  we define a labeling  $\sigma_i$  that assigns to every vertex  $v \in V(G_i)$  a label  $\{(\sigma_i^1(v), \sigma_i^2(v))\}$  defined as follows:

- $\sigma_i^1(v) = 1$  if  $v = \text{root}(G_i)$  and  $\sigma_i^1(v) = 0$  otherwise;
- $\sigma_i^2(v) = B_v^i$  if  $v \in C_i$  and  $\sigma_i^2(v)$  is the one-vertex rooted graph otherwise.

The labeling  $\sigma$  of  $\mathcal{A}$  is then  $\{\sigma_i : i \in \mathbb{N}\}$ . Let us define a quasi-ordering  $\preceq$  on the set of labels  $\Sigma$  assigned by  $\sigma$ . For two labels  $(s_a^1, s_a^2), (s_b^1, s_b^2) \in \Sigma$  we define  $(s_a^1, s_a^2) \preceq (s_b^1, s_b^2)$  iff  $s_a^1 = s_b^1$  and  $s_a^2 \leq_{\text{m.c.}} s_b^2$ . Note that in this situation,  $s_a^2$  and  $s_b^2$  are rooted graphs, so  $\leq_{\text{m.c.}}$  compares rooted graphs. Observe that since  $(\mathcal{B}, \leq_{\text{m.c.}})$  is wqo, then  $(\Sigma, \preceq)$  is wqo. For every  $i \in \mathbb{N}$ , let  $A'_i$  be the  $(\Sigma, \preceq)$ -labeled rooted graph  $(A_i, \sigma_i)$ . We now consider the infinite sequence  $\langle A'_i \rangle_{i \in \mathbb{N}}$ . By our initial assumption,  $(\text{lab}_\Sigma(\mathcal{A}), \leq_{\text{m.c.}})$  is wqo (as  $\mathcal{A}$  consists only in 2-connected graphs), so there is a good pair  $(A'_i, A'_j)$  in the sequence  $\langle A'_i \rangle_{i \in \mathbb{N}}$ .

To complete the proof, we will show that  $A'_i \leq_{\text{m.c.}} A'_j \Rightarrow G_i \leq_{\text{m.c.}} G_j$ . Let  $\mu$  be a contraction model of  $A'_i$  in  $A'_j$ . Then for each cutvertex  $c \in C_i$ ,  $\mu(c)$  contains a vertex  $d \in C_j$  with  $B_c^i \leq_{\text{m.c.}} B_d^j$ . Let  $\mu_c$  denote a root-preserving contraction model of  $B_c^i$  onto  $B_d^j$ . We construct a model  $g$  as follows:

$$\nu: \begin{cases} V(G_i) & \rightarrow \mathcal{P}(V(G_j)) \\ v & \mapsto \mu(v) \text{ if } v \in A_i \setminus C_i \\ v & \mapsto \mu_c(v) \text{ if } v \in B_c^i \setminus C_i \\ v & \mapsto \mu(v) \cup \mu_v(v) \text{ if } v \in C_i \end{cases}$$

We now prove that  $\nu$  is a contraction model of  $G_i$  onto  $G_j$ . First note that by definition of  $\mu$  and each  $\mu_c$ , we have  $\nu(u) \cap \nu(v) = \emptyset$ , for any pair of distinct vertices  $u$  and  $v$  in  $G_i$ , and also every vertex of  $G_j$  is in the image of some vertex of  $G_i$  (items **(M1)** and **(M7)** in the definition of a contraction model). If  $u \in C_i$ , then  $\mu(u)$  contains a vertex  $v \in C_j$  for which  $B_u^i \leq_{\text{m.c.}} B_v^j$  and  $v$  is also contained in  $\mu_v(v)$  since  $\mu_v$  preserves roots. Thus,  $G_j[\nu(u)]$  is connected when  $u \in C_i$  (point **(M2)**). This is obviously true when  $u \notin C_i$  again by the definitions of  $\mu$  and each  $\mu_c$ . Moreover, the endpoints of every edge of  $G_i$  belong either both to  $A_i$ , or both to  $B_c^i$ , so point **(M6)** follows from the properties of  $\mu$  and each  $\mu_c$ . Finally, as the labeling  $\sigma$  ensures that  $\text{root}(G_j) \in \nu(\text{root}(G_i))$ , we establish that  $G_i \leq_{\text{m.c.}} G_j$ . So  $\langle G_i \rangle_{i \in \mathbb{N}}$  has a good pair  $(G_i, G_j)$ , a contradiction.  $\square$

Let us now summarize the tools introduced in this section.

**Lemma 3.33** (Summary of **Section 3.3**). *If  $(A, \preceq_A)$  and  $(B, \preceq_B)$  are wqos, then*

- (i)  $(C, \preceq_A)$  is a wqo for every  $C \subseteq A$  (**Remark 3.24**);
- (ii)  $(A \cup B, \preceq_A \cup \preceq_B)$  is a wqo (**Remark 3.25**);
- (iii)  $(A \times B, \preceq_A \times \preceq_B)$  is a wqo (**Lemma 3.26**);
- (iv)  $(A^*, \preceq^*)$  is a wqo (**Theorem 3.1**);
- (v)  $(\mathcal{P}^{<\omega}(A), \preceq^{\mathcal{P}})$  is a wqo (**Corollary 3.27**);
- (vi)  $(C, \preceq_C)$  is a wqo, for every epi  $f: (A, \preceq_A) \rightarrow (C, \preceq_C)$  (**Remark 3.28**);
- (vii)  $(\{y \in C, x' \not\preceq_C y\}, \preceq_C)$  is a wqo whenever  $(\{y \in C, x \not\preceq_C y\}, \preceq_C)$  is, for every  $x', x$  elements of a qoset  $(C, \preceq_C)$  such that  $x' \preceq x$  (**Remark 3.30**);
- (viii) if for any wqo  $(C, \preceq_C)$  the 2-connected graphs of a graph class  $\mathcal{H}$  labeled by  $(C, \preceq_C)$  are wqo by  $\leq_{\text{i.m.}}$ , then  $(\mathcal{H}, \leq_{\text{i.m.}})$  is a wqo (**Lemma 3.31**);
- (ix) if for any wqo  $(C, \preceq_C)$  the 2-connected multigraphs of a multigraph class  $\mathcal{H}$  labeled by  $(C, \preceq_C)$  are wqo by  $\leq_{\text{m.c.}}$ , then  $(\mathcal{H}, \leq_{\text{m.c.}})$  is a wqo (**Lemma 3.32**).

In the next section we give the main lines of the proofs of **Theorem 3.12**, **Theorem 3.13**, and **Theorem 3.14**.

## 3.4 A high-level view of the proofs

The techniques that we use in the following sections mostly follow a general scheme. The purpose of this section is to provide an informal description of its steps. The general setting is the following: we are given a qoset  $(S, \preceq)$  and the goal is to show that it is a wqo.

Step 1: Define a *compact representation* of the elements of  $S$ . More formally, we choose an injective function  $f$  from  $S$  to some other set  $S'$ . The intuition is that, usually,  $S$  is a subset of a wider set and its elements only span a *restricted area* of this set. Therefore we would like to *project* these elements into a smaller set, that will be easier to well-quasi-order afterwards. This representation can be seen as an encoding, as we will sometime break a graph into parts and represent it as a tuple containing these parts. It may be deduced from structural information about the graphs of  $S$ .

Step 2: Find a quasi-order relation  $\preceq'$  on  $S'$  with good properties. This relation should be chosen such that whenever  $f(x) \preceq' f(y)$  holds, we have  $x \preceq y$ , for every  $x, y \in S$ . We should stress that the choice of  $f$  must be done with this step in mind.

Step 3: The third and last step is to show that  $(S', \preceq')$  is a wqo. This can be done by applying standard results on well-quasi-orders, if both  $S'$  and  $\preceq'$  have been chosen carefully.

Finally, we obtain that  $f$  is a epi, the domain of which is a subset of the wqo  $(S', \preceq')$ . According to [Remark 3.24](#) and [Remark 3.28](#), we get that  $(S, \preceq)$  is a wqo, as desired. As an illustration of this scheme, let us prove the following easy lemma.

**Lemma 3.34.** *Let  $\mathcal{H}$  be a class of connected graphs and let  $\mathcal{G}$  the class of graphs, the connected components of which belong to  $\mathcal{H}$ . If  $(\mathcal{H}, \leq_{\text{i.sg.}})$  is a wqo, then so is  $(\mathcal{G}, \leq_{\text{i.sg.}})$ .*

*Proof.* Let us follow the steps detailed above.

Step 1: We represent every graph  $G$  of  $\mathcal{G}$  as a sequence  $f(G)$  listing its connected components taken in an arbitrary order. The function  $f: \mathcal{G} \rightarrow \mathcal{H}^*$  thus defined is clearly injective.

Step 2: We notice that for every  $G, G' \in \mathcal{G}$ , if  $f(G) \leq_{\text{i.sg.}}^* f(G')$  then  $G \leq_{\text{i.sg.}} G'$ . Therefore the quasi-order relation  $\leq_{\text{i.sg.}}^*$  suits our needs.

Step 3: We show that the codomain of  $f$  is a wqo. For this, we apply Higman's Lemma ([Theorem 3.1](#)) to  $(\mathcal{H}, \leq_{\text{i.sg.}})$ , which is a wqo by our initial assumption.

As explained above, this implies that  $(\mathcal{G}, \leq_{\text{i.sg.}})$  is a wqo and we are done.  $\square$

Let us end this section with a simple but crucial observation in the study of the well-quasi-orderability of classes that are defined by forbidden structures (of any kind). If none of the elements of an infinite antichain  $A$  contains some element  $x$ , then excluding  $x$  does not give a well-quasi-order. Indeed, the class obtained still contains the infinite antichain  $A$ . Let us formally restate this observation.

*Observation 3.35.* Let  $(S, \preceq)$  be a qoset and let  $A \subseteq S$  be an infinite antichain wrt.  $\preceq$ . Let  $x \in S$ . If  $\{y \in S, x \not\preceq y\}$  is well-quasi-ordered by  $\preceq$ , then all but finitely many elements of  $A$  are larger than  $x$  wrt.  $\preceq$ .

For this reason, we devote the next section to a study of infinite antichains for the orderings considered in this chapter.

### 3.5 The bestiary

It is worth noting that none of the orderings we consider in this chapter admits an infinite decreasing sequence. Indeed, for all these relations, the sum of the number of vertices and edges, which is a non-negative integer, is decreasing when considering smaller graphs. Hence, every decreasing sequence of graphs yields a decreasing sequence of positive integers of the same length, which is always finite. Therefore the only obstacle for these relations to be well-quasi-orders is the presence of an infinite antichain.

As noted in [Observation 3.35](#), the study of infinite antichains may provide helpful information when proving dichotomy theorems. In this section, we present several antichains for the orderings we consider.

#### 3.5.1 Antichains for induced minors




Figure 3.5: Thomas' alternating double wheels.

In 1985, Thomas [[Tho85](#)] presented an infinite sequence of planar graphs (also mentioned later in [[RS93](#)]) and proved that it is an antichain for induced minors. He showed that this relation does not well-quasi-order planar graphs. The elements of this antichain, called *alternating double wheels*, are constructed from an even cycle by adding two nonadjacent vertices and connecting one to one color class of the cycle and connecting the other vertex to the other color class (cf. [Figure 3.5](#) for the three first such graphs). This infinite antichain shows that  $(\text{Excl}_{\leq_{i.m.}}(K_5), \leq_{i.m.})$  is not a wqo since no alternating double wheel contains  $K_5$  as (induced) minor. As a consequence,  $(\text{Excl}_{\leq_{i.m.}}(H), \leq_{i.m.})$  is not a wqo as soon as  $H$  contains  $K_5$  as induced minor.

Therefore, in the quest for all graphs  $H$  such that  $(\text{Excl}_{\leq_{i.m.}}(H), \leq_{i.m.})$  is wqo, we can focus the cases where  $H$  is  $K_5$ -induced minor-free.

Let  $K_5^-$  denote the complete graph on five vertices where one edge has been deleted. The infinite antichain  $\mathcal{A}_M$ , depicted in [Figure 3.6](#), has been introduced in [[MNT88](#)]. Notice that graphs in this antichain have bounded maximum degree. As illustrated in [Figure 3.7](#), these graphs are also planar. The authors of [[MNT88](#)] furthermore proved that no graph of  $\mathcal{A}_M$  contains  $K_5^-$  as induced minor. Similarly as the above remark, we deduce that if  $(\text{Excl}_{\leq_{i.m.}}(H), \leq_{i.m.})$  is a wqo then  $K_5^- \not\leq_{i.m.} H$ .

An *interval graph* is the intersection graph of segments of  $\mathbb{R}$ . A well-known property of interval graphs that we will use later is that they do not contain  $C_4$  as induced minor. In order to show that interval graphs are not wqo by  $\leq_{i.m.}$ , Ding introduced




Figure 3.6: The infinite antichain  $\mathcal{A}_M$  of Matoušek, Nešetřil, and Thomas.

in [Din98] an infinite sequence of graphs defined as follows. For every  $n \in \mathbb{N}$ ,  $n > 2$ , let  $T_n$  be the set of closed intervals

- $[i, i]$  for  $i$  in  $\llbracket -2n, -1 \rrbracket \cup \llbracket 1, 2n \rrbracket$ ;
- $[-2, 2], [-4, 1], [-2n + 1, 2n], [-2n + 1, 2n - 1]$ ;
- $[-2i + 1, 2i + 1]$  for  $i$  in  $\llbracket 1, n - 2 \rrbracket$ ;
- $[-2i, 2i - 2]$  for  $i$  in  $\llbracket 3, n \rrbracket$ .

Figure 3.8 depicts the intervals of  $T_6$ : the real axis (solid line) is folded up and an interval  $[a, b]$  is represented by a dashed line between  $a$  and  $b$ .

For every  $n \in \mathbb{N}$ ,  $n > 2$ , let  $A_n^D$  be the intersection graph of segments of  $T_n$ . Let  $\mathcal{A}_D = \langle A_n^D \rangle_{n>2}$ . Ding proved in [Din98] that  $\mathcal{A}_D$  is an antichain for  $\leq_{i.m.}$ , thus showing that interval graphs are not wqo by induced minors.

Let us now present two more infinite antichains. Let  $\mathcal{A}_{\overline{C}} = \langle \overline{C}_n \rangle_{n \geq 5}$  be the sequence of *antiholes* (i.e. complements of cycles of order at least five), whose first elements are represented in Figure 3.9.

**Lemma 3.36.**  $\mathcal{A}_{\overline{C}}$  is an antichain for induced minors.

*Proof.* Let  $\overline{\deg}$  be the function that maps a vertex  $v$  of a graph  $G$  to the value  $|V(G)| - \deg(v)$ , that is, the number of vertices (including  $v$ ) that are not adjacent to  $v$ . Remark that performing edge contractions and vertex deletions in a graph can only decrease the value of  $\overline{\deg}$  of a given vertex. Indeed, during such an operation, the order of the graph decreases by exactly one whereas the degree of a vertex decreases by at most one. Also notice that in every graph  $G$  of  $\mathcal{A}_{\overline{C}}$ , every vertex  $v$  satisfies  $\overline{\deg}(v) = 3$ . An edge contraction in  $G$  yields a vertex  $v$  such that  $\overline{\deg}(v) < 3$  whereas a vertex deletion gives a vertex  $v$  with  $\overline{\deg}(v) = 2$ . By an above remark, this value cannot be increased by further edge contractions or vertex deletions. Therefore there is no sequence of edge contractions and vertex deletions on  $G$  yielding an other graph of  $\mathcal{A}_{\overline{C}}$ . This proves that  $\mathcal{A}_{\overline{C}}$  is an antichain wrt.  $\leq_{i.m.}$   $\square$




Figure 3.7: A planar drawing (right) of the third graph of  $\mathcal{A}_M$  (left).




Figure 3.8: An illustration of the intervals in  $T_6$ .

We will meet again the antichain  $\mathcal{A}_{\bar{C}}$  in the proof of [Theorem 3.12](#). Another infinite antichain which shares with  $\mathcal{A}_M$  the properties of planarity and bounded maximum degree is the antichain of nested lozenges depicted in [Figure 3.10](#). We will not go more into detail about it here as this antichain will not be used in our proofs.

### 3.5.2 Antichains for contractions

Let us first mention that since every contraction of a graph is also an induced minor, all the antichains for induced minors presented in [Subsection 3.5.1](#) are also antichains for the contraction relation. Not surprisingly there are more antichains for the contraction ordering. The most simple is certainly the class  $\mathcal{A}_{\bar{K}} = \{\bar{K}_i\}_{i \in \mathbb{N}}$  of edgeless graphs. We present here one more infinite antichain for contractions, which is the class of complete bipartite graphs with one part of size two:  $\mathcal{A}_K = \{K_{2,r}, r \in \mathbb{N}_{\geq 2}\}$  (cf. [Figure 3.11](#)).




Figure 3.9: Antiholes antichain.




Figure 3.10: Nested lozenges.

**Lemma 3.37.** *For every  $(p, q), (p', q') \in \mathbb{N}_{\geq 2}$  such that  $p \leq p'$  and  $q < q'$ , there is no contraction model of  $K_{p,q}$  in  $K_{p',q'}$ .*

*Proof.* Let us assume for contradiction that there is a contraction model  $\varphi$  of  $K_{p,q}$  in  $K_{p',q'}$ . As  $K_{p',q'}$  has more vertices than  $K_{p,q}$ , there is a vertex  $v$  of  $K_{p,q}$  such that  $|\varphi(v)| \geq 2$ . Observe that every subset of at least two vertices of  $K_{p,q}$  that induced a connected subgraph is dominating. Indeed, such a subset must contain at least a vertex from each part of the bipartition. It follows from the definition of a contraction model that  $K_{p',q'}$  has a dominating vertex, a contradiction. Therefore there is no contraction model of  $K_{p,q}$  in  $K_{p',q'}$ .  $\square$

**Corollary 3.38.**  $\{K_{2,p}, p \in \mathbb{N}_{\geq 2}\}$  is an antichain of  $\leq_c$ .

If we allow multiple edges and consider the multigraph contraction ordering, then the sequence of multigraphs  $\mathcal{A}_\theta = \{\theta_i\}_{i \in \mathbb{N}^*}$  depicted in Figure 3.12 is also an infinite antichain.

As we will show in Subsection 3.8.3, every infinite antichain for multigraph contraction is mainly composed of  $\mathcal{A}_{\overline{K}}$  and  $\mathcal{A}_\theta$ .

## 3.6 Induced minors and well-quasi-ordering

In this section, we prove the dichotomy Theorem 3.12. Let us first recall this result.

**Theorem 3.12** (Dichotomy for induced minors). *Let  $H$  be a graph. The class of  $H$ -induced minor-free simple graphs is wqo by  $\leq_{i.m.}$  iff  $H \leq_{i.m.} \hat{K}_4$  or  $H \leq_{i.m.} \text{Gem}$ .*




Figure 3.11: The antichain  $\mathcal{A}_K$ .




Figure 3.12: The multiedges antichain.

Our proof naturally has two parts: for different values of  $H$ , we need to show wqo of  $H$ -induced minor-free graphs or exhibit an  $H$ -induced minor-free antichain. We recall that, unless otherwise specified, this section deals with simple graphs.

In order to achieve **Step 1** presented in **Section 3.4**, we first prove two decomposition theorems. The following two theorems describe the structure of graphs with  $H$  forbidden as an induced minor, when  $H$  is  $\hat{K}_4$  and the Gem, respectively. Recall that the prism is the Cartesian product of  $K_3$  and  $K_2$ .

**Theorem 3.39** (Decomposition of  $\hat{K}_4$ -induced minor-free graphs). *Let  $G$  be a 2-connected graph such that  $\hat{K}_4 \not\leq_{\text{i.m.}} G$ . Then:*

- either  $K_4 \not\leq_{\text{i.m.}} G$ ;
- or  $G$  is a subdivision of a graph among  $K_4$ ,  $K_{3,3}$ , and the prism;
- or  $V(G)$  has a partition  $(C, M)$  such that  $G[C]$  is an induced cycle,  $G[M]$  is a complete multipartite graph and every vertex of  $C$  is either adjacent in  $G$  to all vertices of  $M$ , or to none of them.

**Theorem 3.40** (Decomposition of Gem-induced minor-free graph). *Let  $G$  be a 2-connected graph such that  $\text{Gem} \not\leq_{\text{i.m.}} G$ . Then  $G$  has a subset  $X \subseteq V(G)$  of at most six vertices such that every connected component of  $G \setminus X$  is either a cograph or a path whose internal vertices are of degree two in  $G$ .*

Using the two above structural results, we are able to show the well-quasi-ordering of the two classes with respect to induced minors by following the steps described in **Section 3.4**.

**Theorem 3.41.** *The class of  $\hat{K}_4$ -induced minor-free graphs is wqo by  $\leq_{\text{i.m.}}$ .*

**Theorem 3.42.** *The class of Gem-induced minor-free graphs is wqo by  $\leq_{\text{i.m.}}$ .*

**Organization of the proof.** Subsection 3.6.1 is devoted to the proof of Theorem 3.12, assuming Theorem 3.41 and Theorem 3.42, the proofs of which are respectively given in Subsection 3.6.2 and Subsection 3.6.3.

Let us first provide some definitions specific to this section. An induced subgraph of a graph  $G$  is said to be *basic in  $G$*  if it is either a cograph, or an induced path whose internal vertices are of degree two in  $G$ .

**Complete multipartite graphs.** A graph  $G$  is said to be *complete multipartite* if its vertex set can be partitioned into sets  $V_1, \dots, V_k$  (for some positive integer  $k$ ) in a way such that two vertices of  $G$  are adjacent iff they belong to different  $V_i$ 's. The class of complete multipartite graphs is referred to as  $\mathcal{K}_{\mathbb{N}^*}$ .

**Containing  $K_4$ -subdivisions.** A graph  $G$  contains  $K_4$  as an induced minor if and only if  $G$  contains  $K_4$ -subdivision as a subgraph. This equivalence is highly specific to the graph  $K_4$  and in general neither implication would be true. We will freely change between those two notions for containing  $K_4$ , depending on which one is more convenient in the given context.

A graph  $G$  will be said to contain a *proper  $K_4$ -subdivision*, if there is some vertex  $v \in V(G)$ , such that  $G \setminus v$  contains a  $K_4$ -subdivision.

**Cycle-multipartite.** Given a graph  $G$ , a pair  $(C, R)$  of induced subgraphs of  $G$  is said to be a *cycle-multipartite decomposition* of  $G$  if the following conditions are satisfied:

- (i)  $(V(C), V(R))$  is a partition of  $V(G)$ ;
- (ii)  $C$  is a cycle and  $R$  is a complete multipartite graph;
- (iii)  $\forall u, v \in V(R), N_C(u) = N_C(v)$ .

The class of graphs having cycle-multipartite decomposition is denoted by  $\mathcal{W}$ .

**Cuts.** In a graph  $G$ , a  $K_2$ -cut (respectively  $\overline{K_2}$ -cut) is a subset  $S \subseteq V(G)$  such that  $G - S$  is not connected and  $G[S]$  is isomorphic to  $K_2$  (respectively  $\overline{K_2}$ ).

### 3.6.1 The dichotomy theorem

The purpose of this section is to prove Theorem 3.12, that is, to characterize all graphs  $H$  such that  $(\text{Excl}_{\leq_{i.m.}}(H), \leq_{i.m.})$  is a wqo. To this end, we will assume Theorem 3.41 and Theorem 3.42, which we will prove later, in Subsection 3.6.2 and Subsection 3.6.3 respectively.

The main ingredients of the proof are the infinite antichains presented in Subsection 3.5.1, together with Theorem 3.41 and Theorem 3.42. Infinite antichains will be used to discard every graph  $H$  that is not induced minor of all but finitely many elements of some infinite antichain. On the other hand, knowing that  $(\text{Excl}_{\leq_{i.m.}}(H), \leq_{i.m.})$

is a wqo gives that  $(\text{Excl}_{\leq_{i.m.}}(H'), \leq_{i.m.})$  is a wqo for every  $H' \leq_{i.m.} H$ , by the virtue of [Remark 3.30](#).

In the statement of the following results, we assume that  $H$  is any graph. Recall that a linear forest is a disjoint union of paths.

**Lemma 3.43.** *If  $(\text{Excl}_{\leq_{i.m.}}(H), \leq_{i.m.})$  is a wqo then  $\overline{H}$  is a linear forest.*

*Proof.* Let us show that  $\text{Excl}_{\leq_{i.m.}}(H)$  has an infinite antichain given that  $\overline{H}$  is not a linear forest. We will in fact show that  $\text{Excl}_{\leq_{i.m.}}(H)$  contains infinitely many antiholes. If  $\overline{H}$  is not a linear forest, then it has a vertex of degree at least 3, or it contains an induced cycle as induced subgraph.

*First case:*  $\overline{H}$  has a vertex  $v$  of degree 3. Let  $x, y, z$  be three neighbors of  $v$ . In the graph  $H[\{v, x, y, z\}]$ , the vertex  $v$  is adjacent to none of  $x, y, z$ . In an antihole (c.f. [Figure 3.9](#)), every vertex has exactly two non-neighbors, so  $H[\{v, x, y, z\}]$  is not an induced minor of any element of  $\mathcal{A}_{\overline{C}}$ . Therefore  $\mathcal{A}_{\overline{C}} \subseteq \text{Excl}_{\leq_{i.m.}}(H)$ .

*Second case:*  $\overline{H}$  contains has a cycle. Let  $k$  be the length of an induced cycle of  $\overline{H}$ . Then we have  $\overline{C}_k \leq_{i.m.} H$ . Let us start with the case where  $k \geq 5$ , and let us consider  $\overline{C}_n$  for some  $n > |H|$ . By the choice of  $n$ , we have  $\overline{C}_n \not\leq_{i.m.} H$ . On the other hand, if  $H \leq_{i.m.} \overline{C}_n$ , then by the fact that  $\overline{C}_k \leq_{i.m.} H$  and transitivity of  $\leq_{i.m.}$ , we would have  $\overline{C}_k \leq_{i.m.} \overline{C}_n$ , which would yield a contradiction with the fact that  $\mathcal{A}_{\overline{C}}$  is an antichain. Hence  $\mathcal{A}_{\overline{C}} \cap \text{Excl}_{\leq_{i.m.}}(H)$  contains all antiholes of size greater than  $|H|$ , and in particular is infinite as required. Let us now consider the case where  $k \in \{3, 4\}$ . Every vertex of an antihole  $A$  has only two non-neighbors, which are adjacent, therefore  $A$  does not contain  $\overline{C}_3$  (i.e. three independent vertices) as an induced subgraph. Moreover, as the neighbors of two distinct vertices of a cycle are distinct, the non-neighbors of two distinct vertices of  $A$  are also distinct and we deduce that  $A$  does not have a pair of independent edges, i.e.  $\overline{C}_4 \not\leq_{i.sg.} A$ . Observe that if  $J$  has maximum degree one, then  $J \leq_{i.sg.} G$  iff  $J \leq_{i.m.} G$ , for any graph  $G$ . Therefore  $\overline{C}_3 \not\leq_{i.sg.} A$  and  $\overline{C}_4 \not\leq_{i.sg.} A$ , and we have  $A \in \text{Excl}_{\leq_{i.m.}}(H)$ . This holds for every antihole  $A$ , hence  $\mathcal{A}_{\overline{C}} \subseteq \text{Excl}_{\leq_{i.m.}}(H)$  and we are done.  $\square$

Due to the interesting properties on  $\overline{H}$  given by [Lemma 3.43](#), we will be led below to work with this graph rather than with  $H$ . The following lemma presents step by step the properties that we can deduce on  $\overline{H}$  by assuming that  $\text{Excl}_{\leq_{i.m.}}(H)$  is wqo by  $\leq_{i.m.}$ . Recall that we denote by  $\text{cc}(G)$  the number of connected components of graph  $G$ .

**Lemma 3.44.** *If  $(\text{Excl}_{\leq_{i.m.}}(H), \leq_{i.m.})$  is a wqo, then we have*

- (R1)  $\overline{H}$  has at most 4 connected components;
- (R2) at most one connected component of  $\overline{H}$  is not a single vertex;
- (R3) the largest connected component of  $\overline{H}$  has at most 4 vertices;
- (R4) if  $n = |V(H)|$  and  $c = \text{cc}(\overline{H})$  then  $n \leq 7$  and  $\overline{H} = (c - 1) \cdot K_1 + P_{n-c+1}$ ;
- (R5) if  $\text{cc}(\overline{H}) = 3$  then  $|V(H)| \leq 5$ .

(R6) if  $\text{cc}(\overline{H}) = 4$  then  $|V(H)| \leq 4$ .

*Proof.* *Proof of item (R1).* The infinite antichain  $\mathcal{A}_M$  does not contain  $K_5$  as an (induced) minor, hence  $K_5 \not\leq_{\text{i.m.}} H$  and so  $\overline{H}$  does not contain  $5 \cdot K_1$  as induced minor. Therefore it has at most 4 connected components.

*Proof of items (R2) and (R3).* The infinite antichain  $\mathcal{A}_D$  does not contain  $C_4$  as induced minor (as it is an interval graph), hence neither does  $H$ . Therefore  $\overline{H}$  does not contain  $2 \cdot P_2$  as induced minor. This implies that  $\overline{H}$  does not contain  $P_5$  as induced minor and that given two connected components of  $\overline{H}$  at least one must be of order one. As connected components of  $H$  are (induced) paths (by Lemma 3.43), the largest connected component of  $H$  has order at most 4.

Item (R4) follows from the above proofs and from the fact that  $\overline{H}$  is a linear forest. *Proof of item (R5).* Similarly as in the proof of item (R1),  $\mathcal{A}_M$  does not contain  $K_5^-$  as induced minor so  $\overline{K_5^-} = K_2 + 3 \cdot K_1$  is not an induced minor of  $\overline{H}$ . If we assume that  $\text{cc}(\overline{H}) = 3$  and  $|V(H)| \geq 6$ , the largest component of  $\overline{H}$  is a path on (a least) 4 vertices, so it contains  $K_1 + K_2$  as induced subgraph. Together with the two other (single vertex) components, this gives an  $K_2 + 3 \cdot K_1$  induced minor, a contradiction. *Proof of item (R6).* Let us assume that  $\text{cc}(\overline{H}) = 4$ . If the largest connected component has more than one vertex, then  $\overline{H}$  contains  $K_2 + 3 \cdot K_1$  as an induced minor, which is not possible (as in the proof of item (R5)). Therefore  $\overline{H} = 4 \cdot K_1$  and so  $|V(\overline{H})| = 4$ .  $\square$

We are now able to describe more precisely graphs  $H$  for which  $(\text{Excl}_{\leq_{\text{i.m.}}}(H), \leq_{\text{i.m.}})$  could be a wqo. Let  $K_3^+$  be the complement of  $P_3 + K_1$  and recall that we call  $D_2$  the complement of  $K_2 + 2 \cdot K_1$ , which is also the graph obtained from  $K_4$  by deleting an edge (sometimes referred as *diamond graph*).

**Lemma 3.45.** *If  $(\text{Excl}_{\leq_{\text{i.m.}}}(H), \leq_{\text{i.m.}})$  is a wqo, then  $H \leq_{\text{i.m.}} \hat{K}_4$  or  $H \leq_{\text{i.m.}} \text{Gem}$ .*

*Proof.* Using the information on  $\overline{H}$  given by Lemma 3.44, we can build a table of possible graphs  $\overline{H}$  depending on  $\text{cc}(\overline{H})$  and  $|V(\overline{H})|$ . Table 3.2 is such a table: each column corresponds to a number of connected components (between one and four according to item (R1)) and each line corresponds to an order (at most seven, by item (R4)). A grey cell means either that there is no such graph (for instance a graph with one vertex and two connected components), or that for all graphs  $\overline{H}$  matching the number of connected components and the order associated with this cell, the qoset  $(\text{Excl}_{\leq_{\text{i.m.}}}(H), \leq_{\text{i.m.}})$  is not a wqo.

From Table 3.2 we can easily deduce Table 3.3 of the corresponding graphs.

Remark that we have

- $K_1 \leq_{\text{i.m.}} 2 \cdot K_1 \leq_{\text{i.m.}} K_2 + K_1 \leq_{\text{i.m.}} P_4 \leq_{\text{i.m.}} \text{Gem}$ ;
- $K_2 \leq_{\text{i.m.}} P_3 \leq_{\text{i.m.}} K_3^+ \leq_{\text{i.m.}} \text{Gem}$ ;
- $K_3 \leq_{\text{i.m.}} D_2 \leq_{\text{i.m.}} \hat{K}_4$ ; and
- $K_4 \leq_{\text{i.m.}} \hat{K}_4$ .

This concludes the proof.  $\square$

$ V(H)  \setminus \text{cc}(\overline{H})$	1	2	3	4
1	$K_1$			
2	$K_2$	$2 \cdot K_1$		
3	$P_3$	$K_2 + K_1$	$3 \cdot K_1$	
4	$P_4$	$P_3 + K_1$	$K_2 + 2 \cdot K_1$	$4 \cdot K_1$
5	(R3)	$P_4 + K_1$	$P_3 + 2 \cdot K_1$	(R6)
6	(R3)	(R3)	(R5)	(R6)
7	(R3)	(R3)	(R5)	(R6)

Table 3.2: If  $(\text{Excl}_{\leq_{i.m.}}(H), \leq_{i.m.})$  is a wqo, then  $\overline{H}$  belongs to this table.

$ V(H)  \setminus \text{cc}(\overline{H})$	1	2	3	4
1	$K_1$			
2	$2 \cdot K_1$	$K_2$		
3	$K_2 + K_1$	$P_3$	$K_3$	
4	$P_4$	$K_3^+$	$D_2$	$K_4$
5	(R3)	Gem	$\hat{K}_4$	(R6)
6	(R3)	(R3)	(R5)	(R6)
7	(R3)	(R3)	(R5)	(R6)

Table 3.3: If  $(\text{Excl}_{\leq_{i.m.}}(H), \leq_{i.m.})$  is a wqo, then  $H$  belongs to this table.

We are now ready to give the proof of **Theorem 3.12** that we first restate.

**Theorem 3.12** (Dichotomy for induced minors). *Let  $H$  be a graph. The class of  $H$ -induced minor-free simple graphs is wqo by  $\leq_{i.m.}$  iff  $H \leq_{i.m.} \hat{K}_4$  or  $H \leq_{i.m.} \text{Gem}$ .*

*Proof.* If  $H \not\leq_{i.m.} \text{Gem}$  and  $H \not\leq_{i.m.} \hat{K}_4$ , then by **Lemma 3.45**  $(\text{Excl}_{\leq_{i.m.}}(H), \leq_{i.m.})$  is not a wqo. On the other hand, by **Theorem 3.41** and **Theorem 3.42** we know that both  $\text{Excl}_{\leq_{i.m.}}(\hat{K}_4)$  and  $\text{Excl}_{\leq_{i.m.}}(\text{Gem})$  are wqo by  $\leq_{i.m.}$ . Consequently, by **Remark 3.30**,  $(\text{Excl}_{\leq_{i.m.}}(H), \leq_{i.m.})$  is wqo as soon as  $H \leq_{i.m.} \text{Gem}$  or  $H \leq_{i.m.} \hat{K}_4$ .  $\square$

### 3.6.2 Graphs not containing $\hat{K}_4$

The main goal of this section is to provide a proof to the following theorem, that we already stated above.

**Theorem 3.41.** *The class of  $\hat{K}_4$ -induced minor-free graphs is wqo by  $\leq_{i.m.}$ .*

To this purpose, we first prove that the graphs of  $\text{Excl}_{\leq_{i.m.}}(\hat{K}_4)$  admit a simple structural decomposition. This structure is then used to show that graphs of  $\text{Excl}_{\leq_{i.m.}}(\hat{K}_4)$  are well-quasi-ordered by the relation  $\leq_{i.m.}$ .

### A decomposition theorem for $\text{Excl}_{\leq i.m.}(\hat{K}_4)$

The main topic of this section is the proof of [Theorem 3.39](#). This theorem states that every graph in the class  $\text{Excl}_{\leq i.m.}(\hat{K}_4)$ , either does not have even  $K_4$  as induced minor, or is a subdivision of some small graph, or has a cycle-multipartite decomposition. Most of the time, we show that some property  $P$  is not satisfied by graphs of  $\text{Excl}_{\leq i.m.}(\hat{K}_4)$  by showing an induced minor model of  $\hat{K}_4$  in graphs satisfying  $P$ . We first assume that  $G$  contains a proper  $K_4$ -subdivision, and we show in [Lemma 3.62](#) how to deal with the other case.

**Lemma 3.46.** *If  $G$  contains as induced minor any graph  $H$  consisting of:*

- *a  $K_4$ -subdivision  $S$ ;*
- *an extra vertex  $x$  linked by exactly two paths  $L_1$  and  $L_2$  to two distinct vertices  $s_1, s_2 \in V(S)$ , where the only common vertex of  $L_1$  and  $L_2$  is  $x$ ;*
- *and possibly extra edges between the vertices of  $S$ , or between  $L_1$  and  $L_2$ , or between the interior of the paths and  $S$ ,*

*then  $\hat{K}_4 \leq_{i.m.} G$ .*

*Proof.* We assume that  $L_1$  and  $L_2$  do not share internal vertices with  $S$ , what we can do without loss of generality by considering the paths of minimum length satisfying the requirements of the lemma. Let us call  $V = \{v_1, v_2, v_3, v_4\}$  the non-subdivision vertices of  $S$ , i.e. vertices corresponding to vertices of  $K_4$ . We present here a sequence of edge contractions in  $H$  leading to  $\hat{K}_4$ . Let us iteratively repeat the following procedure: as long as there is a path on at least two edges between two elements of  $V \cup \{s_1, s_2, x\}$  that is internally disjoint with this set, contract the whole path to a single edge.

Once we can not apply this contraction any more, we end up with a graph that has two parts: the  $K_4$ -subdivision with at most 2 subdivisions (with vertex set  $V \cup \{s_1, s_2\}$ ) and the vertex  $x$ , which is now only adjacent to  $s_1$  and  $s_2$ .

*First case:*  $s_1, s_2 \in V$ . The graph  $H$  is isomorphic to  $\hat{K}_4$ : it is  $K_4$  plus a vertex of degree two.

*Second case:*  $s_1 \in V$  and  $s_2 \notin V$  (and the symmetric case). As vertices of  $V$  are the only vertices of  $H$  that have degree 3 in  $S$ ,  $s_2$  is of degree 2 in  $S$  (it is introduced by subdivision). The contraction of the edge between  $s_2$  and one of its neighbors in  $S$  that is different to  $s_1$  leads to first case.

*Third case:*  $s_1, s_2 \notin V$ . As in the second case, these two vertices have degree two in  $S$ . Since no two different edges of  $K_4$  can have the same endpoints, the neighborhoods of  $s_1$  and  $s_2$  have at most one common vertex. Then for every  $i \in \{1, 2\}$  there is a neighbor  $t_i$  of  $s_i$  that is not adjacent to  $s_{3-i}$ . Contracting the edges  $\{s_1, t_1\}$  and  $\{s_2, t_2\}$  leads to first case.  $\square$

**Corollary 3.47.** *Let  $G \in \text{Excl}_{\leq i.m.}(\hat{K}_4)$  be a 2-connected graph containing a proper  $K_4$ -subdivision. For every subdivision  $S$  of  $K_4$  in  $G$ , and for every vertex  $x \in V(G) \setminus V(S)$ ,  $N_S(x) \geq 3$ .*

*Proof.* Let  $S$  be a proper  $K_4$ -subdivision in  $G$  and  $x \in V(G) \setminus V(S)$ . Let  $L_1, L_2$  be two shortest paths from  $x$  to  $S$  meeting only in  $x$ . Such paths exist by the 2-connectivity of  $G$ . Remark that if  $|N_G(x) \cap V(S)| \leq 2$ , then the graph induced by  $S, L_1$ , and  $L_2$  satisfies conditions of [Lemma 3.46](#). Therefore,  $N_S(x) \geq 3$ .  $\square$

*Remark 3.48.* For every two edges of  $K_4$  there is a Hamiltonian cycle using these edges.

*Remark 3.49.* Three edges of  $K_4$  are not contained into a same cycle iff they are incident with the same vertex.

Recall that a *3-wheel* a graph that can be obtained by connecting three distinct vertices of a cycle to a new vertex.

**Lemma 3.50.** *Every 2-connected graph  $G \in \text{Excl}_{\leq i.m.}(\hat{K}_4)$  containing a proper  $K_4$ -subdivision has a 3-wheel as subgraph.*

*Proof.* Let  $S$  be a minimum (proper)  $K_4$ -subdivision in  $G$  and let  $x \in V(G) \setminus V(S)$ . We define  $V$  as in the proof of [Lemma 3.46](#) and we say that two neighbours of  $x$  in  $S$  are equivalent if they lie on the same path between two elements of  $V$  (intuitively they correspond to the same edge of  $K_4$ ). By [Corollary 3.47](#), we only have to consider the case  $|N_S(x)| \geq 3$ .

First of all, observe that if some three neighbors of  $x$  lie on a cycle of  $S$ , then we are done. Let us assume from now on, that there is no cycle of  $S$  containing three neighbors of  $x$ . This implies that no two neighbors of  $x$  are equivalent (by [Remark 3.48](#)), no neighbor of  $x$  belongs to  $V$  (by the same remark), and that  $|N_S(x)| = 3$  (by [Remark 3.49](#)). Let us consider the induced minor  $H$  of  $S + x$  obtained by contracting all edges not incident with two vertices of  $V \cup N_S(x)$ . By [Remark 3.49](#) and since the three neighbors of  $x$  do not belong to a cycle, there is a vertex of  $V(H) \setminus \{x\}$  adjacent to the three neighbors of  $x$ . Contracting two of the edges incident with this vertex merges two neighbors of  $x$  and the graph we obtain is a  $K_4$ -subdivision (corresponding to  $S$ ) together with a vertex of degree 2 (corresponding to  $x$ ). By [Lemma 3.46](#), this would imply that  $\hat{K}_4 \leq_{i.m.} G$ , a contradiction. Therefore three neighbors of  $x$  lie on a cycle of  $S$  and this concludes the proof.  $\square$

Now we will deal with a graph  $G$  that satisfies conditions of [Lemma 3.50](#); namely:  $G$  is a 2-connected graph, without  $\hat{K}_4$  as an induced minor, containing a proper  $K_4$ -subdivision;  $C$  denotes the cycle of a minimum (in terms of number of vertices) 3-wheel in  $G$ , and  $R$  the graph induced by the remaining vertices.

*Remark 3.51.* As this 3-wheel is a subdivision of  $K_4$  as subgraph of  $G$ , by [Corollary 3.47](#), every vertex of  $R$  has at least three neighbors in  $C$ .

*Remark 3.52.* This 3-wheel contains no more vertices than the  $K_4$ -subdivision that we assumed to be contained in  $G$ . Therefore, every minimum  $K_4$ -subdivision of  $G$  (in terms of vertices) is a 3-wheel.

**Lemma 3.53.** *Let  $G$  be a 2-connected graph of  $\text{Excl}_{\leq i.m.}(\hat{K}_4)$ . Every minimum (in terms of number of vertices) 3-wheel  $W$  of cycle  $C$  and center  $r$  that is a subgraph of  $G$  is such that, if  $C$  is not an induced cycle in  $G$ ,*

- (i) the endpoints of every chord are both adjacent to some  $u \in N_C(r)$ ;
- (ii) every two distinct  $v, w \in N_C(r) \setminus \{u\}$  are adjacent on  $C$ ;
- (iii)  $C$  has exactly one chord in  $G$
- (iv)  $|N_C(r)| = 3$ .




Figure 3.13: Forbidden configurations in the proof of [Lemma 3.53](#).

*Proof.* Let  $u, v, w \in N_C(r)$  be three distinct neighbours of  $r$  in  $C$  and let  $C_u$  be the path of  $C$  between  $v$  and  $w$  that does not contain the vertex  $u$ , and similarly for  $C_v$  and  $C_w$ . First of all, notice that no proper subgraph of  $W$  can be a subdivision of  $K_4$ , otherwise  $G$  would contain a graph smaller than  $W$  but meeting the same requirements, according to [Lemma 3.50](#). Below we will show that when conditions (i)-(iv) are not fulfilled,  $W$  is not minimal, i.e. that when some vertices are deleted in  $W$ , it still contains a  $K_4$ -subdivision. [Figure 3.13](#) illustrates such configurations, where white vertices can be deleted.

Let  $\{x, y\}$  be a chord of  $C$  in  $G$ . Remark that the endpoints of a chord cannot belong to the same path  $C_l$  for any  $l \in \{u, v, w\}$  without violating the minimality of  $S$ , as deleting vertices of  $C_l$  that are between  $x$  and  $y$  would still lead to a 3-wheel (first configuration of [Figure 3.13](#)). Therefore,  $x$  and  $y$  belong to different  $C_l$ 's, say without loss of generality that  $x \in V(C_w)$  and  $y \in V(C_v)$ .

Let us prove that  $x$  and  $y$  must both be adjacent to  $u$ . By contradiction, we assume that, say,  $y$  and  $u$  are not adjacent. Let us consider the induced subgraph of  $W$  obtained by the deletion of the interior of the path  $yC_vu$  (containing at least one vertex). Notice that contracting into an edge each of the paths  $vC_uw$ ,  $vC_wx$ ,  $yC_vw$  and  $rpathtuC_wx$  gives  $K_4$ , a contradiction (cf. the second configuration of [Figure 3.13](#)). The case where  $x$  and  $u$  are not adjacent is symmetric. This proves that every chord of  $C$  in  $G$  has endpoints adjacent to a same neighbor of  $r$  on  $C$ , that is (i).

Now, we show that the path  $C_u$  must be an edge. To see this, assume by contradiction that it has length at least 3. The subgraph of  $W$  induced by the six paths  $rvC_wx$ ,

$\{r, u\}, rwC_vy, \{x, y\}, uC_wx$  and  $uC_vy$  does not contain the internal vertices of  $C_u$ , thus it is smaller than  $W$  (third configuration of [Figure 3.13](#)). However, it contains a subdivision of  $K_4$  as subgraph, that can for instance be obtained by contracting each of these six paths to an edge. This contradicts our first remark, therefore the two neighbors  $v$  and  $w$  of  $r$  on the cycle are adjacent. This proves item (ii).

Let now assume that there  $C$  has a second chord  $\{z, t\}$  in  $G$ . In the light of the previous remark,  $C_u$  is an edge, hence the only paths to which  $z$  and  $t$  can belong are the paths  $xC_wv$  and  $yC_vw$  and, according to our first remark, they do not both belong to the same of these two paths. Also, as  $\{z, t\} \neq \{x, y\}$ , one of  $z, t$  does not belong to  $\{x, y\}$ . We can thus assume without loss of generality that  $z \in V(xC_wv)$ ,  $t \in V(yC_vw)$  and  $z \neq x$ . This case is represented by the fourth configuration of [Figure 3.13](#). Let us consider the cycle  $C'$  obtained by the concatenation of the paths  $zC_wv, C_u, wC_vt$  and  $(t, z)$  and the vertex  $r$ , which is connected to the cycle by the three paths  $(r, v), (r, w)$  and  $(r, u) - uC_vt$  that only share the vertex  $r$ . This subgraph is smaller than  $H$  since it does not contain vertex  $x$ , but it is a subdivision of  $K_4$  (three paths issued from the same vertex  $a$  and meeting the cycle  $C'$ ). By one of the above remarks, this configuration is impossible and thus the chord  $(z, t)$  cannot exist. Hence we proved item (iii):  $C$  can have at most one chord in  $G$ .

Notice that we have  $|N_C(r)| \geq 3$  since  $W$  is a 3-wheel. We now assume that  $|N_C(r)| > 3$ . Let  $u, v, w, z$  be four different neighbors of  $r$  such that  $z$  is a common neighbor of the endpoints of the chord  $x$  and  $y$ . This case is depicted in the fifth configuration of [Figure 3.13](#). Then  $r$  has at least three neighbors ( $u, v$ , and  $w$ ) on the cycle going through the edge  $\{x, y\}$  and following  $C$  up to  $x$  without using the vertex  $z$ . As this contradicts the minimality of  $W$ , we have  $|N_C(r)| = 3$ , that is item (iv), and this concludes the proof.  $\square$

**Corollary 3.54.** *According to [Remark 3.52](#), every minimum  $K_4$ -subdivision in  $G$  is a 3-wheel, so [Lemma 3.53](#) is still true when replacing 3-wheel by  $K_4$ -subdivision in its statement.*

**Lemma 3.55.** *Every two non-adjacent vertices of  $R$  have the same neighborhood in  $C$ .*

*Proof.* By contradiction, we assume that there are two non-adjacent vertices  $s, t \in V(R)$  and a vertex  $u_1 \in V(C)$  such that  $\{s, u_1\} \in E(G)$  but  $\{t, u_1\} \notin E(G)$ . By [Remark 3.51](#),  $s$  and  $t$  have (at least) three neighbors in  $C$ . Let  $U = \{u_1, u_2, u_3\}$  and  $V = \{v_1, v_2, v_3\}$  be the respective neighbors of  $s$  and  $t$ . We consider the graph  $H$  induced in  $G$  by  $C$  and  $\{s, t\}$  where we iteratively contracted every edge of  $S$  that is not incident with two vertices of  $\{u_1, u_2, u_3, v_1, v_2, v_3\}$ . This graph consists of a cycle on at most 6 vertices that may have chords; we call this cycle  $C'$ ; plus the two non-adjacent vertices of degree at least three  $s$  and  $t$ . Remark that while two neighbors of  $s$  are adjacent and are not both neighbors of  $t$ , we can contract the edge between them and decrease the degree of  $s$ , without changing degree of  $t$ . If the degree of  $s$  reaches two by such means, then by [Lemma 3.46](#),  $K_4 \leq_{i.m.} H$ , a contradiction. We can thus assume that every vertex of  $C'$  adjacent to a neighbor of  $s$  is a neighbor of  $t$ . This is also true when  $s$  and  $t$  are swapped since this argument can be applied to  $t$  too. This observation implies that  $N_S(s) \cap N_S(t) = \emptyset$  (as  $u_1$  is adjacent to  $s$  but not to  $t$ , none of its neighbors on  $C$  can be adjacent to  $s$ , and so on along the cycle)

and that the neighbors of  $s$  and  $t$  are alternating on  $C'$ . Without loss of generality, we suppose that  $C' = u_1v_1u_2v_2u_3v_3$ . We consider now the five following sets of vertices of  $H$  :  $M_1 = \{u_1\}$ ,  $M_2 = \{s\}$ ,  $M_3 = \{u_2, v_1\}$ ,  $M_4 = \{v_2, u_3, v_3\}$ ,  $M_5 = \{t\}$ . They are depicted on [Figure 3.14](#). Let  $\mu: \hat{K}_4 \rightarrow \mathcal{P}^{<\omega}(V(H))$  be the function defined as follows:  $\forall i \in \llbracket 1, 5 \rrbracket$ ,  $\mu(v_i) = M_i$  (using the names of vertices of  $\hat{K}_4$  defined on [Figure 3.2](#)). Now, remark that  $\mu$  is an induced minor model of  $\hat{K}_4$  in  $H$ : for every  $i \in \llbracket 1, 5 \rrbracket$ , the set  $M_i$  is connected,  $M_1, M_3, M_4$  forms a cycle (using edges  $\{u_1, v_1\}, \{u_2, v_2\}, \{v_3, u_1\}$  is this order),  $M_2$  is adjacent to any of these three sets (by edges  $\{s, u_1\}, \{s, u_2\}, \{s, u_3\}$ ) and the set  $M_5$  is only adjacent to  $M_3$  and  $M_4$  (by edges  $\{t, v_1\}, \{t, v_2\}$ ). Remark that the previous statement holds even when  $C'$  is not an induced cycle, as any possible chord of  $C'$  will be between vertices of the sets  $M_1, M_2, M_3, M_4$  (as  $M_5$  is reduced to  $t \notin V(C')$ ) and these sets are already all pairwise adjacent. We assumed our initial graph to be  $\hat{K}_4$ -induced minor-free but we proved that it contains an induced minor model of  $\hat{K}_4$  : this is the contradiction we were looking for.  $\square$




Figure 3.14: Graph  $H$  (left) used in [Lemma 3.55](#) (middle) and in [Lemma 3.56](#) (right).

**Lemma 3.56.** *Every two adjacent vertices of  $R$  have the same neighborhood in  $C$ .*

*Proof.* By contradiction, we assume that there are two adjacent vertices  $s, t \in V(R)$  and  $u_1 \in V(C)$  such that  $\{s, u_1\} \in E(G)$  but  $\{t, u_1\} \notin E(G)$ . As this proof is very similar to the proof of [Lemma 3.55](#), we define  $u_1, u_2, u_3, v_1, v_2, v_3, U, V, H, C'$  in the same way here.

*First case:*  $C'$  is an induced cycle. In this case, the graph  $H$  is the (induced) cycle  $C' = u_1v_1u_2v_2u_3v_3$  plus the two adjacent vertices  $s$  and  $t$ . Let us define five vertex sets:  $M_1 = \{v_2\}$ ,  $M_2 = \{t\}$ ,  $M_3 = \{u_3, v_3\}$ ,  $M_4 = \{v_1, u_2, s\}$ ,  $M_5 = \{u_1\}$ . They are depicted on [Figure 3.14](#). Now, remark that the function that sends the vertex of  $\hat{K}_4$  labeled  $i$  on [Figure 3.2](#) to  $M_i$  is an induced minor model of  $\hat{K}_4$  in  $H$ : every set  $M_i$  is connected,  $M_1, M_3, M_4$  forms a cycle (using edges  $\{v_2, u_3\}, \{u_3, s\}, \{u_2, v_2\}$  is this order),  $M_2$  is adjacent to any of these three sets (by edges  $\{t, v_2\}, \{t, v_3\}, \{t, v_1\}$ ), and the set  $M_5$  is only adjacent to  $M_3$  and  $M_4$  (by edges  $\{u_1, v_3\}, \{u_1, v_1\}$ ). As we assumed our initial graph to be  $\hat{K}_4$ -induced minor-free, this is a contradiction.

*Second case:* the cycle  $C'$  is not induced. By [Lemma 3.53](#) and as  $C$  is supposed to be a minimal 3-wheel of  $G$ , the cycle  $C$  has only one chord. In this case, the graph  $H$  is the (induced) cycle  $C' = u_1v_1u_2v_2u_3v_3$  plus the two adjacent vertices  $s$  and  $t$  and an edge

$e$  between two vertices of  $C'$ . In  $H$ , both  $H \setminus \{s\}$  and  $H \setminus \{t\}$  are minimal 3-wheels, sharing the same cycle  $C'$ . By applying [Lemma 3.53](#) on these two 3-wheels, we obtain that the endpoints of  $e$  must both be adjacent to a vertex of  $u_1, u_2, u_3$  (neighbor of  $s$  on  $C'$ ) and to a vertex of  $v_1, v_2, v_3$  (neighbor of  $t$  on  $C'$ ). Such a configuration is impossible.  $\square$

**Lemma 3.57.** *If  $C$  is not an induced cycle of  $G$ , then  $|V(R)| = 1$ .*

*Proof.* Let  $r \in V(R)$  be the center of a minimum 3-wheel of cycle  $C$ . By contradiction, let us assume that  $R$  contains a vertex  $s \neq r$ . By [Lemma 3.53](#),  $r$  has exactly three neighbors on  $C$ , one of which, that we call  $u$ , is adjacent to both endpoints of the only chord of  $C$ . Furthermore the two other neighbors of  $r$ , that we denote by  $\{v, w\}$ , are adjacent. According to [Lemma 3.55](#) and [Lemma 3.56](#),  $r$  and  $s$  have the same neighborhood in  $C$ . There are now two different cases to consider depending whether  $\{r, s\}$  is an edge or not. The case  $\{r, s\} \in E(G)$  (respectively  $\{r, s\} \notin E(G)$ ) is depicted on the right (respectively left) of [Figure 3.15](#). Remark that if  $\{r, s\} \in E(G)$ , the graph  $G$  satisfies conditions of [Lemma 3.46](#) (with  $G[\{r, s, u, v, w\}]$  for  $S$ ,  $\{u, v\}$  for  $\{s_1, s_2\}$  and  $x$  for  $x$ ), so  $G \leq_{i.m.} \hat{K}_4$ , what is contradictory. In the other hand, when  $\{r, s\} \notin E(G)$ , the induced subgraph  $G[\{r, s, u, v, w\}]$  is isomorphic to  $\hat{K}_4$ , so we also have  $G \leq_{i.m.} \hat{K}_4$ . Hence  $|V(R)| < 2$ . By definition of  $C$  and  $R$ , the subgraph  $R$  contains at least one vertex (that is the center of a 3-wheel of cycle  $C$ ), thus  $|V(R)| = 1$  as required.




Figure 3.15: Two different cases in the proof of [Lemma 3.57](#)

$\square$

Recall that the prism is the Cartesian product of  $K_3$  and  $K_2$ , see [Figure 3.16](#) for a picture.




Figure 3.16: The prism.

**Corollary 3.58.** *If  $C$  is not an induced cycle of  $G$ , then  $G$  is a subdivision of the prism.*

**Lemma 3.59.**  *$R$  is complete multipartite.*

*Proof.* As a graph is complete multipartite iff it does not contain  $K_1 + K_2$  as induced subgraph, we only need to show that the case where  $R = K_1 + K_2$  is not possible. Consequently, let us assume that we are in this case, and let  $u, v, w$  be the vertices of  $R$ ,  $\{u, v\}$  being the only edge in  $R$ . As  $R$  is an induced subgraph of  $G$ ,  $u$  and  $w$  are not adjacent in  $G$  neither. By [Lemma 3.55](#), they have the same neighborhood on  $C$ . The same argument can be applied to  $v$  and  $w$  to show that  $N_C(u) = N_C(v) = N_C(w)$ . Let  $x, y, z \in N_C(u)$  be three different vertices (they exist by [Corollary 3.47](#)) and let  $H$  be the graph obtained from  $G$  by contracting every edge of  $C$  that is not incident with two vertices of  $\{x, y, z\}$ . Such a graph is a triangle (obtained by contracting  $C$ ) and the three vertices  $u, v, w$  each adjacent to every vertex of the triangle, as drawn in [Figure 3.17](#). Deleting vertex  $y$  gives a graph isomorphic to  $\hat{K}_4$ , as one can easily check (cf. [Lemma 3.59](#)).  $\square$




Figure 3.17: The graph  $H$  of [Lemma 3.59](#) (left) and the graph obtained after deletion of  $y$  (right).

We now need to show that every graph containing a  $K_4$ -subdivision either has a proper  $K_4$ -subdivision, or fall in the possible cases of the statement of [Theorem 3.39](#).

**Lemma 3.60.** *If  $G$  can be obtained by adding an edge between two vertices of a  $K_{3,3}$ -subdivision, then  $G$  has a proper  $K_4$ -subdivision.*

*Proof.* Let  $S$  be the spanning subgraph of  $G$  which is a  $K_{3,3}$ -subdivision. A *branch* of  $S$  is a maximal path, the internal vertices of which are of degree two. In  $S$ , non-subdivision vertices are connected by branches. Let us call  $a, b, c, x, y, z$  the non-subdivision vertices of  $S$  in a way such that there is neither a branch between any two vertices of  $\{a, b, c\}$  nor between any two vertices of  $\{x, y, z\}$  (intuitively  $\{a, b, c\}$  and  $\{x, y, z\}$  correspond to the two maximum independent sets of  $K_{3,3}$ ). Observe that every  $K_{3,3}$ -subdivision contains a  $K_4$ -subdivision (but not a proper one). Let us now consider all the possible endpoints of the only edge  $e$  of  $E(G) \setminus E(S)$ .

*First case:* both endpoints of  $e$  belong to the same branch  $B$  of  $S$ . Let  $X$  be the set of internal vertices of the subpath of  $B$  starting at the one endpoint of  $e$  and ending at the other one. As  $G$  is a simple graph,  $|X| \geq 1$ . Then  $G \setminus X$  has a  $K_4$ -subdivision (as it is a  $K_{3,3}$ -subdivision), which is a proper  $K_4$ -subdivision of  $G$ .

*Second case:*  $e$  is incident with two non-subdivision vertices. Observe that the case where  $e$  is incident with a vertex from  $\{a, b, c\}$  and the other from  $\{x, y, z\}$  is contained in the previous case. Let us assume without loss of generality that  $e = \{a, b\}$ . Then  $G \setminus \{x\}$  has a  $K_4$ -subdivision. Indeed, if  $B_{s,t}$  denotes the branch with endpoints the vertices  $s$  and  $t$  (for  $(s, t) \in \{a, b, c\} \times \{x, y, z\}$ ), then the vertices of the paths  $B_{b,y}, B_{b,z}$ ,

$B_{c,z}$  and  $B_{c,y}$  induce a cycle in  $G$ . The vertex  $a$  is then connected to this cycle with the paths  $B_{a,y}$ ,  $B_{a,z}$  and the edge  $e$ . Hence  $G$  has a proper  $K_4$ -subdivision, as required.

*Third case:*  $e$  is incident with two subdivision vertices. If the two endpoints of  $e$  belong to the same branch, then we are in the first case. Otherwise, we can easily reach the second case as follows. If we contract all the edges on the path connecting the first endpoint of  $e$  to a vertex of  $\{a, b, c\}$  and all the edges on the path connecting the second endpoint of  $e$  to a vertex of  $\{x, y, z\}$ , we get a  $K_4$ -subdivision (because we never contracted an edge incident with two non-subdivision vertices of  $S$ ) plus the edge  $e$  which is now incident with two non-subdivision vertices. This concludes the proof.  $\square$

**Lemma 3.61.** *If  $G$  can be obtained by adding an edge between two vertices of a prism-subdivision, then  $G$  has a proper  $K_4$ -subdivision.*

*Proof.* Let  $S$  be a prism-subdivision in  $G$  and let  $e \in E(G) \setminus E(S)$ . We will use the concept of *branch* defined in the proof of Lemma 3.60, which is very similar to this one. Let us call  $a, b, c, x, y, z$  the non-subdivision vertices in a way such that there are branches between every pair of vertices of  $\{a, b, c\}$  (respectively  $\{x, y, z\}$ ) and between vertices of the pairs  $(a, x)$ ,  $(b, y)$ , and  $(c, z)$ . Intuitively  $\{a, b, c\}$  and  $\{x, y, z\}$  correspond to the two triangles of the prism. Let us consider the positions of the endpoints of  $e$ .

*First case:* both endpoints of  $e$  belong to the same branch of  $S$ . Since the prism contains a  $K_4$  subdivision (but not a proper one), we can in this case find a smaller prism subdivision as in the first case of the proof of Lemma 3.60, and thus a proper  $K_4$ -subdivision.

*Second case:*  $e$  is incident with two non-subdivision vertices. Let us assume without loss of generality that  $e = \{a, y\}$  (the cases  $e \subseteq \{a, b, c\}$ ,  $e \subseteq \{x, y, z\}$ , and  $e \in \{\{a, x\}, \{b, y\}, \{c, z\}\}$  are subcases of the first one). Then in  $G \setminus \{x\}$ , the paths  $B_{a,b}$ ,  $B_{b,z}$  and  $B_{x,y}$  together with the edge  $e$  induces a cycle to which the vertex  $c$  is connected via the paths  $B_{c,b}$ ,  $B_{c,a}$ , and  $B_{c,y}$ . Hence  $G$  has a proper  $K_4$ -subdivision.

*Third case:*  $e$  is incident with two branches between  $a, b$ , and  $c$  (and the symmetric case with branches between  $x, y$ , and  $z$ ). Let us assume without loss of generality that  $e$  has one endpoint  $r$  among the interior vertices of  $B_{a,c}$  and the other one  $s$  among the interior vertices of  $B_{b,c}$ . Then  $(r, s)$ ,  $sB_{b,c}c$ , and  $cB_{a,c}r$  induce in  $G \setminus \{x\}$  a cycle to which the vertex  $b$  is connected via the paths  $bB_{b,c}s$ ,  $bB_{a,b}a$  together with  $aB_{a,c}r$ , and  $B_{b,z}$  together with  $B_{z,y}$  and  $B_{y,b}$ . Again,  $G$  has a proper  $K_4$ -subdivision.

*Fourth case:*  $e$  is incident with two branches, the one connected to a vertex in  $\{a, b, c\}$  and the other one connected to a vertex in  $\{x, y, z\}$ . In this case, by contracting the edges of the first branch that are between the endpoint of  $e$  and a vertex of  $\{a, b, c\}$  and similarly with the second branch and a vertex of  $\{x, y, z\}$  gives a graph with a prism-subdivision plus an edge between two non-subdivision vertices (that is, first case), exactly as in the proof of Lemma 3.60.  $\square$

**Lemma 3.62.** *If graph  $G$  contains a  $K_4$ -subdivision, then either  $G$  has a proper  $K_4$ -subdivision, or  $G$  is a wheel, or a subdivision of one of the following graphs:  $K_4$ ,  $K_{3,3}$ , and the prism.*

*Proof.* Looking for a contradiction, let  $G$  be a counterexample with the minimum number of vertices and, subject to that, the minimum number of edges. Let  $S$  be a  $K_4$ -subdivision in  $G$  and let  $e \in E(G) \setminus E(S)$ . Observe that since  $G$  has no proper  $K_4$ -subdivision,  $S$  is a spanning subgraph of  $G$ . Also,  $e$  is well defined as we assume that  $G$  is not a  $K_4$ -subdivision. Notice that since the minimum degree of  $K_4$  is 3, contracting an edge incident with a vertex of degree 2 in  $G$  would yield a smaller counterexample. Therefore  $G$  has minimum degree at least 3. Let  $G' = G \setminus \{e\}$ . This graph clearly contains  $S$ . By minimality of  $G$ , the graph  $G'$  is either a wheel, or a subdivision of a graph among  $K_4$ ,  $K_{3,3}$ , and the prism. Observe that  $G'$  cannot have a proper  $K_4$ -subdivision because it would also be a proper  $K_4$ -subdivision in  $G$ .

*First case:*  $G'$  is a wheel. Let  $C$  be the cycle of the wheel and let  $r$  be its center. Obviously, in  $G$  the edge  $e$  has not  $r$  as endpoint otherwise  $G$  would also be a wheel. Therefore  $e$  is incident with two vertices of  $C$ . Let  $P$  and  $P'$  be the two subpaths of  $C$  whose endpoints are then endpoints of  $e$ . Observe that none of  $P$  and  $P'$  contains more than two neighbors of  $r$ . Indeed, if, say,  $P$  contained at least three neighbors of  $r$ , then the subgraph of  $G$  induced by the vertices of  $P$ ,  $e$ , and  $r$  would contain a  $K_4$ -subdivision, hence contradicting the fact that  $G$  has no proper  $K_4$ -subdivision.

Therefore  $G$  is the cycle  $C$  with exactly one chord,  $e$ , and the vertex  $r$  which has at most 4 neighbors on  $C$ . Because  $G$  has maximum degree at least 3, it has at most 7 vertices. If  $r$  has three neighbors on  $C$ , then necessarily  $P$  contains one of them and  $P'$  the other two (or the other way around). We can easily check in this case that  $G$  is a subdivision of the prism. If  $r$  has four neighbors on  $C$ , the interior of  $P$  and  $P'$  must each contain two of them according to the above remarks. The deletion of any neighbor of  $r$  in this graph yields a  $K_4$ -subdivision of non-subdivision vertices  $r$  and the remaining neighbors. Observe that both cases contradict the assumptions made on  $G$ .

*Second case:*  $G'$  is a subdivision of  $K_4$ , or  $K_{3,3}$ , or the prism. If  $G'$  is a subdivision of  $K_{3,3}$  or of the prism, then the result follows by [Lemma 3.60](#) and [Lemma 3.61](#). Let us now assume that  $G'$  is a subdivision of  $K_4$  and let us consider branches of this subdivision as defined in the proof of [Lemma 3.60](#). If  $e$  has endpoints in the same branch, then as in the first cases of the aforementioned lemmas we can find in  $G'$  a  $K_4$ -subdivision with fewer vertices and thus a proper  $K_4$ -subdivision in  $G$ . In the case where the endpoints of  $e$  belong to the interior of two different branches, then it is easy to see that  $G$  is a prism-subdivision. Let  $\{x, y, z, t\}$  be the non-subdivision vertices of the  $K_4$ -subdivision. Finally, let us assume that the one endpoint of  $e$  is a non-subdivision vertex, say  $x$ , and the other one, that we call  $u$ , is a subdivision vertex of a branch, say  $B_{y,z}$  (using the same notation as in the proof of [Lemma 3.60](#)). If  $X$  is set set of interior vertices of one of  $B_{x,y}$ ,  $B_{x,z}$ , or  $B_{x,t}$ , then the graph  $G \setminus X$  has a  $K_4$ -subdivision of non-subdivision vertices  $x, u, z, t$ ,  $x, y, y, t$  or  $x, y, u, z$  respectively. In this case  $G$  has a proper  $K_4$ -subdivision. If none of  $B_{x,y}$ ,  $B_{x,z}$ , and  $B_{x,t}$  has internal vertices, then  $G$  is a wheel of center  $x$ .

In all the possible cases we reached the contradiction we were looking for. This concludes the proof.  $\square$

We are now ready to prove [Theorem 3.39](#), that we restate here.

**Theorem 3.39** (Decomposition of  $\hat{K}_4$ -induced minor-free graphs). *Let  $G$  be a 2-connected graph such that  $\hat{K}_4 \not\leq_{i.m.} G$ . Then:*

- *either  $K_4 \not\leq_{i.m.} G$ ;*
- *or  $G$  is a subdivision of a graph among  $K_4$ ,  $K_{3,3}$ , and the prism;*
- *or  $V(G)$  has a partition  $(C, M)$  such that  $G[C]$  is an induced cycle,  $G[M]$  is a complete multipartite graph and every vertex of  $C$  is either adjacent in  $G$  to all vertices of  $M$ , or to none of them.*

*Proof.* Let  $G \in \text{Excl}_{\leq i.m.}(\hat{K}_4)$  be a 2-connected graph. If  $G$  does not contain a  $K_4$ -subdivision, then the theorem is trivially true for  $G$ . If graph  $G$  contains a  $K_4$ -subdivision but not a proper one, from [Lemma 3.62](#) we get that  $G$  is a subdivision of one of  $K_4$ ,  $K_{3,3}$ , or the prism, in which case the theorem holds, or that  $G$  is a wheel, which has a trivial cycle-multipartite decomposition, with the center being the multipartite part.

Finally, let us assume that  $G$  contains a proper  $K_4$ -subdivision. By [Lemma 3.50](#),  $G$  contains a 3-wheel. Let  $C$  be the cycle of a minimum 3-wheel in  $G$  and  $R$  the subgraph of  $G$  induced by  $V(G) \setminus V(C)$ . According to [Corollary 3.58](#), if  $C$  is not an induced cycle, then  $G$  is a subdivision of the prism. When  $C$  is induced, then by [Lemma 3.59](#),  $R$  is complete multipartite. Furthermore, vertices of  $R$  have the same neighborhood on  $C$ , as proved in [Lemma 3.55](#) and [Lemma 3.56](#). Therefore,  $(C, R)$  is a cycle-multipartite decomposition of  $G$  and we are done.  $\square$

### From a decomposition theorem to well-quasi-ordering

This section is devoted to the proof of [Theorem 3.41](#).

The proof relies on the two following lemmas which are proved in the next subsections.

**Lemma 3.63.** *For every (unlabeled) graph  $G$  and every wqo  $(S, \preceq)$ , the class of  $(S, \preceq)$ -labeled  $G$ -subdivisions is well-quasi-ordered by the contraction relation.*

**Lemma 3.64.** *For every wqo  $(S, \preceq)$ , the class of  $(S, \preceq)$ -labeled graphs having a cycle-multipartite decomposition is well-quasi-ordered by induced minors.*

Let us recall the statement of [Theorem 3.41](#).

**Theorem 3.41.** *The class of  $\hat{K}_4$ -induced minor-free graphs is wqo by  $\leq_{i.m.}$ .*

*Proof.* The class of graphs not containing  $K_4$  as minor (or, equivalently, as induced minor) has been shown to be well-quasi-ordered by induced minors in [[Tho85](#)], cf. [Theorem 3.11](#). According to [Remark 3.25](#), we can then restrict our attention to graphs of  $\text{Excl}_{\leq i.m.}(\hat{K}_4)$  that contain  $K_4$  as minor. As some of these graphs might not be 2-connected, we use [Lemma 3.31](#): it is enough to show that for every wqo  $(S, \preceq)$ , the class of  $(S, \preceq)$ -labeled 2-connected graphs containing  $K_4$  as minor are wqo by induced minors. By [Theorem 3.39](#), this class can be divided into two subclasses:

- (2-connected) subdivisions of a graph among  $K_4$ ,  $K_{3,3}$ , and the prism;
- graphs having a cycle-multipartite decomposition.

**Lemma 3.63** and **Lemma 3.64** handle these two cases, hence by **Remark 3.25** the class of  $(S, \preceq)$ -labeled 2-connected graphs containing  $K_4$  as minor are wqo by induced minors for every wqo  $(S, \preceq)$ . This concludes the proof.  $\square$

The following subsection contains the proofs of **Lemma 3.63** and **Lemma 3.64**, that follow the steps described in **Section 3.4**.

### Well-quasi-ordering subdivisions

Let  $\mathcal{OP}$  denote the class of paths whose endpoints are distinguished, i.e. one end is said to be the beginning and the other one the end. In the sequel,  $\text{fst}(P)$  denotes the first vertex of the path  $P$  and  $\text{lst}(P)$  its last vertex. We extend the relation  $\leq_{\text{i.m.}}$  to  $\mathcal{OP}$  as follows: for every  $G, H \in \mathcal{OP}$ ,  $G \leq_{\text{i.m.}} H$  if there is an induced minor model  $\mu$  of  $G$  in  $H$  such that  $\text{fst}(H) \in \mu(\text{fst}(G))$  and  $\text{lst}(H) \in \mu(\text{lst}(G))$ , and similarly for  $\leq_{\text{c.}}$ .

**Lemma 3.65.** *If the qoset  $(Q, \preceq_Q)$  is a wqo, then the qoset  $(\text{lab}_{(Q, \preceq_Q)}(\mathcal{OP}), \leq_{\text{c.}})$  also is a wqo.*

*Proof.* Let us take  $(S, \preceq)$  to be  $(\mathcal{P}^{<\omega}(Q), \preceq_Q^{\mathcal{P}})$ . By **Corollary 3.27**  $(S, \preceq)$  is a wqo. We consider the function

$$f: (S^*, \preceq^*) \rightarrow (\text{lab}_{(S, \preceq)}(\mathcal{OP}), \leq_{\text{c.}})$$

that, given a sequence  $\langle s_1, \dots, s_k \rangle \in S^*$  of elements of  $S$ , returns the path  $P$  on  $k$  vertices whose  $i$ -th vertex  $v_i$  is labeled by  $s_i$  for every  $i \in \llbracket 1, k \rrbracket$  and where  $\text{fst}(P) = v_1$  and  $\text{lst}(P) = v_k$ . The image of this function is clearly  $\text{lab}_{(S, \preceq)}(\mathcal{OP})$  and by **Theorem 3.1** its domain is well-quasi-ordered by  $\preceq^*$ . By the virtue of **Remark 3.28**, it is thus enough to show that

$$f: (S^*, \preceq^*) \rightarrow (\text{lab}_{(S, \preceq)}(\mathcal{OP}), \leq_{\text{c.}})$$

is monotone in order to prove that  $(\text{lab}_{(S, \preceq)}(\mathcal{OP}), \leq_{\text{c.}})$  is wqo.

Let  $R = \langle r_1, \dots, r_k \rangle, S = \langle s_1, \dots, s_l \rangle \in S^*$ , be two sequences such that  $R \preceq^* S$  and let us show that  $f(R) \leq_{\text{c.}} f(S)$ . We will use the following notation:  $f(R)$  is the path  $v_1 \dots v_k$  labeled by  $\lambda_R$  and similarly for  $f(S)$ ,  $u_1 \dots u_l$  and  $\lambda_S$ . Let  $\varphi: \llbracket 1, k \rrbracket \rightarrow \llbracket 1, l \rrbracket$  be an increasing function such that  $\forall i \in \llbracket 1, k \rrbracket$ , we have  $r_i \preceq_{s_{\varphi(i)}}$  (such a function exists since  $R \preceq^* S$ ). Let us consider the path obtained from  $f(S)$  by, for every  $i \notin \{\varphi(j)\}_{j \in \llbracket 1, k \rrbracket}$ , contracting the label<sup>4</sup> of  $u_i$  to the empty set and then dissolving  $u_i$ . Remark that this graph is a path on  $k$  vertices  $p_1 p_2 \dots p_k$  such that  $\forall i \in \llbracket 1, k \rrbracket$ ,  $\lambda_R(v_i) = r_i \preceq \lambda_S(p_i) = s_{\varphi(i)}$ . Furthermore, this path is a contraction of  $f(S)$  where either  $u_1 = p_1$  (respectively  $u_l = p_k$ ) or this vertex has been contracted to  $p_1$  (respectively  $p_k$ ), hence  $f(R) \leq_{\text{c.}} f(S)$ , as desired.  $\square$

We can now prove **Lemma 3.63**, that we first restate.

<sup>4</sup>Recall that a label contraction is the operation of relabeling with a label that is smaller (in the qoset used for labelling).

**Lemma 3.63.** *For every (unlabeled) graph  $G$  and every wqo  $(S, \preceq)$ , the class of  $(S, \preceq)$ -labeled  $G$ -subdivisions is well-quasi-ordered by the contraction relation.*

*Proof.* Let  $G$  be a non-labeled graph, let  $(S, \preceq)$  be a wqo and let  $\mathcal{G}$  be the class of all  $(S, \preceq)$ -labeled  $G$ -subdivisions. We set  $m = |E(G)|$ . Let us show that  $(\mathcal{G}, \leq_c)$  is a wqo. First, we arbitrarily choose an orientation to every edge of  $G$  and an enumeration  $e_1, \dots, e_m$  of these edges. We now consider the function  $f$  that, given a tuple  $(Q_1, \dots, Q_m)$  of  $m$  paths of  $\text{lab}_{(S, \preceq)}(\mathcal{OP})$ , returns the graph constructed from  $G$  by, for every  $i \in \llbracket 1, m \rrbracket$ , replacing the edge  $e_i$  by the path  $Q_i$ , while respecting the orientation, i.e. the first (respectively last) vertex of  $Q_i$  goes to the first (respectively last) vertex of  $e_i$ . By [Lemma 3.26](#) on Cartesian products of wqos and since  $(\text{lab}_{(S, \preceq)}(\mathcal{OP}), \leq_c)$  is a wqo ([Lemma 3.65](#)), the domain  $\text{lab}_{(S, \preceq)}(\mathcal{OP})^m$  of  $f$  is well-quasi-ordered by  $\leq_c^m$ . Notice that every element of the codomain of  $f$  is an  $G$ -subdivision (by definitions of  $f$ ), and moreover that  $f$  is surjective on  $\mathcal{G}$ : for every  $(S, \preceq)$ -labeled  $G$ -subdivision  $H$  we can construct a tuple  $(Q_1, \dots, Q_m)$  of  $m$  paths of  $\text{lab}_{(S, \preceq)}(\mathcal{OP})$ , such that  $f(Q_1, \dots, Q_m) = H$ .

In order to show that  $(\mathcal{G}, \leq_c)$  is a wqo, it is enough to prove that

$$f: (\text{lab}_{(S, \preceq)}(\mathcal{OP}), \leq_c^m) \rightarrow (\mathcal{G}, \leq_c)$$

is an epi, as explained in [Remark 3.28](#), that is, to prove that for every two tuples  $\mathcal{Q}, \mathcal{R} \in \text{lab}_{(S, \preceq)}(\mathcal{OP})^m$  such that  $\mathcal{Q} \leq_c^m \mathcal{R}$ , we have  $f(\mathcal{Q}) \leq_c f(\mathcal{R})$ . According to [Remark 3.29](#), we only need to care, for every  $i \in \llbracket 1, m \rrbracket$ , of the case where  $\mathcal{Q}$  and  $\mathcal{R}$  only differs by the  $i$ -th coordinate. It is at this point of the proof important to remark the symmetry of the definition of  $f$ : since the different coordinates any element of the domain of  $f$  are playing the same role, we only have to deal with the case where  $\mathcal{Q}$  and  $\mathcal{R}$  differs by one (fixed) coordinate, say the first one. Therefore, let us consider two tuples  $\mathcal{Q} = (Q, Q_2, \dots, Q_m)$  and  $\mathcal{R} = (R, Q_2, \dots, Q_m)$  of  $\text{lab}_{(S, \preceq)}(\mathcal{OP})^m$  such that  $\mathcal{Q} \leq_c^m \mathcal{R}$ , i.e. satisfying  $Q \leq_c R$ . Let  $\mu: V(Q) \rightarrow \mathcal{P}^{<\omega}(V(R))$  be a contraction model of  $Q$  in  $R$  and let  $\mu'$  be the trivial contraction model of  $f(\mathcal{Q})$  in itself, that is,

$$\mu': \begin{cases} V(f(\mathcal{Q})) & \rightarrow \mathcal{P}^{<\omega}(V(f(\mathcal{Q}))) \\ u & \mapsto \{u\}. \end{cases}$$

We now consider the function  $\nu$  defined as follows:

$$\nu: \begin{cases} V(f(\mathcal{Q})) & \rightarrow \mathcal{P}^{<\omega}(V(f(\mathcal{R}))) \\ u & \mapsto \mu(u) \quad \text{if } u \in V(Q) \\ u & \mapsto \mu'(u) \quad \text{otherwise.} \end{cases}$$

Let us show that  $\nu$  is a contraction model of  $f(\mathcal{Q})$  in  $f(\mathcal{R})$ . First, notice that since both  $\mu$  and  $\mu'$  are contraction models,  $\nu$  inherits some of their properties: for every  $u \in V(f(\mathcal{Q}))$ , the subgraph induced in  $f(\mathcal{R})$  by  $\nu(u)$  is connected and  $\lambda_{f(\mathcal{Q})}(u) \subseteq \bigcup_{v \in \mu(u)} \lambda_{f(\mathcal{R})}(v)$ . For the same reason, we have:

$$\begin{aligned} \bigcup_{u \in V(f(\mathcal{Q}))} \nu(u) &= \bigcup_{u \in V(Q)} \mu(u) \cup \bigcup_{u \in V(f(\mathcal{Q}) \setminus V(Q))} \mu'(u) \\ &= V(R) \cup V(f(\mathcal{Q})) \setminus V(Q) \\ &= V(f(\mathcal{R})). \end{aligned}$$

Let us now consider two distinct vertices  $u$  and  $v$  of  $f(Q)$ .

*First case:*  $u$  and  $v$  both belong to the same set among  $V(Q)$  and  $V(f(Q)) \setminus V(Q)$ . In this case  $\nu(u)$  and  $\nu(v)$  are disjoint and they are adjacent iff  $\{u, v\} \in E(f(Q))$  since both  $\mu$  and  $\mu'$  are contraction models.

*Second case:*  $u \in V(Q)$  and  $v \in V(f(Q)) \setminus V(Q)$  (or the symmetric case). As in the previous case,  $\nu(u)$  and  $\nu(v)$  are disjoint. Assume that  $\{u, v\}$  is an edge of  $f(Q)$ . Notice that we necessarily have either  $u = \text{fst}(Q)$  and  $v \in N_{f(Q) \setminus V(Q)}(\text{fst}(Q))$ , or  $u = \text{lst}(Q)$  and  $v \in N_{f(Q) \setminus V(Q)}(\text{lst}(Q))$ . Let us assume, without loss of generality, that we are in the first of these two subcases. By definition of  $f(\mathcal{R})$ ,  $\{\text{fst}(R), v\}$  is an edge. Since  $\mu$  is a contraction model, we then also have  $\text{fst}(R) \in \nu(u)$  and therefore  $\nu(v)$  and  $\nu(u)$  are adjacent in  $f(\mathcal{R})$ .

We just proved that  $\nu$  is an induced minor model of  $f(Q)$  in  $f(\mathcal{R})$ . As explained above, this is enough in order to show that  $f$  is monotone. Hence  $(\mathcal{G}, \leq_c)$  is a wqo and this concludes the proof.  $\square$

### Well-quasi-ordering cycle-multipartite decompositions

In this section, we show that graphs having a cycle-multipartite decomposition are well-quasi-ordered by induced minors.

**Lemma 3.66.** *If  $(Q, \preceq_Q)$  is wqo then the class of  $(Q, \preceq_Q)$ -labeled independent sets is wqo by the induced subgraph relation.*

*Proof.* We will again define  $(S, \preceq) := (\mathcal{P}^{<\omega}(Q), \preceq_Q^{\mathcal{P}})$ , and observe that it is a wqo.

The function  $f$  that maps every sequence  $\langle x_1, \dots, x_k \rangle$  (for some positive integer  $k$ ) of elements of  $S$  to the  $(S, \preceq)$ -labeled independent set on vertex set  $\{v_1, \dots, v_k\}$  where  $v_i$  have label  $x_i$  for every  $i \in \llbracket 1, k \rrbracket$  has clearly the class of  $(S, \preceq)$ -labeled independent sets as codomain. Let us show that  $f$  is an epi. Let  $X = \langle x_1, \dots, x_k \rangle, Y = \langle x_1, \dots, x_l \rangle \in S^*$  be two sequences such that  $X \preceq^* Y$ . By definition of the relation  $\preceq^*$ , there is an increasing function  $\varphi: \llbracket 1, k \rrbracket \rightarrow \llbracket 1, l \rrbracket$  such that  $\forall i \in \llbracket 1, k \rrbracket, x_i \preceq_{\varphi(i)}$ . Therefore the function  $\mu: V(f(X)) \rightarrow V(f(Y))$  that maps the vertex  $v_i$  of  $f(X)$  to the singleton  $\{v_{\varphi(i)}\}$  of  $f(Y)$  is an induced subgraph model of  $f(X)$  in  $f(Y)$  and this proves the monotonicity of  $f$ . By the virtue of [Remark 3.28](#) and since  $(S^*, \preceq^*)$  is a wqo, we get that the class of  $(S, \preceq)$ -labeled independent sets is wqo by the induced subgraph relation.  $\square$

**Corollary 3.67.** *With a very similar proof, we can also show that if  $(S, \preceq)$  is wqo then the class of  $(S, \preceq)$ -labeled cliques is wqo by the induced subgraph relation.*

The *join* of two graphs  $H$  and  $G$  is the graph obtained by taking the disjoint union of  $H$  and  $G$  and adding all possible edges (with multiplicity one, as we deal here with simple graphs) with the one endpoint in  $G$  and the other one in  $H$ .

**Corollary 3.68.** *If a class of  $(S, \preceq)$ -labeled graphs  $(\mathcal{G}, \leq_{\text{i.m.}})$  is wqo, then so is its closure by finite disjoint union (respectively join).*

*Proof.* Let  $\mathcal{U}$  be the closure of  $(\mathcal{G}, \leq_{i.m.})$  by disjoint union. Remark that every graph of  $\mathcal{U}$  can be partitioned in a family of pairwise non-adjacent graphs of  $\mathcal{G}$ . Therefore we can define a function mapping every  $\mathcal{G}$ -labeled independent set to the graph of  $\mathcal{U}$  obtained from  $G$  by replacing each vertex by its label (which is an  $(S, \preceq)$ -labeled graph). It is easy to check that this function is an epi of  $(\mathcal{G}, \leq_{i.m.}) \rightarrow (\mathcal{U}, \leq_{i.m.})$ . Together with [Remark 3.28](#) and [Lemma 3.66](#), this yields the desired result.  $\square$

**Corollary 3.69.** *If  $(S, \preceq)$  is a wqo then the class of  $(S, \preceq)$ -labeled complete multipartite graphs are wqo by the induced subgraph relation.*

We now prove [Lemma 3.64](#). Let us restate it.

**Lemma 3.64.** *For every wqo  $(S, \preceq)$ , the class of  $(S, \preceq)$ -labeled graphs having a cycle-multipartite decomposition is well-quasi-ordered by induced minors.*

*Proof.* We consider the function

$$f: (\text{lab}_{(S, \preceq)}(\mathcal{OP})^* \times \text{lab}_{(S, \preceq)}(\mathcal{K}_{\mathbb{N}^*}), \leq_{c.^*} \times \leq_{i.sg.}) \rightarrow (\text{lab}_{(S, \preceq)}(\mathcal{W}), \leq_{i.m.})$$

that, given a sequence  $[R_0, \dots, R_{k-1}] \in \text{lab}_{(S, \preceq)}(\mathcal{OP})$  of  $(S, \preceq)$ -labeled paths of  $\mathcal{OP}$  and a  $(S, \preceq)$ -labeled complete multipartite graph  $K$ , returns the graph constructed as follows.

1. consider the disjoint union of  $K$  and the paths of  $\{R_i\}_{i \in \llbracket 0, k-1 \rrbracket}$  and call  $v_i$  the vertex obtained by identifying the two vertices  $\text{lst}(R_i)$  and  $\text{fst}(R_{(i+1) \bmod k})$ , for every  $i \in \llbracket 0, k-1 \rrbracket$  (informally, this graph is the disjoint union of  $K$  and the cycle built by putting  $R_i$ 's end-to-end);
2. for every element  $v$  of  $\{v_i\}_{i \in \llbracket 0, k-1 \rrbracket}$ , add all possible edges between  $v$  and the vertices of  $K$ .

Remark that the codomain of  $f$  is  $\mathcal{W}$ . Indeed, every element of the image of  $f$  has a cycle-multipartite decomposition (by construction) and conversely, if  $G \in \mathcal{W}$  is of cycle-multipartite decomposition  $(C, K)$ , one can construct a sequence of  $R_0, \dots, R_k$  of subpaths of  $C$  meeting only on endpoints and whose interior vertices are of degree two such that  $G = f(R_0, \dots, R_{k-1}, K)$ . Let us show that the domain of  $f$  is well-quasi-ordered by  $\leq_{c.^*} \times \leq_{i.sg.}$ . We proved in [Lemma 3.65](#) that  $(\text{lab}_{(S, \preceq)}(\mathcal{OP}), \leq_{c.^*})$  is a wqo and [Corollary 3.69](#) shows that  $(\mathcal{K}_{\mathbb{N}^*}, \leq_{i.sg.})$  is a wqo, so by applying [Theorem 3.1](#) we get first that  $(\text{lab}_{(S, \preceq)}(\mathcal{OP})^*, \leq_{c.^*})$  is a wqo, and then by [Lemma 3.26](#) together with [Corollary 3.69](#) that  $(\text{lab}_{(S, \preceq)}(\mathcal{OP})^* \times \mathcal{K}_{\mathbb{N}^*}, \leq_{c.^*} \times \leq_{i.sg.})$  is a wqo.

According to [Remark 3.28](#), it is enough to show that the function

$$f: (\text{lab}_{(S, \preceq)}(\mathcal{OP})^* \times \text{lab}_{(S, \preceq)}(\mathcal{K}_{\mathbb{N}^*}), \leq_{c.^*} \times \leq_{i.sg.}) \rightarrow (\mathcal{W}, \leq_{i.m.})$$

is an epi in order to prove that  $(\text{lab}_{(S, \preceq)}(\mathcal{W}), \leq_{i.sg.})$  is wqo. We show the monotonicity of  $f$  in two steps: the first by showing

$$\forall R \in \text{lab}_{(S, \preceq)}(\mathcal{OP})^*, \forall H, H' \in \text{lab}_{(S, \preceq)}(\mathcal{K}_{\mathbb{N}^*}), H \leq_{i.sg.} H' \Rightarrow f(R, H) \leq_{i.m.} f(R, H')$$

and the second by proving

$$\forall Q, R \in \text{lab}_{(S, \preceq)}(\mathcal{OP})^*, \forall H \in \text{lab}_{(S, \preceq)}(\mathcal{K}_{\mathbb{N}^*}), Q \leq_{c.}^* R \Rightarrow f(Q, H) \leq_{i.m.} f(R, H).$$

According to **Remark 3.29**, the desired result follows from these two assertions.

*First step.* Let  $R = \langle R_0, \dots, R_{k-1} \rangle \in \text{lab}_{(S, \preceq)}(\mathcal{OP})^*$  and  $H, H' \in \text{lab}_{(S, \preceq)}(\mathcal{K}_{\mathbb{N}^*})$  be such that  $H \leq_{i.sg.} H'$ . We therefore have  $(R, H) \leq_{c.}^* \times \leq_{i.sg.} (R, H')$ . Let us show that  $f(R, H) \leq_{i.m.} f(Q, H')$ . Since  $H \leq_{i.sg.} H'$ , there is a subset  $A \subseteq V(H')$  such that  $H' \setminus A = H$ . Let  $C$  denote the cycle obtained from the disjoint union of  $\{R_i\}_{i \in \llbracket 0, k-1 \rrbracket}$  by the identification of the two vertices  $\text{lst}(R_i)$  and  $\text{fst}(R_{(i+1) \bmod k})$ , for every  $i \in \llbracket 0, k-1 \rrbracket$ , where we call  $v_i$  the vertex resulting from this identification. Let us consider the graph  $f(Q, H) \setminus A$ : to construct this graph we started with the disjoint union of  $C$  and  $H$ , then added all possible edges between  $v_i$  and  $V(H)$  for every  $i \in \llbracket 0, k-1 \rrbracket$ , and at last deleted the vertices of  $A \setminus H$ . Remark that this graph is isomorphic to  $f(R, H)$  and therefore  $f(R, H) \leq_{i.m.} f(R, H')$ , as desired.

*Second step.* Let  $Q = \langle Q_0, \dots, Q_{k-1} \rangle$  and  $R = \langle R_0, \dots, R_{l-1} \rangle$  be two elements of  $\text{lab}_{(S, \preceq)}(\mathcal{OP})^*$  such that  $Q \leq_{c.}^* R$  and let  $H \in \text{lab}_{(S, \preceq)}(\mathcal{K}_{\mathbb{N}^*})$ . We thus have  $(Q, H) \leq_{c.}^* \times \leq_{i.sg.} (R, H)$ . Let us show that  $f(Q, H) \leq_{i.m.} f(R, H)$ . By definition of the relation  $\leq_{c.}^*$ , there is an increasing function  $\varphi: \llbracket 0, k-1 \rrbracket \rightarrow \llbracket 0, l-1 \rrbracket$  such that

$$\forall i \in \llbracket 0, k-1 \rrbracket, Q_i \leq_{c.} R_{\varphi(i)}.$$

For every  $i \in \llbracket 0, k-1 \rrbracket$ , let  $\mu_i: V(Q_i) \rightarrow \mathcal{P}^{<\omega}(V(R_{\varphi(i)}))$  be a contraction model of  $Q_i$  in  $R_i$ . Recall that since  $Q_i$  and  $R_{\varphi(i)}$  are oriented paths, the contraction sending  $R_{\varphi(i)}$  on  $Q_i$  preserves endpoints. We now consider the function  $\mu$  defined as follows

$$\mu \begin{cases} V(f(Q, H)) & \rightarrow \mathcal{P}^{<\omega}(V(f(R, H))) \\ x & \rightarrow \{x\} & \text{if } x \in V(H) \\ \text{lst}(Q_i) & \rightarrow \mu_i(\text{lst}(Q_i)) \cup \bigcup_{j=\varphi(i)+1}^{\varphi(i+1)-1} V(R_j) \setminus \{\text{fst}(R_{\varphi(i+1)})\} \\ x & \rightarrow \mu_i(x) \subseteq R_{\varphi(i)} & \text{if } x \in Q_i \setminus \{\text{lst}(Q_i)\}. \end{cases}$$

We will show that  $\mu$  is an induced minor model of  $f(Q, H)$  in  $f(R, H)$ . First at all, remark that every element of the image of  $f$  induces in  $f(R, H)$  a connected subgraph:

- either  $x \in V(H)$  and  $\mu(x)$  is a singleton;
- or  $x \in \text{lst}(Q_i) \setminus \{\text{lst}(Q_i)\}$  and  $f(R, H)[\mu(x)]$  is connected since  $\mu_i(x) = \mu(x)$  is an induced minor model;
- or  $x = \text{lst}(Q_i)$  and  $\mu_i(\text{lst}(Q_i)) \cup \bigcup_{j=\varphi(i)+1}^{\varphi(i+1)-1} V(R_j) \setminus \{\text{fst}(R_{\varphi(i+1)})\}$  induces in  $f(R, H)$  a connected subgraph because  $f(R, H)[\mu_i(\text{lst}(Q_i))]$  is connected and the other vertices are consecutive on the cycle.

Let us now show that adjacencies are preserved by  $\mu$ . Let  $u, v$  be two distinct vertices of  $f(Q, H)$ . If  $u, v \in H$ , then  $\mu(u)$  and  $\mu(v)$  are adjacent in  $f(R, H)$  iff  $u$  and  $v$  are in  $f(Q, H)$ , as  $\mu(u) = \{u\}$  and  $\mu(v) = \{v\}$  (informally, the “ $H$ -part” of  $f(R, H)$  is not changed by the model). If  $u, v \in Q$ , observe that  $u$  and  $v$  are adjacent in  $f(Q, H)$

iff they belong to the same path of  $\{Q_i\}_{i \in [0, k-1]}$ . Thus in this case, the property that  $u$  and  $v$  are adjacent in  $f(Q, H)$  iff  $\mu(u)$  is adjacent to  $\mu(v)$  in  $f(R, H)$  is given by the fact that  $\{\mu_i\}_{i \in [0, k-1]}$  are contraction models.

If  $u \in Q$  and  $v \in H$ , then  $\{u, v\} \in f(Q, H)$  (respectively  $\mu(u)$  is adjacent to  $\mu(v)$  in  $f(R, H)$ ) iff  $u$  is an endpoint of a path of  $\{Q_i\}_{i \in [0, k-1]}$  (respectively  $\mu(u)$  contains an endpoint of a path of  $\{R_i\}_{i \in [0, l-1]}$ ), by definition of  $f$ . As the contraction relation on oriented paths of  $\mathcal{OP}$  is required to contract endpoints to endpoints, the image  $\mu(u)$  must contain the endpoint of a path of  $\{R_i\}_{i \in [0, l-1]}$  iff  $u$  is the endpoint of a path of  $\{Q_i\}_{i \in [0, k-1]}$ . Therefore  $u$  and  $v$  are adjacent in  $f(Q, H)$  iff  $\mu(u)$  is adjacent to  $\mu(v)$  in  $f(R, H)$ , as required. We finally proved that  $f$  is monotone. This was the only missing step in order to prove that  $(\text{lab}_{(S, \preceq)}(\mathcal{W}), \leq_{i.\text{sg.}})$  is a wqo.  $\square$

### 3.6.3 Graphs not containing Gem

The purpose of this section to give a proof to [Theorem 3.42](#). This will be done by first proving a decomposition theorem for graphs of  $\text{Excl}_{\leq_{i.m.}}(\text{Gem})$ , and then using this theorem to prove that  $(\text{Excl}_{\leq_{i.m.}}(\text{Gem}), \leq_{i.m.})$  is a wqo.

#### A Decomposition theorem for $\text{Excl}_{\leq_{i.m.}}(\text{Gem})$

This section is devoted to the proof of [Theorem 3.40](#), which is split in several lemmas. In the sequel,  $G$  is a 2-connected graph of  $\text{Excl}_{\leq_{i.m.}}(\text{Gem})$ . When  $G$  is 3-connected, we will rely on the following result originally proved by Ponomarenko.

**Proposition 3.70** ([\[Pon91\]](#)). *Every 3-connected Gem-induced minor-free graph is either a cograph or has an induced subgraph  $S$  isomorphic to  $P_4$ , such that every connected component of  $G \setminus S$  is a cograph.*

Therefore we will here focus on the case where  $G$  is 2-connected but not 3-connected. In this section, we deal with *edge-rooted* graphs, that are graphs with one distinguished edge. For two edge rooted graphs  $G, H$ , we say that  $H$  is an induced minor of  $G$  if there is an induced minor model  $\mu$  of  $H$  in  $G$  such that, if  $\{u, v\}$  is the root of  $H$ , then the root of  $G$  has the one endpoint in  $\mu(u)$  and the other one in  $\mu(v)$ . Intuitively, the model preserves the root. A *rooted diamond* is a graph which can be constructed from a rooted  $C_4$  by adding a chord incident with exactly one endpoint of the root (cf. [Figure 3.18](#)).



Figure 3.18: A rooted diamond, the root being the thick edge.

**Lemma 3.71.** *Let  $S = \{v_1, v_2\}$  be a cutset in a graph  $G$  and let  $C$  be a component of  $G \setminus S$ . Let  $H$  be the graph  $G[V(C) \cup \{v_1, v_2\}]$  rooted at  $\{v_1, v_2\}$ . If  $C$  has a rooted diamond as an induced minor, then  $\text{Gem}_{\leq_{i.m.}} G$ .*

*Proof.* Let  $C'$  be a component of  $G \setminus S$  other than  $C$  and let  $G'$  be the graph obtained from  $G$  by:

1. applying the necessary operations (contractions and vertex deletions) to transform  $G[V(C) \cup \{v_1, v_2\}]$  into a rooted diamond;
2. deleting every vertex not belonging to  $V(C) \cup V(C') \cup \{v_1, v_2\}$ ;
3. contracting  $C'$  to a single vertex.

The graph  $G'$  is then a rooted diamond and a vertex adjacent to both endpoints of its root, that is,  $G'$  is isomorphic to Gem.  $\square$

Let us now characterize the 2-connected graphs avoiding rooted diamonds.

**Lemma 3.72.** *Let  $G$  be a 2-connected graph rooted at  $\{u, v\} \in E(G)$ . If  $\{u, v\}$  is not a cut of  $G$  and  $G$  does not contain a rooted diamond as an induced minor, then either  $G$  is an induced cycle or both  $u$  and  $v$  are dominating in  $G$ .*

*Proof.* Assuming that  $u$  is not dominating and  $G$  is not an induced cycle, let us prove that  $G$  contains a rooted diamond as induced minor. Let  $w \in V(G)$  be a vertex such that  $\{u, w\} \notin E(G)$ . Such a vertex always exists given that  $u$  is not dominating. Let  $C$  be a shortest cycle using the edge  $\{u, v\}$  and the vertex  $w$  (which exists since  $G$  is 2-connected), let  $P_u$  be the subpath of  $C$  linking  $u$  to  $w$  without meeting  $v$  and similarly let  $P_v$  be the subpath of  $C$  linking  $v$  to  $w$  without meeting  $u$ . By the choice of  $C$ , both  $P_u$  and  $P_v$  are induced paths. Notice that if there is an edge connecting a vertex of  $P_u \setminus \{w\}$  to vertex of  $P_v \setminus \{w\}$ , then  $G$  contains a rooted diamond as induced minor. Therefore we can now assume that  $C$  is an induced cycle.

Recall that we initially assumed that  $G$  is not an induced cycle. Therefore  $G$  contains a vertex not belonging to  $C$ . Let  $G'$  be the graph obtained from  $G$  by contracting to one vertex  $x$  any connected component of  $G \setminus C$  and deleting all the other components. Obviously we have  $G' \leq_{\text{i.m.}} G$ . Let us show that  $G'$  contains a rooted diamond as induced minor.

Remark that the neighborhood of  $x$ , which is of size at least two (as  $G$  is 2-connected), is not equal to  $\{u, v\}$ , otherwise  $\{u, v\}$  would be a cut in  $G$ . Now contract in  $G'$  all the edges of  $C \setminus \{u, v\}$  except three in a way such that  $|N(x)| \geq 2$  and  $N(x) \neq \{u, v\}$ . Let  $G''$  be the obtained graph, which consists of a cycle of length four rooted at  $\{u, v\}$  and a vertex  $x$  adjacent to at least two vertices of this cycle. We shall here recall that since this cycle is a contraction of the induced cycle  $C$ , it is induced too. If  $x$  is adjacent (among others) to two vertices at distance two on this cycle, then, by contracting the edge between  $x$  and one of these vertices, we get a rooted diamond. The remaining case is when  $x$  is only adjacent to the vertices of the cycle which are not  $u$  and  $v$ . The contraction of the edge between  $v$  and one of these vertices gives a rooted diamond, and this concludes the proof.  $\square$

*Remark 3.73.* In a Gem-induced minor-free graph  $G$ , every induced subgraph  $H$  dominated by a vertex  $v \in V(G) \setminus V(H)$  is a cograph.

Indeed, assuming that  $H$  is not a cograph, let  $P$  be an path on four vertices which is induced subgraph of  $H$ . Then  $G[V(P) \cup \{v\}]$  is isomorphic to Gem, a contradiction.

Recall that we say that an induced subgraph of a graph  $G$  is basic in  $G$  if it is either a cograph, or an induced path whose internal vertices are of degree two in  $G$ .

**Lemma 3.74.** *If  $G$  has a  $K_2$ -cut  $S = \{v_1, v_2\}$ , then every connected component of  $G \setminus S$  is basic in  $G$ .*

*Proof.* By **Lemma 3.71**, for every connected component  $C$  of  $G \setminus S$  we know that the graph  $G[V(C) \cup S]$  rooted at  $\{u, v\}$  contains no rooted diamond. By the virtue of **Lemma 3.72**, this graph either is an induced cycle or has a dominating vertex among  $u$  and  $v$ . In the first case,  $C$  is a path whose all internal vertices are of degree two in  $G$ , hence  $H$  is basic. If one of  $u$  and  $v$  is dominating, then  $C$  is a cograph according to **Remark 3.73**. Therefore in both cases  $C$  is basic in  $G$ .  $\square$

Let us now focus on 2-connected graphs with a  $\overline{K_2}$ -cut, which is the last case in our characterization theorem.

**Corollary 3.75.** *If  $G$  has a  $\overline{K_2}$ -cut  $S$  such that  $G \setminus S$  contains more than two connected components, then every connected component of  $G \setminus S$  is basic in  $G$ .*

*Proof.* It follows directly from **Lemma 3.74**. Indeed, if the connected components of  $G \setminus S$  are  $C_1, C_2, \dots, C_k$ , let us contract  $C_1$  to an edge between the two vertices of  $S$ . The obtained graph fulfills the assumptions of **Lemma 3.74**:  $S$  is a  $K_2$ -cut. Therefore each of the components  $C_2, \dots, C_k$  is basic in  $G$ . Applying the same argument with  $C_2$  instead of  $C_1$  yields that  $C_1$  is basic in  $G$  as well.  $\square$

**Lemma 3.76.** *Let  $S = \{u, v\}$  be a  $\overline{K_2}$ -cut, such that and  $G \setminus S$  has only two connected components  $H_1$  and  $H_2$ . Then  $G$  contains a cycle  $C$  as induced subgraph such that every connected component of  $G \setminus C$  is basic in  $G$ .*

*Proof.* For every  $i \in \{1, 2\}$ , let  $Q_i$  be a shortest path linking  $u$  to  $v$  in  $G[V(H_i) \cup \{u, v\}]$ . Notice that the cycle  $C = G[V(Q_1) \cup V(Q_2)]$  is then an induced cycle. For contradiction, let us assume that some connected component  $J$  of  $G[V \setminus C]$  is not basic in  $G$ . By symmetry, we can assume that  $J \subset H_1$ .

Notice that since  $G$  is 2-connected,  $J$  has at least two distinct neighbors  $x, y$  on  $C$ . Let  $G'$  be the graph obtained from  $G[V(H_1) \cup V(C) \cup V(H_2)]$  by contracting  $Q_1$  to an edge between  $u$  and  $v$  in a way such that  $x$  is not contracted to  $y$  (that is,  $x$  is contracted to one of  $u, v$  and  $y$  to the other one). In  $G'$ ,  $\{u, v\}$  is a  $K_2$ -cut, therefore by **Lemma 3.74**, every connected component of  $G \setminus S$  is basic in  $G'$ . As this consequence holds for every choice of  $J$  and  $G'$  is an induced minor of  $G$ , we eventually get that every connected component of  $G \setminus C$  is basic in  $G$ .  $\square$

In the sequel,  $S, u, v$  and  $C$  follow the definitions of the statement of **Lemma 3.76**. In order to be more accurate on how the connected components of  $G \setminus C$  are connected to  $C$ , we will prove the following lemma according to which most of the vertices of  $C$  have degree 2 in  $G$ .

*Remark 3.77.* Every connected component  $J$  of  $G \setminus C$  has at least two and at most three neighbours on  $C$ .

Indeed, it has at least two neighbours on  $C$  because  $G$  is 2-connected. Besides if  $J$  has at least four neighbours on  $C$ , then contracting in  $G[V(C) \cup V(J)]$  the component  $J$  to a single vertex, deleting a vertex of  $C$  not belonging to  $N(J)$  (which exists since  $J$  belongs to only one of the components of  $G \setminus S$ ) and then contracting every edge incident with a vertex of degree two would yield Gem.

**Lemma 3.78.** *If  $C$  has at least one vertex of degree two, then for every distinct connected components  $J_1$  and  $J_2$  of  $G \setminus C$  we have  $N_C(J_1) \subseteq N_C(J_2)$  or  $N_C(J_2) \subseteq N_C(J_1)$ .*

*Proof.* Let us assume, for contradiction, that the claim is not true and let  $G$  be a minimal counterexample with respect to induced minors. In such a case both  $J_1$  and  $J_2$  are single vertices (say  $j_1$  and  $j_2$  respectively) and they are the only connected components of  $G \setminus C$ . We now would like to argue that any such minimal counterexample must contain as induced minor one of graphs presented on [Figure 3.19](#) (where thick edges represent the cycle  $C$ ). This would conclude the proof as each of these graphs contains Gem as induced minor, as shown in [Figure 3.20](#).

First of all, in such a minimal counterexample there is only one vertex in  $C$  of degree 2, let us call it  $c$ . We will consider all the ways that the vertices  $j_1$  and  $j_2$  can be connected to the neighbors of  $c$  and show that in every such case we can contract our graph to one of the graphs on [Figure 3.19](#). According to [Remark 3.77](#), each of  $j_1$  and  $j_2$  will have either two or three neighbors on  $C$ .

*First case:* both  $j_1$  and  $j_2$  are connected with both neighbours of  $c$ . As  $N(j_1) \not\subseteq N(j_2)$  and  $N(j_2) \not\subseteq N(j_1)$ , each of  $j_1, j_2$  has a neighbor which is not adjacent to the other. But since  $j_1$  and  $j_2$  can have at most three neighbors, the neighborhood of  $j_1$  and  $j_2$  is now completely characterized. The leftmost part of [Figure 3.19](#) presents the only possible graph for this case.




Figure 3.19: Induced minor-minimal counterexamples in [Lemma 3.78](#).

*Second case:*  $j_1$  is connected with exactly one of neighbours of  $c$  and  $j_2$  is connected with the other one. In this case, as each of  $j_1, j_2$  has at least two neighbors on  $C$ , contracting all the edges of  $C$  whose both endpoints are at distance at least two from  $c$  gives the graph depicted in the center of [Figure 3.19](#).

*Third case:*  $j_1$  is connected with both neighbours of  $c$ , and  $j_2$  is connected with at most one of them. In this case, as long as  $C$  has more than 4 edges, we can contract an edge of  $C$  to find a smaller counterexample. Precisely, if there are at least 4 edges, there are two edges  $e_1, e_2$  in  $C$  within distance exactly 1 to  $c$  and those two do not share an endpoint. Moreover  $j_2$  has a neighbour  $s$  in  $C \setminus N(c)$ , say  $x$ , which is not a neighbour

of  $j_1$ . Now one of the edges  $e_1, e_2$  is not incident to  $s$  and contracting this edge would yield a smaller counterexample.

Therefore, we only have to care about the case where  $C$  has exactly 4 edges and this case is exactly the graph represented on the right of [Figure 3.19](#).

We have considered all possible induced minor-minimal counterexamples (up to symmetry between  $j_1$  and  $j_2$ ). For each of these cases, which are presented on [Figure 3.19](#), we will now give an induced minor model of Gem, which proves that they all contain Gem as induced minor. For each graph depicted on [Figure 3.20](#) we consider the model mapping the vertex  $v_i$  of Gem to the set of vertices labeled  $M_i$ . It is easy to check that each of these sets induces a connected subgraph and that the adjacencies between two sets correspond to the ones between the corresponding vertices of Gem. This concludes the proof.  $\square$




Figure 3.20: Models of Gem in graphs from [Figure 3.19](#).

**Corollary 3.79.** *If  $C$  has at least one vertex of degree two, then it has at most three vertices of degree greater than two.*

*Proof.* Notice that the set of vertices of  $C$  that have degree greater than two is exactly the union of  $N_C(J)$  over all connected components  $J$  of  $G \setminus C$ . We just saw in [Lemma 3.78](#) that for every two connected components of  $G \setminus C$ , the neighborhood on  $C$  of one is contained in the neighborhood on  $C$  of the other and that these neighborhoods have size at most three. Therefore their union have size at most three as well.  $\square$

**Corollary 3.80.** *Every connected component of  $G \setminus C$  is basic and  $C$  has at most six vertices of degree greater than two.*

*Proof.* Remark that contracting  $H_1$  to a single vertex  $h$  in  $G$  gives a graph  $G'$  and a cycle  $C'$  (contraction of  $C$ ) such that every connected component of  $G' \setminus C'$  is basic and  $C'$  has at least one vertex of degree 2, which is  $h$ . By [Corollary 3.79](#),  $C'$  has at most three vertices of degree greater than two. Notice that these vertices belong to  $G' \setminus h$  which is isomorphic to  $G \setminus H_1$ . Hence  $G \setminus H_1$  has at most three vertices of degree greater than two. Applying the same argument with  $H_2$  instead of  $H_1$  we get the desired result.  $\square$

Now we are ready to prove main decomposition theorem for Gem-induced minor-free graphs. Let us remind its statement.

**Theorem 3.40** (Decomposition of Gem-induced minor-free graph). *Let  $G$  be a 2-connected graph such that  $\text{Gem} \not\prec_{\text{i.m.}} G$ . Then  $G$  has a subset  $X \subseteq V(G)$  of at most six vertices such that every connected component of  $G \setminus X$  is either a cograph or a path whose internal vertices are of degree two in  $G$ .*

*Proof.* Recall that we are looking for a subset  $X$  of  $V(G)$  of size at most 6 such that each component of  $G \setminus X$  is basic in  $G$ .

If  $G$  is 3-connected, by [Proposition 3.70](#) it is either a cograph, or has a subset  $X$  of four vertices such that every connected component of  $G \setminus X$  is a cograph. Let us now assume that  $G$  is not 3-connected.

In the case where  $G$  has a  $K_2$ -cut  $S$ , or if  $G$  has a  $\overline{K_2}$ -cut  $S$  such that  $G \setminus S$  has more than two connected components, then according to [Lemma 3.74](#) and [Corollary 3.75](#) respectively,  $S$  satisfies the required properties. In the remaining case, by [Corollary 3.80](#)  $G$  has a cycle  $C$  such that every connected component of  $G \setminus C$  is basic in  $G$  and which has at most six vertices of degree more than two in  $G$ . Let  $X$  be the set containing those vertices of degree more than two. Observe that every connected component of  $G \setminus X$  is either a connected component of  $G \setminus C$  (hence it is basic) or a part of  $C$ , i.e. a path whose internal vertices are of degree two in  $G$  (which is basic as well). As  $|X| \leq 6$ ,  $X$  satisfies the desired properties.  $\square$

## Well-quasi-ordering of labelled cographs

We were able to show that structure of 2-connected Gem-induced minor-free graphs is essentially very simple, with building blocks being cographs and long induced paths. To conclude that labelled 2-connected Gem-induced minor-free graphs are wqo by induced minor relation we will need the fact that the building blocks, in particular labelled cographs, are themselves well-quasi-ordered by the induced minor relation.

This following result has been proven by Damaschke in [[Dam90](#)] in the unlabelled case. The proof for the labelled case follows the same general approach. We present below the sketch of the proof.

Let us denote  $\mathcal{C}$  to be a class of all cographs.

**Theorem 3.81.** *For any wqo  $(Q, \preceq_Q)$ , the class  $\text{lab}_{(Q, \preceq_Q)}(\mathcal{C})$  is wqo with respect to  $\leq_{\text{i.sg.}}$ .*

*Proof.* Let us define as usual  $(S, \preceq)$  to be  $(\mathcal{P}^{<\omega} Q, \preceq^{\mathcal{P}})$ . Define  $(S^+, \preceq^+)$  such that  $S^+$  is disjoint union of  $S$  and  $\{0, 1\}$ ; the order  $\preceq^+$  is such that  $\preceq^+$  is just  $\preceq$  when restricted to  $S$ , but 0, 1 and elements of  $S$  are incomparable.

By virtue of [Remark 3.25](#) and [Corollary 3.27](#), we know that  $(S^+, \preceq^+)$  is wqo. By the labelled version of the famous Kruskal theorem (see [[Kru60](#)]), the class of all finite trees labelled by  $(S^+, \preceq^+)$  is wqo, with respect to a topological minor relation. In particular, we can consider class  $\mathcal{T}$  of finite trees, labelled by  $(S^+, \preceq^+)$ , such that all internal nodes have labels  $\{0, 1\}$ , and all leaves has label from  $S$ . We will consider this class again with the ordering by a labelled topological minor relation. As it is a subclass of a wqo, the class  $\mathcal{T}$  itself is also wqo. We will now provide a epi  $\phi : \mathcal{T} \rightarrow \text{lab}_{(Q, \preceq_Q)}(\mathcal{C})$ , and we will conclude by [Remark 3.28](#), that  $\text{lab}_{(Q, \preceq_Q)}(\mathcal{C})$  is wqo.

The function  $\phi$  is defined as follows: given a labelled tree  $T$ , if the whole tree is only a single leaf, it produces a graph with a single vertex, and with the same label as the one the leaf has in  $T$ . If the tree is larger than a single vertex, it has root  $r$  with label  $s$ , and subtrees  $T_1, T_2, \dots, T_k$ , all rooted at some children of  $r$ . Then  $\phi(T)$  is defined as disjoint union of  $\phi(T_i)$  if label  $s$  were 0, or join of  $\phi(T_i)$  of label of  $s$  were 1. It is well-known, that every cograph has such a presentation, i.e. that function  $\phi$  indeed is surjective. The tree  $T$  which is mapped  $G$  by  $\phi$  is usually called *cotree* of  $G$ .

Now we only need to prove that  $\phi$  is monotone. Indeed, consider two labelled trees  $T_1, T_2 \in \mathcal{T}$ , such that  $T_1 \leq T_2$ , and let  $i : V(T_1) \mapsto V(T_2)$  to be an embedding of  $T_1$  in  $T_2$  as a topological minor, such that  $\lambda_{T_1}(u) \preceq^+ \lambda_{T_2}(i(u))$ . In particular, by the second property, we conclude that  $i$  maps leaves of  $T_1$  to leaves of  $T_2$ . Therefore we can consider a corresponding mapping  $\tilde{i}$  from  $V(\phi(T_1))$  to  $V(\phi(T_2))$ . Clearly it is injective, and has the property that  $\lambda_{\phi(T_1)}(u) \preceq \lambda_{\phi(T_2)}(\tilde{i}(u))$ . To show that it defines an induced minor model of  $\phi(T_1)$  as an induced subgraph of  $\phi(T_2)$ , we only need to prove that  $\tilde{i}(u)$  and  $\tilde{i}(v)$  are connected by an edge iff  $u$  and  $v$  are connected by an edge.

Remark that two vertices  $u, v \in V(\phi(T))$  are connected by an edge in  $\phi(T)$  iff the label of a lowest common ancestor of the corresponding leaves in  $T$  is 1. But  $i$  was an embedding of  $T_1$  in  $T_2$  as a topological minor, so in particular lowest common ancestor of  $i(u)$  and  $i(v)$  is the same as an image of lowest common ancestor of  $u, v$ . Moreover, by the definition of the order  $S^+$ , embedding  $i$  preserves exactly labels of internal nodes. Hence  $\phi$  is indeed monotone and this concludes the proof of the theorem.  $\square$

## Well-quasi-ordering Gem-induced minor-free graphs

In this section we will give a proof of [Theorem 3.42](#).

We define  $\mathcal{B}$  as the class of graphs which are disjoint unions of induced paths and cographs, and  $\mathcal{C}$  as the class of cographs.

**Lemma 3.82.** *Let  $k \in \mathbb{N}$ , let  $(S, \preceq)$  be a wqo and let  $\mathcal{G}$  be the class of  $(S, \preceq)$ -labelled graphs such that the removal of at most  $k$  vertices yields a graph of  $\mathcal{B}$ . Then  $(\mathcal{G}, \leq_{i.m.})$  is a wqo.*

*Proof.* Let  $k$  be fixed.

For every  $G \in \mathcal{G}$ , let  $X_G$  be a set of at most  $k$  vertices of  $G$  such that  $G \setminus X \in \mathcal{B}$ .

For every graph  $H$  on at most  $k$  vertices, let  $\mathcal{G}_H = \{G \in \mathcal{G}, G[X_G] = H\}$ . Observe that this gives a partition of  $\mathcal{G}$  into a finite number of subclasses. By the virtue of [Remark 3.25](#), we only need to focus on one of these classes. For the sake of simplicity, we assume that  $H$  has exactly  $k$  vertices  $\{v_1, \dots, v_k\}$ .

Informally, our goal is now to define a function  $f$  which constructs a graph of  $\mathcal{G}_H$ , given an encoding in terms of graphs of  $\mathcal{B}$ . We will then show that  $f$  is an epi.

Let  $f$  be the function whose domain is the quasi-order

$$(\mathcal{D}, \preceq_{\mathcal{D}}) = (S, \preceq)^k \times (\mathcal{P}^{<\omega}(\text{lab}_{(S, \preceq)}(\mathcal{OP})), \leq_{c. \star})^{\binom{k}{2}} \times (\text{lab}_{(S, \preceq) \times (2^{\llbracket 1, k \rrbracket}, =)}(\mathcal{C}), \leq_{i.sg.})$$

(where  $\preceq_{\mathcal{D}} = \preceq^k \times (\leq_{c. \star})^{\binom{k}{2}} \times \leq_{i.sg.}$ ) and which, given a tuple

$$((s_i)_{i \in \llbracket 1, k \rrbracket}, (L_{i,j})_{i,j \in \llbracket 1, k \rrbracket, i < j}, J)$$

such that

- $(s_i)_{i \in \llbracket 1, k \rrbracket} \in S^k$  is a tuple of  $k$  labels from  $S$ ,
- $(L_{i,j})_{i,j \in \llbracket 1, k \rrbracket, i < j}$  is a tuple of  $\binom{k}{2}$  subsets of  $(S, \preceq)$ -labeled oriented paths, and
- $J \in \text{lab}_{(S, \preceq) \times (2^{\llbracket 1, k \rrbracket}, =)}(\mathcal{C})$ , is a  $(S, \preceq) \times (2^{\llbracket 1, k \rrbracket}, =)$ -labeled cograph,

returns the graph constructed as follows, starting from  $H$ :

1. label  $s_i$  the vertex  $v_i$ , for every  $i \in \llbracket 1, k \rrbracket$ ;
2. for every  $i, j \in \llbracket 1, k \rrbracket^2$ ,  $i < j$ , and for every path  $L \in L_{i,j}$ , add a copy of  $L$  to the current graph, connect  $v_i$  to  $\text{fst}(L)$  and  $v_j$  to  $\text{lst}(L)$ ;
3. add to the current graph a copy of the underlying graph of  $J$  and, for every vertex  $v$  labeled  $(s, \{e_1, \dots, e_l\})$  (for some  $l \in \llbracket 1, k \rrbracket$ ), give the label  $s$  to  $v$  in the current graph and make  $v$  adjacent to vertices  $v_{e_1}, \dots, v_{e_l}$ .

By construction, the codomain of  $f$  is included in  $\text{lab}_{(S, \preceq)}(\mathcal{G}_H)$ . Let us now show that  $f$  is surjective on  $\text{lab}_{(S, \preceq)}(\mathcal{G}_H)$ . Let  $G \in \text{lab}_{(S, \preceq)}(\mathcal{G}_H)$  and let us consider the connected components of  $G \setminus X_G$ . Let  $J$  be the disjoint union of all such components that are cographs. Note that  $J$  is a cograph as well. For every vertex  $v$  of  $J$  of label  $s$ , we relabel  $v$  with the label  $(s, \{e_1, \dots, e_l\})$ , where  $\{e_1, \dots, e_l\}$  are all the integers  $i \in \llbracket 1, k \rrbracket$  such that  $v$  is adjacent to  $v_i$ . For every  $i, j \in \llbracket 1, k \rrbracket$ ,  $i < j$ , let  $L_{i,j}$  be the set of paths of  $G \setminus X_H$  which are neighbors in  $G$  of  $v_i$  and  $v_j$ , to which we give the following orientation: the first vertex of such a path is the one which is adjacent to  $v_i$  and its last vertex is the one adjacent to  $v_j$ . Last, let  $s_i$  be the label of  $v_i$  for every  $i \in \llbracket 1, k \rrbracket$ . Then it is clear that  $G$  is isomorphic to  $f(\{s_i\}_{i \in \llbracket 1, k \rrbracket}, \{L_{i,j}\}_{i,j \in \llbracket 1, k \rrbracket, i < j}, J)$ . Consequently  $f$  is surjective on  $\text{lab}_{(S, \preceq)}(\mathcal{G}_H)$ .

Our current goal is now, in order to show that  $f: (\mathcal{B}, \preceq_{\mathcal{B}}) \rightarrow (\text{lab}_{(S, \preceq)}(\mathcal{G}_H), \leq_{\text{i.m.}})$  is an epi, is to prove that it is monotone. Let  $A, B$  be two elements of

$$S^k \times (\mathcal{P}^{<\omega}(\text{lab}_{(S, \preceq)}(\mathcal{OP})))^{\binom{k}{2}} \times \text{lab}_{(S, \preceq) \times (2^{\llbracket 1, k \rrbracket}, =)}(\mathcal{C})$$

such that  $A \preceq_{\mathcal{D}} B$ . Let us show that  $f(A) \leq_{\text{i.m.}} f(B)$ . According to [Remark 3.29](#), it is enough to focus on the cases where  $A$  and  $B$  differ by only one coordinate.

*First case:*  $A$  and  $B$  differ by the  $i$ -th coordinate, for  $i \in \llbracket 1, k \rrbracket$ . Let  $s_A$  (respectively  $s_B$ ) be the value of the  $i$ -th coordinate of  $A$  (respectively of  $B$ ). According to the definition of  $f$ , the graphs  $f(A)$  and  $f(B)$  differ only by the label of vertex  $v_i$ : this label is  $s_A$  in  $f(A)$  whereas it equals  $s_B$  in  $f(B)$ . But since we have  $s_A \preceq s_B$  (as  $A \preceq_B B$ ), we get  $f(A) \leq_{\text{i.m.}} f(B)$ .

*Second case:*  $A$  and  $B$  differ by the last coordinate. Let  $J_A$  (respectively  $J_B$ ) be the value of the last coordinate of  $A$  (respectively of  $B$ ). As previously,  $A \preceq_B B$  gives  $J_A \leq_{\text{i.sg.}} J_B$ , therefore we can obtain  $J_A$  by removing vertices of  $J_B$  and contracting labels. As the adjacencies of vertices of  $J_A$  and  $J_B$  to the rest of  $f(A)$  and  $f(B)$  (respectively) depend only on the label of their vertices, the same deletion and contraction operations in  $f(B)$  give  $f(A)$ , hence  $f(A) \leq_{\text{i.m.}} f(B)$ .

*Third case:*  $A$  and  $B$  differ by the  $i$ -th coordinate, for some  $i \in \llbracket k+1, k + \binom{k}{2} \rrbracket$ . Let  $L_A$  (respectively  $L_B$ ) be the value of this coordinate in  $A$  (respectively in  $B$ ). As previously again,  $A \preceq_B B$  gives  $L_A \leq_c L_B$ , consequently we can obtain  $L_A$  by contracting edges of  $J_B$  and contracting labels. Since the contraction relation on  $\mathcal{OP}$  requires that endpoints (beginning and end of a path) are preserved, the same contraction operations in  $f(B)$  give  $f(A)$ , thus we again get  $f(A) \leq_{i.m.} f(B)$ .

We just proved that  $f$  is monotone, therefore it is an epi. By [Remark 3.28](#), it is enough to show that  $(\mathcal{B}, \preceq_B)$  is a wqo in order to prove that  $(\text{lab}_{(S, \preceq)}(\mathcal{G}_H), \leq_{i.m.})$  is a wqo.

Notice that  $(\mathcal{B}, \preceq_B)$  is a Cartesian product of wqos and of the set of finite subsets of a wqo. Indeed, we assumed that  $(S, \preceq)$  is a wqo. Furthermore, we proved in [Lemma 3.65](#) that for every wqo  $(S, \preceq)$ , the quasi-order  $(\text{lab}_{(S, \preceq)}(\mathcal{OP}), \leq_c)$  is a wqo, and hence so is  $(\mathcal{P}^{<\omega}(\text{lab}_{(S, \preceq)}(\mathcal{OP})), \leq_c^*)$  (cf. [Corollary 3.27](#)). Last, we proved in [Theorem 3.81](#) that the class of cographs labeled by a wqo is well-quasi-ordered by the induced subgraph relation. Therefore,  $(\mathcal{B}, \preceq_B)$  is a wqo, which concludes the proof.  $\square$

We can now prove [Theorem 3.42](#), that we restate here as a reminder.

**Theorem 3.42.** *The class of Gem-induced minor-free graphs is wqo by  $\leq_{i.m.}$ .*

*Proof.* According to [Lemma 3.31](#), it is enough to prove that for every wqo  $(S, \preceq)$ , the class of  $(S, \preceq)$ -labeled 2-connected graphs which does not contain Gem as induced minor is well-quasi-ordered by induced minors. By [Theorem 3.40](#), these graphs can be turned into a disjoint union of paths and cographs by the deletion of at most six vertices. A consequence of [Lemma 3.82](#) (for  $k = 6$ ), these graphs are well-quasi-ordered by induced minors and we are done.  $\square$

## 3.7 Contractions and well-quasi-ordering

This section is devoted to the proof of [Theorem 3.13](#). We recall that, unless otherwise specified, this section deals with simple graphs.

### 3.7.1 On graphs with no diamond

In this section we show that graphs in  $\text{Excl}_{\leq_c}(D_2)$  have a simple structure. The *clique-cactus* graphs are the graphs whose blocks are cliques and cliques (cf. [Figure 3.21](#) for an example).

Recall that for every positive integer  $r$ , we denote by  $D_r$  the graph obtained by contracting an edge of  $K_{2,r+1}$ . In particular,  $D_2$  is the graph sometimes called diamond. For this section, we set  $\mathcal{D} = \{D_r, r \in \mathbb{N}\}$  (cf. [Figure 3.22](#)) and  $\mathcal{S} = \{K_{1,r}, r \in \mathbb{N}\}$ .

We prove the following lemma.

**Lemma 3.83** (Characterization of clique-cactus graphs). *Graphs of  $\text{Excl}_{\leq_c}(D_2)$  are exactly the clique-cactus graphs.*




Figure 3.21: A clique-cactus graph.




Figure 3.22: Graphs of  $\mathcal{D}$ .

The proof of **Lemma 3.83** will be given after a few lemmas. If  $C$  is a cycle of a graph  $G$  and  $\{u, v\}, \{u', v'\} \subseteq V(C)$ , we say that  $\{u, v\}$  and  $\{u', v'\}$  are *crossing in  $C$*  if  $u, u', v, v'$  are distinct and are appearing in this order on the cycle. A *non-chord* in a cycle is a set of two vertices that are not adjacent.

**Lemma 3.84.** *Let  $G$  be a graph and let  $C$  be a cycle in  $G$ . If  $C$  has at least one chord and one non-chord in  $G$ , then it has one chord and one non-chord that are crossing in  $C$ .*

*Proof.* Let  $\{x, x'\}$  be a non-chord of  $C$  in  $G$  and let  $P$  and  $Q$  be the connected components of  $C \setminus \{x, x'\}$  which obviously are paths. Let us assume that every chord of  $C$  in  $G$  has both endpoints either in  $P$  or in  $Q$  (otherwise we are done) and let  $\{y, y'\}$  be a chord of  $C$  in  $G$ , the endpoints of which belong, say, to  $P$ . Let  $z$  be a vertex of the subpath of  $P$  delimited by  $y$  and  $y'$  such that  $z \notin \{y, y'\}$  and let  $z'$  be a vertex of  $Q$ . If  $\{z, z'\}$  is a chord of  $C$  in  $G$ , then  $\{x, x'\}$  and  $\{z, z'\}$  are satisfying the required property. Otherwise,  $\{z, z'\}$  is a non-chord and now  $\{y, y'\}$  and  $\{z, z'\}$  are crossing.  $\square$

**Lemma 3.85.** *Let  $G \in \text{Excl}_{\leq c.}(D_2)$ . Every cycle of  $G$  is either an induced cycle, or it induces a clique in  $G$ .*

*Proof.* Let  $G$  be a graph of  $\text{Excl}_{\leq c.}(D_2)$  and let  $C$  be a cycle of  $G$ . For contradiction, let us assume that  $C$  has at least one chord  $\{u, u'\}$  and one non-chord  $\{v, v'\}$ . According to **Lemma 3.84** we can assume without loss of generality that they are crossing in  $C$ . Let  $P$  and  $Q$  be the two connected components of  $C \setminus \{v, v'\}$ . In the graph  $G[C]$ , contracting  $P$  to a single vertex  $x$  and  $Q$  to  $y$  yields a graph on four vertices  $v, v', x, y$  such that

1.  $v, x, v', y$  lie on the cycle in this order;
2.  $\{v, v'\} \notin E(G)$ ; and

3.  $\{x, y\} \in E(G)$  (as  $\{u, u'\}$  connects the subgraphs that are respectively contracted to  $x$  and  $y$ ).

The obtained graph is  $D_2$ , a contradiction. Therefore  $C$  has either no chords or no non-chords in  $G$ . It is clear that in the first case  $C$  is an induced cycle of  $G$  and that in the second case it induces a clique.  $\square$

**Lemma 3.86.** *Let  $G \in \text{Excl}_{\leq c.}(D_2)$  be a 2-connected graph. Then  $G$  is either a cycle, or a clique.*

*Proof.* We assume that  $|V(G)| > 1$ , otherwise the result is trivial. Let  $C$  be a longest cycle of  $G$ . By [Lemma 3.85](#) the cycle  $C$  is either an induced cycle or it induces a clique in  $G$ . Let us treat these two cases separately. For contradiction, we assume that  $V(G) \setminus V(C)$  is not empty and we call  $H_1, \dots, H_t$  the connected components of  $G \setminus C$ , for some  $t \in \mathbb{N}_{\geq 1}$ . Let us consider the graph  $G'$  where  $H_i$ , which is connected, has been contracted to a single vertex  $h_i$ , for every  $i \in \llbracket 1, t \rrbracket$ . Observe that  $G'$  is 2-connected, given that  $G$  is 2-connected. Also,  $G' \in \text{Excl}_{\leq c.}(D_2)$ .

*First case:*  $C$  induces a clique in  $G'$ . Notice that  $C$  is then a maximal clique. Let  $u = h_1$ . As  $C$  is maximal, there is a vertex  $v \in V(C)$  such that  $\{u, v\} \notin E(G')$ . Let  $x$  and  $y$  be two neighbors of  $u$  on  $C$  (they exist since  $G'$  is 2-connected). These vertices define two subpaths of  $C$ . Let  $R$  be the longest of these paths that contains  $v$ . Observe that in this case,  $R$  has at least three vertices. The union of  $\{u, x\}$ ,  $\{u, y\}$  and  $R$  is a cycle of  $G'$  that we call  $C'$ . According to [Lemma 3.85](#), this cycle is either induced or it induces a clique. As  $\{u, v\} \notin E(G')$ ,  $C'$  cannot induce a clique in  $G$ . On the other hand,  $C$  is not an induced cycle as every pair of vertices of  $R$  are adjacent (and  $|V(R)| \geq 3$  as mentioned earlier). We reached the contradiction we were looking for.

*Second case:*  $C$  is an induced cycle and has at least 4 vertices. Let  $i \in \llbracket 1, t \rrbracket$ . As  $G'$  is 2-connected,  $h_i$  has at least two neighbors on  $C$ : let  $x$  and  $y$  be two of them.

*Claim 3.87.*  $x$  and  $y$  are not adjacent.

*Proof.* Let us assume that  $\{x, y\} \in E(G')$ . Let  $C'$  be the cycle obtained from  $C$  by replacing the edge  $\{x, y\}$  by the path  $xh_iy$ . This cycle is not induced as  $x, y$  are not adjacent in  $C'$  whereas  $\{x, y\} \in E(G)$ . It does not induce a clique neither since  $x$  is not adjacent with the other neighbor of  $y$  on  $C$  (which is not  $x$  as we assume that  $C$  has at least 4 vertices). This contradicts [Lemma 3.85](#) and therefore proves that  $\{x, y\} \notin E(G)$ .  $\square$

Every pair of distinct vertices of the cycle  $C$  defines two subpaths of  $C$  meeting only at these vertices. Let  $u$  and  $v$  be two vertices of  $C$  such that  $h_i$  has at least one neighbor in the interior of each of the subpaths of  $C$  defined by  $u$  and  $v$ , that we will respectively call  $P$  and  $Q$ . Such vertices exists, as a consequence of [Claim 3.87](#).

Let us consider the contraction  $H$  of  $G'$  obtained by contracting the interior path of  $P$  (respectively  $Q$ ) to a single vertex  $w_P$  (respectively  $w_Q$ ) and then by contracting the edge connecting  $h_i$  to  $w_P$ . This edge exists by definition of  $u$  and  $v$ . Then  $uw_Pvw_Q$  is a cycle of  $H$  where  $\{w_P, w_Q\}$  is a chord (because we contracted to  $w_P$  the vertex  $h_1$  which was adjacent to both  $w_P$  and  $w_Q$ ) and  $\{u, v\}$  is a non-chord (as they were

non-adjacent vertices of the induced cycle  $C$  and that noting has been contracted to them). According to [Lemma 3.85](#), the graph  $H$  contains  $D_2$  as contraction. As  $H$  is a contraction of  $G$ , then  $D_2 \leq_c G$ , a contradiction.

In both cases we reached a contradiction, therefore  $V(G) \setminus V(C)$  is empty:  $G$  is a clique or an induced cycle.  $\square$

We are now ready to prove [Lemma 3.83](#). Let us recall its statement.

**Lemma 3.83** (Characterization of clique-cactus graphs). *Graphs of  $\text{Excl}_{\leq_c}(D_2)$  are exactly the clique-cactus graphs.*

*Proof.* The fact that a graph of  $\text{Excl}_{\leq_c}(D_2)$  is a clique-cactus is a straightforward corollary of [Lemma 3.86](#). It is easy to see that a cactus graph does not contain  $D_2$  as contraction by noticing that  $D_2$  is a contraction of a graph if and only if it is a contraction of one of its 2-connected components. As  $D_2$  is neither a contraction of a cycle, nor of a clique, we get the desired result.  $\square$

### 3.7.2 Well-quasi-ordering clique-cactus graphs

We proved in the previous section that graphs of  $\text{Excl}_{\leq_c}(D_2)$  are exactly the connected clique-cactus graphs. This section contains the last part of the proof of [Theorem 3.13](#), which is the following lemma.

**Lemma 3.88.** *Connected clique-cactus graphs are well-quasi-ordered by  $\leq_c$ .*

In this section we deal with rooted graphs. Let us denote by  $\mathcal{C}$  the class of rooted connected clique-cactus graphs. In this class, two isomorphic graphs with a different root are seen as different. It is clear that proving that  $(\mathcal{C}, \leq_c)$  is a wqo implies [Lemma 3.88](#). This is what we will do.

**Building blocks.** Let us define three functions  $\text{stick}: \mathcal{C}^* \rightarrow \mathcal{C}$ ,  $\text{cycle}: \mathcal{C}^* \rightarrow \mathcal{C}$ , and  $\text{clique}: \mathcal{C}^* \rightarrow \mathcal{C}$  that will output graphs obtained by combining together the graphs given as parameters, in the following fashion. Given a sequence  $\langle G_0, \dots, G_{p-1} \rangle \in \mathcal{C}^*$  (for some  $p \in \mathbb{N}$ ), if  $U$  denote the union of the graphs  $G_1, \dots, G_{p-1}$ , then we define;

- $\text{stick}(G_0, \dots, G_{p-1})$  is the graph obtained from  $U$  by identifying the vertices

$$\text{root}(G_0), \dots, \text{root}(G_{p-1});$$

- $\text{cycle}(G_0, \dots, G_{p-1})$  is the graph obtained from  $U$  by adding the edges

$$\{\text{root}(G_i), \text{root}(G_{(i+1) \bmod p})\}$$

for every  $i \in \llbracket 0, p-1 \rrbracket$ ; and

- $\text{clique}(G_0, \dots, G_{p-1})$  is the graph obtained from  $U$  by adding the edges

$$\{\text{root}(G_i), \text{root}(G_j)\}$$

for every distinct  $i, j \in \llbracket 0, p-1 \rrbracket$ .

The root of  $\text{stick}(G_0, \dots, G_{p-1})$ ,  $\text{cycle}(G_0, \dots, G_{p-1})$  and  $\text{clique}(G_0, \dots, G_{p-1})$  is the vertex that is the root of  $G_0$ . These functions will allow us to encode graphs of  $\mathcal{C}$  into sequences.

We will now decompose graphs of  $\mathcal{C}$  along blocks.

For every block  $B$  of a graph  $G$ , let  $\text{dec}_B(G)$  denote the collection of all the graphs  $H$  that can be constructed from some connected component  $C$  of  $G \setminus V(B)$  by adding a new vertex  $v$  adjacent to the vertices of  $C$  that are adjacent to a vertex of  $B$  in  $G$  and setting  $\text{root}(H) = v$ .

Observe that, as soon as  $\text{root}(G) \in V(B)$ , every graph of  $\text{dec}_B(G)$  is a proper contraction of  $G$ . Let  $\text{dec}(G)$  denote the union of the sets  $\text{dec}_B(G)$  for every block  $B$  of  $G$  containing the root of  $G$ . The following observation is a consequence of [Lemma 3.83](#).

*Observation 3.89.* For every graph  $G \in \mathcal{C}$  there is a (non necessarily unique) sequence  $\langle \mathcal{G}_0, \dots, \mathcal{G}_{p-1} \rangle \in \text{dec}(G)^*$  (for some  $p \in \mathbb{N}$ ) such that either

$$\begin{aligned} G &= \text{cycle}(\text{stick}(\mathcal{G}_0), \dots, \text{stick}(\mathcal{G}_{p-1})), \text{ or} \\ G &= \text{clique}(\text{stick}(\mathcal{G}_0), \dots, \text{stick}(\mathcal{G}_{p-1})). \end{aligned}$$

**From encodings to well-quasi-ordering.** The following lemma will allow us to work on sequences in order to show that two graphs are comparable.

**Lemma 3.90.** *Let  $\sigma, \tau \in \mathcal{C}^*$ . If  $\sigma \leq_{c.*} \tau$ , then*

- (i)  $\text{cycle}(\sigma) \leq_c \text{cycle}(\tau)$ ;
- (ii)  $\text{clique}(\sigma) \leq_c \text{clique}(\tau)$ ; and
- (iii)  $\text{stick}(\sigma) \leq_c \text{stick}(\tau)$ .

*Proof.* Let  $\sigma = \langle H_1, \dots, H_p \rangle$  and  $\tau = \langle G_1, \dots, G_q \rangle$  (for some positive integers  $p, q$ ) and let  $H = \text{cycle}(\sigma)$  and  $G = \text{cycle}(\tau)$ . For the sake of readability we will refer to  $H_i$ 's (respectively  $G_i$ 's) either as elements of  $\sigma$  (respectively  $\tau$ ) or as subgraphs of  $H$  (respectively  $G$ ).

If  $\sigma \leq_{c.*} \tau$ , then there is, by definition of  $\leq_{c.*}$ , an increasing function  $\varphi: \llbracket 1, p \rrbracket \rightarrow \llbracket 1, q \rrbracket$  such that  $\forall i \in \llbracket 1, p \rrbracket, H_i \leq_c G_{\varphi(i)}$ . Therefore there is a sequence of edge contractions transforming  $G_{\varphi(i)}$  into  $H_i$  for every  $i \in \llbracket 1, p \rrbracket$ . Let us perform the following operations on  $G$ :

1. for every  $j \in \llbracket 1, q \rrbracket \setminus \{\varphi(i), i \in \llbracket 1, p \rrbracket\}$  we contract the subgraph  $G_j$  to a single vertex  $v_j$  and we then contract some edge incident with  $v_j$ ;
2. for every  $i \in \llbracket 1, p \rrbracket$  we contract the subgraph  $G_i$  in order to obtain the subgraph  $H_{\varphi(i)}$ .

Observe that after step 1., we obtain the graph  $\text{cycle}(\tau^-)$ , where  $\tau^-$  can be obtained from  $\tau$  by deleting elements of indices in  $\llbracket 1, q \rrbracket \setminus \{\varphi(i), i \in \llbracket 1, p \rrbracket\}$ . Intuitively, we contracted the graphs that do not appear in  $H$  and removed their attachment point from the cycle. Then we replace in step 2. every graph of  $\tau^-$  by its corresponding

contraction of  $\sigma$ . Therefore the graph obtained at the end is  $\text{cycle}(\sigma)$ , that is  $H$ , as required.

The cases (ii) and (iii) are very similar:  $H$  can be obtained from  $G$  by following the same operations as above.  $\square$

Let us restate [Lemma 3.88](#) before we give its proof.

**Lemma 3.88.** *Connected clique-cactus graphs are well-quasi-ordered by  $\leq_c$ .*

*Proof.* Let us assume by contradiction that  $(\mathcal{C}, \leq_c)$  is not a wqo. All decreasing sequences of this quasi-order are finite (as each contraction decreases the number of edges by one), therefore  $(\mathcal{C}, \leq_c)$  contains an infinite antichain. Let us consider an antichain  $\{A_i\}_{i \in \mathbb{N}}$  of  $(\mathcal{C}, \leq_c)$  with the property that, for every  $i \in \mathbb{N}$ ,  $A_i$  is a minimal element (with respect to  $\leq_c$ ) such that there is an infinite antichain starting with  $\{A_j\}_{j \in [0, i]}$ . Let  $\mathcal{B} = \bigcup_{i \in \mathbb{N}} \text{dec}(A_i)$  and let us show that  $(\mathcal{B}, \leq_c)$  is a wqo. For contradiction, let us assume that it is not a wqo and let  $\{B_i\}_{i \in \mathbb{N}}$  be an antichain of this quasi-order, with the same minimality requirement as above.

By definition of  $\mathcal{B}$ , for every  $H \in \mathcal{B}$  there is an integer  $i \in \mathbb{N}$  such that  $H \leq_c A_i$  (for instance, an integer  $i$  such that  $H \in \text{dec}(A_i)$ ). Therefore for every  $i \in \mathbb{N}$  there is an integer  $\pi(i)$  such that  $B_i \leq_c A_{\pi(i)}$ . Let  $k \in \mathbb{N}$  be the integer where  $\pi$  is minimum. Then the following sequence

$$\mathcal{A} = A_0, \dots, A_{\pi(k)-1}, B_k, B_{k+1}, \dots$$

is an infinite antichain of  $(\mathcal{C}, \leq_c)$ . Indeed, as both  $\{A_i\}_{i \in \mathbb{N}}$  and  $\{B_i\}_{i \in \mathbb{N}}$  are antichains, every pair of comparable graphs of  $\mathcal{A}$  involves one graph of  $\{A_i\}_{i \in [1, \pi(k)-1]}$  and one graph of  $\{B_i\}_{i \in \mathbb{N}_{\geq k}}$ . Let us assume that for some  $i \in [0, \pi(k) - 1]$  and  $j \in \mathbb{N}_{\geq k}$  we have  $A_i \leq B_j$ . Then  $A_i \leq B_j \leq A_{\pi(i)}$ , a contradiction with the fact that  $\{A_i\}_{i \in \mathbb{N}}$  is an antichain. The case  $B_j \leq A_i$  is not possible by the choice of  $k$ . This proves that  $(\mathcal{B}, \leq_c)$  is a wqo. According to [Theorem 3.1](#),  $(\mathcal{B}^*, \leq_c^*)$  is also a wqo. Let  $\mathcal{B}' = \{\text{stick}(\sigma), \sigma \in \mathcal{B}^*\}$ . Item (iii) of [Lemma 3.90](#) implies that any antichain in  $(\mathcal{B}', \leq_c)$  can be translated into an antichain of the same length in  $(\mathcal{B}^*, \leq_c^*)$ , hence  $(\mathcal{B}', \leq_c)$  is a wqo. By the same argument (now using items (i) and (ii) of [Lemma 3.90](#)), we deduce that the quasi-orders

$$(\{\text{cycle}(\sigma), \sigma \in \mathcal{B}'^*\}, \leq_c) \quad \text{and} \quad (\{\text{clique}(\sigma), \sigma \in \mathcal{B}'^*\}, \leq_c)$$

are well-quasi-orders. Therefore  $\mathcal{U} = \{\text{cycle}(\sigma), \sigma \in \mathcal{B}'^*\} \cup \{\text{clique}(\sigma), \sigma \in \mathcal{B}'^*\}$  is well-quasi-ordered by  $\leq_c$ , as a consequence of [Remark 3.25](#). According to [Observation 3.89](#), we have  $\{A_i\}_{i \in \mathbb{N}} \subseteq \mathcal{U}$ . This contradicts the fact that  $\{A_i\}_{i \in \mathbb{N}}$  is an infinite antichain. Therefore  $(\mathcal{C}, \leq_c)$  is a wqo and we are done.  $\square$

### 3.7.3 The dichotomy

The next observations will allow us to give the the proof of [Theorem 3.13](#), using the results obtained in the previous section. Recall that our goal is to characterize the graphs  $H$  such that  $(\text{Excl}_{\leq_c}(H), \leq_c)$  is a wqo.

*Observation 3.91.* For every  $p \in \mathbb{N}_{\geq 1}$ , contracting one edge in  $D_p$  gives either  $D_{p-1}$ , or  $K_{1,p}$ , depending on which edge is contracted.

To remind the definition of  $D_r$ , the reader may refer to [Figure 3.22](#). We want to identify graphs  $H$  such that  $(\text{Excl}_{\leq c.}(H), \leq c.)$  is a wqo, therefore we must consider every graph  $H$  such that  $\mathcal{A} \cap \text{Excl}_{\leq c.}(H)$  is finite, for every antichain  $\mathcal{A}$ . Let us set  $\mathcal{D} = \{D_r, r \in \mathbb{N}\}$  and  $\mathcal{S} = \{K_{1,r}, r \in \mathbb{N}\}$ . A first step towards this goal is the following observation.

**Lemma 3.92.** *Let  $p \in \mathbb{N}_{\geq 2}$ . If  $H \leq c. K_{2,p}$  then  $H \in \mathcal{D} \cup \mathcal{S}$ .*

*Proof.* Given that  $H \leq c. K_{2,p}$ , there is a sequence of contractions transforming  $K_{2,p}$  into  $H$ . If this sequence contains only one contraction, then it is straightforward that  $H = D_{p-1}$ . Therefore in the other cases  $H$  is a contraction of  $D_{p-1}$ . We get the result from [Observation 3.91](#) and the observation that every contraction of a graph of  $\mathcal{S}$  belongs to  $\mathcal{S}$ .  $\square$

*Observation 3.93.* For every positive integers  $p, q$  such that  $p < q$ , we have  $D_p \leq c. K_{2,q}$ .

Indeed, if  $F$  is a collection of  $q - p$  edges that are pairwise not incident with the same vertex of degree 2 of  $K_{2,p}$ , then it is easy to check that contracting  $F$  in  $K_{2,p}$  yields  $D_p$ . An immediate consequence of [Observation 3.93](#) is that  $\mathcal{A}_K \cap \text{Excl}_{\leq c.}(D_p)$  is finite for every positive integer  $p$ .

From the fact that every graph of  $\mathcal{D} \cup \mathcal{S}$  is a contraction of  $D_p$  for some positive integer  $p$ , [Observation 3.93](#) gives.

*Observation 3.94.* If  $(\text{Excl}_{\leq c.}(H), \leq)$  is a wqo, then  $H \leq c. D_p$  for some  $p \in \mathbb{N}_{\geq 1}$

However we will need an other antichain in order to find more properties that  $H$  must satisfy. Let us consider the antichain  $\mathcal{A}_{\overline{C}}$  of antiholes, that has been presented in [Subsection 3.5.1](#).

Again, we look at graphs  $H$  such that  $\text{Excl}_{\leq c.}(H) \cap \mathcal{A}_{\overline{C}}$  is finite. As a wqo must contain none of  $\mathcal{A}_K$  and  $\mathcal{A}_{\overline{C}}$ , it is enough to consider graphs such that  $\text{Excl}_{\leq c.}(H) \cap \mathcal{A}_{\overline{C}}$  is finite among those for which  $\text{Excl}_{\leq c.}(H) \cap \mathcal{A}_K$  is finite.

**Lemma 3.95.** *If  $p \geq 3$  then  $\text{Excl}_{\leq c.}(D_p) \cap \mathcal{A}_{\overline{C}}$  is infinite.*

*Proof.* For every  $p \geq 3$ , then graph  $D_p$  has independence number at least 3. Let  $q > p$ . As contracting edges can only decrease the independence number, there is no sequence of contractions transforming  $\overline{C}_q$  (which has independent number 2) to  $D_p$ , for every integer  $q > p$ . Therefore  $\overline{C}_q \in \text{Excl}_{\leq c.}(D_p)$ , for every integer  $q > p$ .  $\square$

**Corollary 3.96.** *If  $(\text{Excl}_{\leq c.}(H), \leq c.)$  is a wqo, then  $H \leq c. D_2$ .*

We are now ready to give the proof of [Theorem 3.13](#). We first restate it, as a reminder.

**Theorem 3.13** (Dichotomy for contractions). *Let  $H$  be a graph. The class of  $H$ -contraction-free simple graphs is wqo by  $\leq c.$  iff  $H \leq c. D_2$ .*

*Proof.* Let  $H$  be a graph such that  $\text{Excl}_{\leq c.}(H)$  is a wqo. Then  $H \leq c. D_2$ , by **Corollary 3.96**. On the other hand, if  $H \leq c. D_2$  then  $\text{Excl}_{\leq c.}(H) \subseteq \text{Excl}_{\leq c.}(D_2)$ . Observe that every antichain (respectively decreasing sequence) of  $(\text{Excl}_{\leq c.}(H), \leq c.)$  is an antichain (respectively a decreasing sequence) of  $(\text{Excl}_{\leq c.}(D_2), \leq c.)$ . As a consequence of **Lemma 3.88** we get that  $(\text{Excl}_{\leq c.}(H), \leq c.)$  is a wqo and we are done.  $\square$

### 3.7.4 Canonical antichains and contractions

In this section, we will use the following result of Ding in order to prove **Theorem 3.21**.

**Lemma 3.97** ([Din09]). *Let  $(S, \preceq)$  be a quasi-order, let  $\{A_i\}_{i \in \mathbb{N}}$  be a sequence of elements of  $S$  and let  $\{\mathcal{W}_i\}_{i \in \mathbb{N}}$  be a sequence of sequences of elements of  $S$ . If we have*

- (i)  $\{A_i\}_{i \in \mathbb{N}}$  is a fundamental infinite antichain; and
- (ii) for every  $i \in \mathbb{N}$ ,  $\mathcal{W}_i$  is a fundamental infinite antichain; and
- (iii) for every  $i \in \mathbb{N}$  and every  $H \in \mathcal{W}_i$ ,  $A_i \preceq H$ ,

then  $(S, \preceq)$  does not have a canonical antichain.

We will now define some sequences of graphs and show that they satisfy the properties of **Lemma 3.97**.

For every  $p, q \in \mathbb{N}$ , let  $W_{p,q}$  be the graph obtained by adding two non-adjacent dominating vertices to the disjoint union of  $\overline{K_p}$  and  $K_{2,q}$ . These two vertices are called *poles*, and the two vertices corresponding to the part of  $K_{2,q}$  of size 2 are called *semipoles*. Observe that the other vertices either have degree two (in which case they are adjacent to the two poles, only), or have degree four (and they are adjacent to both poles and both semipoles).

**Lemma 3.98.** *For every  $p, p', q, q' \in \mathbb{N}_{\geq 3}$ , there is no contraction model of  $W_{p,q}$  in  $W_{p',q'}$  if  $(p, q) \neq (p', q')$ .*

*Proof.* Let us assume that there is a contraction model  $\varphi$  of  $W_{p,q}$  in  $W_{p',q'}$ . Let  $u$  be a vertex of  $W_{p,q}$  of degree two. By definition of a contraction model, its image by  $\varphi$  must be a subset of degree 2. In  $W_{p',q'}$ , the connected subsets of degree 2 are either of the form  $\{v\}$ , or  $V(W_{p',q'}) \setminus \{v\}$ , where  $v \in V(W_{p',q'})$  has degree 2. As  $W_{p,q}$  has more than two vertices, the only possible form for  $\varphi(u)$  is  $\{v\}$  for some vertex  $v \in V(W_{p',q'})$  of degree 2. Therefore we have  $p \leq p'$ . The same argument applied to vertices of degree 4 yields  $q \leq q'$ . Let us now consider poles and semipoles.

Let  $u$  be a pole. Observe that according to the above remarks,  $\varphi(u)$  must be adjacent to vertices of degree two, so it should contain a pole of  $W_{p',q'}$ . If  $\varphi(u)$  contains in addition a vertex of degree 2 or 4 of  $W_{p',q'}$ , then  $\varphi(u)$  is dominating. This is not possible since  $u$  is not dominating, therefore  $\varphi(u) = \{v\}$  for some pole  $v$  of  $W_{p',q'}$ . Let us now assume that  $u$  is a semipole of  $W_{p,q}$ . As previously, the above remarks imply that  $\varphi(u)$  is adjacent to vertices of degree 4 of  $W_{p',q'}$ . Hence  $\varphi(u)$  contains a semipole of

$W_{p',q'}$  (it cannot contain a pole as both belong to the image of poles of  $W_{p,q}$ ). Therefore each semipole of  $W_{p,q}$  is sent to a subset of  $V(W_{p',q'})$  containing a semipole. Observe that  $\varphi(u)$  cannot contain a vertex of degree two otherwise it would not be connected. Besides, it cannot contain a vertex of degree 4 otherwise it would be adjacent to the image by  $\varphi$  of the other semipole of  $W_{p,q}$ . Consequently  $\varphi(u)$  contains a semipole of  $W_{p',q'}$  and no other vertex. We proved that for every  $u \in V(W_{p,q})$ , the set  $\varphi(u)$  is a singleton. Therefore  $|V(W_{p,q})| = |V(W_{p',q'})|$ . Given that  $p \leq p'$  and  $q \leq q'$  (as proved above), this is possible only if  $p = p'$  and  $q = q'$ . This concludes the proof.  $\square$

**Corollary 3.99.**  $\{W_{p,q}\}_{p,q \geq 3}$  is an antichain for  $\leq_c$ .

For every  $i \in \mathbb{N}_{\geq 3}$ , let  $\mathcal{W}_i = \{W_{i,q}\}_{q \in \mathbb{N}_{\geq 3}}$ .

**Lemma 3.100.** For every  $p, q \in \mathbb{N}_{\geq 3}$ ,  $K_{2,p+1} \leq_c W_{p,q}$ .

*Proof.* Let  $S$  be the set of vertices of degree 4 and semipoles of  $W_{p,q}$  (i.e. the vertices of the copy of  $K_{2,q}$  used in the construction of  $W_{p,q}$ ). These vertices induced a connected subgraph as every vertex of degree 4 is adjacent to both semipoles. It is not hard to see that contracting  $S$  to a single vertex yields  $K_{2,p+1}$ .  $\square$

*Observation 3.101.* Let  $p, q \in \mathbb{N}_{\geq 3}$ . There is no path containing four independent vertices in  $W_{p,q}$ .

**Corollary 3.102.** For every  $p, q \in \mathbb{N}_{\geq 3}$ , the graph  $W_{p,q}$  does not contain the gem as induced minor.

**Corollary 3.103.** No graph of  $\text{Incl}_{\leq_c}(\mathcal{W}_i)$  contains the gem as induced minor, for every  $i \in \mathbb{N}_{\geq 3}$ .

The following observation will allow us to use [Theorem 3.42](#), which deals with induced minors.

*Observation 3.104.* Let  $H$  and  $G$  be two graphs. If both of them have a dominating vertex, then  $H \leq_c G \iff H \leq_{\text{i.m.}} G$ .

The following corollary is a direct consequence of [Theorem 3.42](#), [Observation 3.104](#) and [Corollary 3.103](#).

**Corollary 3.105.** Graphs of  $\text{Incl}_{\leq_c}(\mathcal{W}_i)$  with a dominating vertex are wqo by  $\leq_c$ , for every  $i \in \mathbb{N}_{\geq 3}$ .

**Lemma 3.106.**  $\mathcal{W}_i$  is a fundamental antichain, for every  $i \in \mathbb{N}_{\geq 3}$ .

*Proof.* Let  $i \in \mathbb{N}_{\geq 3}$ . We need to show that  $(\text{Incl}_{\leq_c}(\mathcal{W}_i), \leq_c)$  is a wqo. Let us call *inner edge* every edge of  $W_{p,q}$  that is not incident with a pole, for every  $p, q \in \mathbb{N}_{\geq 3}$ . Observe that if a graph  $H \in \text{Incl}_{\leq_c}(\mathcal{W}_i)$  has been obtained by contracting at least one edge incident with a pole, then  $H$  has a dominating vertex. According to [Corollary 3.105](#), these graphs are wqo by  $\leq_c$ , therefore we will here consider graphs of  $\text{Incl}_{\leq_c}(\mathcal{W}_i)$  that have been obtained by only contracting inner edges. We call  $\mathcal{I}$  this class.

We first show that  $\mathcal{I}$  is the union of the two following classes:

- the class  $\mathcal{I}_0$  of graphs that can be obtained by adding two non-adjacent dominating vertices to  $\overline{K}_i + D_q$  for some  $q \in \mathbb{N}_{\geq 0}$ ; and
- the class  $\mathcal{I}_1$  of graphs that can be obtained by adding two non-adjacent dominating vertices to  $\overline{K}_i + K_{1,q}$  for some  $q \in \mathbb{N}_{\geq 0}$ .

Again we use the notion of *poles* to denote the two dominating vertices added to construct graphs of  $\mathcal{I}_0$  and  $\mathcal{I}_1$ . A *semipole* is either a dominating vertex of  $D_q$  (when dealing with graphs of  $\mathcal{I}_0$ ), or the dominating vertex of  $K_{1,q}$  (when dealing with graphs of  $\mathcal{I}_1$ ).

Contracting an inner edge in  $W_{i,q}$  clearly yields a graph of  $\mathcal{I}_0$ . Now, observe that any further contraction of an edge connecting a vertex of degree 4 to a semipole gives a graph of  $\mathcal{I}_0$  again. If, on the other hand, we contract the edge connecting the two semipoles, then we get a graph of  $\mathcal{I}_1$ . On a graph of  $\mathcal{I}_1$ , contracting an edge of the star (used in the construction of this graph) still gives a graph of  $\mathcal{I}_1$ . Therefore  $\mathcal{I} = \mathcal{I}_0 \cup \mathcal{I}_1$ .

Let us assume that  $\mathcal{I}$  is not wqo by  $\leq_c$ . Therefore it has an infinite antichain. As  $\mathcal{I} = \mathcal{I}_0 \cup \mathcal{I}_1$ , one of  $\mathcal{I}_0$  and  $\mathcal{I}_1$  (at least) has an infinite antichain. Let  $\mathcal{A}$  be such an infinite antichain.

We now look at vertices of graphs of  $\mathcal{A}$  that are neither poles, nor semipoles, nor have degree 2. These vertices are the vertices of degree 2 of the copy of  $D_q$  or the vertices of degree one of the copy of  $K_{1,q}$  used in the construction of the graphs of  $\mathcal{A}$  (depending whether  $\mathcal{A} \subseteq \mathcal{I}_0$  or  $\mathcal{A} \subseteq \mathcal{I}_1$ ). We call them *inner vertices*.

Let  $A$  and  $A'$  be two graphs of  $\mathcal{A}$  such that  $A$  has less inner vertices than  $A'$ . These graphs exist since the elements of  $\mathcal{A}$  are distinct. Let  $q$  be the number of inner vertices of  $A$  and  $q'$  the one of  $A'$ .

In both cases  $\mathcal{A} \subseteq \mathcal{I}_0$  and  $\mathcal{A} \subseteq \mathcal{I}_1$  we can obtain  $A$  from  $A'$  by contracting  $q' - q$  inner vertices of  $A'$  to a semipole. This contradicts the fact that  $\mathcal{A}$  is an antichain. Therefore  $(\mathcal{I}, \leq_c)$  is a wqo. This implies that  $\mathcal{W}_i$  is fundamental, as required.  $\square$

We are now ready to prove [Theorem 3.21](#), that we first restate.

**Theorem 3.21.** *Under the contraction relation, the class of all graphs does not have a canonical antichain.*

*Proof.* Let  $A_i = K_{2,i+1}$  for every  $i \in \mathbb{N}_{\geq 3}$ . According to [Lemma 3.106](#) and [Lemma 3.100](#), these sequences of graphs satisfy the requirements of [Lemma 3.97](#). Therefore there is no canonical antichain for the contraction relation.  $\square$

### 3.8 Multigraph contractions and well-quasi-ordering

In this section, we prove [Theorem 3.14](#) and [Theorem 3.22](#). Let us introduce some definitions and present the intermediate results that we use. First of all, as opposite to the previous sections of this chapter, we deal here with multigraphs: multiple edges are allowed, but not loops.

A *bond* is a minimal non-empty edge cut, i.e. a minimal set of edges whose removal increases the number of connected components (cf. [Figure 3.23](#)).




Figure 3.23: A bond of size 3 (dashed edges) in the house graph.

For every  $p, k \in \mathbb{N}$ , let  $\mathcal{G}_{p,k}$  be the class of graphs having at most  $p$  connected components and not containing a bond of order more than  $k$ . What we prove in this section is the following result.

**Theorem 3.107.** *For every  $p, k \in \mathbb{N}$ , the class  $\mathcal{G}_{p,k}$  is well-quasi-ordered by  $\leq_{\text{m.c.}}$ .*

It is easy to see that **Theorem 3.14** is a consequence of **Theorem 3.107**. Remark that a graph has a bond of order  $k$  iff it contains  $\theta_k$  as contraction, and that it has  $p$  connected components iff it can be contracted to  $\overline{K}_p$ . **Theorem 3.107** is proven in **Subsection 3.8.2** and results on canonical antichains appear in **Subsection 3.8.3**.

Let us now introduce the notation and definitions we will use in this section. As we have to handle many objects with several indices, we find more convenient to use the dot notation  $A.b$ , informally meaning “object  $b$  related to object  $A$ ”. For every  $i \in \{2, 3\}$  we denote by  $\mathcal{H}^{(i)}$  the class of all  $i$ -connected graphs in a class  $\mathcal{H}$ .

**2-rooted graphs.** We define a 2-rooted graph in a very similar way as a rooted graph is defined. A 2-rooted graph is a triple  $(G, r, s)$  where  $G$  is a graph and  $r$  and  $s$  are two distinct vertices of  $G$ . Given two 2-rooted graphs  $(G, r, s), (H, r', s')$ , we say that  $(H, r', s')$  is a contraction of  $(G, r, s)$ , what we denote by  $(H, r', s') \leq_{\text{m.c.}} (G, r, s)$ , if there is a contraction model  $\mu$  of  $H$  in  $G$  such that  $r' \in \mu(r)$  and  $s' \in \mu(s)$ . For the sake of simplicity, we sometimes denote by  $G$  the 2-rooted graph  $(G, r, s)$  and refer to its first (respectively second) root by  $G.r$  (respectively  $G.s$ ). For every rooted graph  $G$ , we define  $\text{root}(G) = \{G.r, G.s\}$ . A 2-rooted graph  $G$  is *edge-rooted* if  $\{G.r, G.s\} \in E(G)$ .




Figure 3.24: Attaching  $H$  to vertices  $(u, v)$  of  $G$  (roots are the white vertices).

The operation of *attaching* a 2-rooted graph  $H$  on the pair of vertices  $(u, v)$  of graph  $G$ , denoted  $G \oplus_u^v H$ , yields the graph rooted in  $(G.r, G.s)$  obtained by identifying  $u$

with  $H.r$  and  $v$  with  $H.s$  in the disjoint union of  $G$  and  $H$  (see [Figure 3.24](#) for an illustration). If both  $G$  and  $H$  are  $(\Sigma, \preceq)$ -labeled (for some qoset  $(\Sigma, \preceq)$ ), then the labeling function  $\lambda$  of the graph  $G \oplus_u^v H$  is defined as follows:

$$\lambda: \begin{cases} V(G \oplus_u^v H) & \rightarrow \mathcal{P}(\Sigma) \\ w & \mapsto G.\lambda(w) \quad \text{if } w \in V(G) \setminus \{u, v\} \\ w & \mapsto H.\lambda(w) \quad \text{if } w \in V(H) \setminus \{H.r, H.s\} \\ w & \mapsto G.\lambda(w) \cup H.\lambda(w) \quad \text{otherwise, i.e. when } w \in \{u, v\}. \end{cases}$$

Now we state several results that we will use. The first one is a decomposition theorem for 2-connected graphs by Tutte.

**Proposition 3.108** ([\[Tut61\]](#), see also [\[Die05, Exercise 20 of Chapter 12\]](#)). *Every 2-connected simple graph has a tree-decomposition  $(T, \mathcal{X})$  such that  $|X_t \cap X_{t'}| = 2$  for every edge  $\{t, t'\} \in T$  and all torsos are either 3-connected or a cycle.*

**Proposition 3.109** ([\[OOT93\]](#)). *For every  $k \in \mathbb{N}$  there is a positive integer  $\zeta_k$  such that every 3-connected simple graph of order at least  $\zeta_k$  contains a wheel of order  $k$  or a  $K_{3,k}$  as minor.*

### 3.8.1 Gluing graphs

This section is devoted to building larger wqos from smaller ones in classes of labeled graphs that are rooted by two vertices. Step by step, we will construct wqos that will be directly used in the proof of the main result, as in [Step 3](#) of the general scheme of [Section 3.4](#). Labels will be used to reduce the study of (unlabeled) graphs to the case of 2-connected graphs with labels (by the virtue of [Lemma 3.32](#)), whereas roots enable us to construct graphs using the operation  $\oplus$ . In this section, let  $(\Sigma, \preceq)$  be any qoset.

**Lemma 3.110.** *Let  $H, H', G, G'$  be four  $(\Sigma, \preceq)$ -labeled 2-rooted graphs. If  $H \leq_{\text{m.c.}} H'$  and  $G \leq_{\text{m.c.}}^\mu G'$ , then for every distinct  $u, v$  in  $V(G)$  and  $u' \in \mu(u), v' \in \mu(v)$  we have*

$$G \oplus_u^v H \leq_{\text{m.c.}} G' \oplus_{u'}^{v'} H'.$$

*Proof.* Let  $\mu_H: V(H) \rightarrow \mathcal{P}(V(H'))$  (respectively  $\mu_G: V(G) \rightarrow \mathcal{P}(V(G'))$ ) be a contraction model of  $H$  in  $H'$  (respectively of  $G$  in  $G'$ ). We consider the following function:

$$\nu: \begin{cases} V(G \oplus_u^v H) & \rightarrow \mathcal{P}(V(G' \oplus_{u'}^{v'} H')) \\ v & \mapsto \mu_H(v) & \text{if } v \in H \setminus \text{root}(H) \\ v & \mapsto \mu_G(v) & \text{if } v \in G \setminus \{u, v\} \\ v & \mapsto \mu_H(v) \cup \mu_G(v) & \text{otherwise.} \end{cases}$$

Let us check that  $\nu$  is a contraction model of  $G \oplus_u^v H$  in  $G' \oplus_{u'}^{v'} H'$ . First, observe that for every  $x \in V(G \oplus_u^v H)$ , the subgraph induced in  $G' \oplus_{u'}^{v'} H'$  by  $\nu(x)$  is connected ([M2](#)): either  $\nu(x) = \mu_H(x)$  or  $\nu(x) = \mu_G(x)$  (and in these cases it follows from the fact that  $\mu_H$  and  $\mu_G$  are models) or  $\nu(x) = \mu_H(x) \cup \mu_G(x)$  (if  $x \in \{u, v\}$ ) and  $(G' \oplus_{u'}^{v'} H')[\nu(x)]$  is connected because both  $\mu_H(x)$  and  $\mu_G(x)$  induce a connected subgraph and both

contain the root of  $H'$ . Furthermore, the images through  $\nu$  of two distinct vertices are always disjoint (M1), and every vertex of  $G' \oplus_u^{v'} H'$  belongs to the image of a vertex (M7), again because  $\mu_H$  and  $\mu_G$  are models. Let us now show point (M6). For every distinct  $x, y \in V(G \oplus_u^v H)$ ,

- either  $x, y \in V(H)$  and  $\{x, y\} \neq \text{root}(H)$  and then

$$\text{mult}_{G \oplus_u^v H}(x, y) = \sum_{(x', y') \in \nu(x) \times \nu(y)} \text{mult}_{G' \oplus_u^{v'} H'}(x', y')$$

as  $\mu_H$  is a contraction model (and symmetrically for the case  $x, y \in V(G)$  and  $\{x, y\} \neq \{u, v\}$ );

- or  $x \in V(H) \setminus \text{root}(H)$  and  $y \in V(G) \setminus \{u, v\}$ : there are no edges between  $x$  and  $y$  because every edge of  $G \oplus_u^v H$  is either an edge of  $H$  or an edge of  $G$ , neither between  $\nu(x)$  and  $\nu(y)$  since  $\nu(x) \subseteq V(H) \setminus \text{root}(H)$  and  $\nu(y) \subseteq V(G) \setminus \{u, v\}$ , therefore we get

$$\text{mult}_{G \oplus_u^v H}(x, y) = \sum_{(x', y') \in \nu(x) \times \nu(y)} \text{mult}_{G' \oplus_u^{v'} H'}(x', y') = 0;$$

- or  $\{x, y\} = \{u, v\} = \text{root}(H)$ :

$$\begin{aligned} \text{mult}_{G \oplus_u^v H}(x, y) &= \text{mult}_G(x, y) + \text{mult}_H(x, y) \quad (\text{by definition of } \oplus) \\ &= \sum_{(x', y') \in \mu_G(x) \times \mu_G(y)} \text{mult}_{G'}(x', y') \\ &\quad + \sum_{(x', y') \in \mu_H(x) \times \mu_H(y)} \text{mult}_{H'}(x', y') \\ &= \sum_{(x', y') \in \nu(x) \times \nu(y)} \text{mult}_{G' \oplus_u^{v'} H'}(x', y'). \end{aligned}$$

Besides, as a consequence that  $\mu_G$  is root-preserving,  $\nu$  also has this property. Last, let us check that  $\nu$  is label-preserving. Let  $x \in V(G \oplus_u^v H)$ . If  $x \notin \{u, v\}$ , then  $(G \oplus_u^v H).\lambda(x) = G.\lambda(x)$  or  $(G \oplus_u^v H).\lambda(x) = H.\lambda(x)$  (depending whether  $x \in V(G) \setminus \{u, v\}$  or  $x \in H \setminus \text{root}(H)$ ) and in these cases labels are preserved, since  $\mu_G$  and  $\mu_H$  are label-preserving. If  $x \in \{u, v\}$ , then, as  $\mu_G$  and  $\mu_H$  are label-preserving we have:

$$\begin{aligned} (G \oplus_u^v H).\lambda(x) &= G.\lambda(x) \cup H.\lambda(y) \\ &\leq^* \bigcup_{x' \in \mu_G(x)} G'.\lambda(x') \cup \bigcup_{x' \in \mu_H(x)} H'.\lambda(x') \\ &\leq^* \bigcup_{x' \in \nu(x)} (G' \oplus_u^{v'} H').\lambda(x') \end{aligned}$$

and thus  $\nu$  is label-preserving as well. We just proved that  $\nu$  is a contraction model of  $G \oplus_u^v H$  in  $G' \oplus_u^{v'} H'$ . Consequently,  $G \oplus_u^v H \leq_{\text{m.c.}} G' \oplus_u^{v'} H'$ , as desired.  $\square$

**Corollary 3.111.** Let  $l \in \mathbb{N}^*$ , let  $J$  be a  $(\Sigma, \preceq)$ -labeled 2-rooted graph and  $\langle (u_i, v_i) \rangle_{i \in \llbracket 1, l \rrbracket}$  be a sequence of pairs of distinct vertices of  $J$ . Let  $\mathcal{H}$  be a class of  $(\Sigma, \preceq)$ -labeled 2-rooted graphs,  $\langle G_1, \dots, G_l \rangle, \langle H_1, \dots, H_l \rangle \in \mathcal{H}^l$  and let  $G$  (respectively  $H$ ) be the graph constructed by attaching  $G_i$  (respectively  $H_i$ ) to the vertices  $(u_i, v_i)$  of  $J$ , for every  $i \in \llbracket 1, l \rrbracket$ . If  $\langle H_1, \dots, H_l \rangle \leq_{\text{m.c.}}^l \langle G_1, \dots, G_l \rangle$ , then  $H \leq_{\text{m.c.}} G$ .

*Proof.* By induction on  $l$ . The case  $l = 1$  follows from [Lemma 3.110](#). If  $l \geq 2$ , then, let  $G'$  (respectively  $H'$ ) be the graph constructed by attaching  $G_i$  (respectively  $H_i$ ) to the vertices  $(u_i, v_i)$  of  $J$ , for every  $i \in \llbracket 1, l-1 \rrbracket$ . By induction hypothesis, we have  $H' \leq_{\text{m.c.}} G'$ . Since  $H$  (respectively  $G$ ) is isomorphic to  $H' \oplus_{u_l}^{v_l} H_l$  (respectively  $G' \oplus_{u_l}^{v_l} G_l$ ) and  $H_l \leq_{\text{m.c.}} G_l$ , by [Lemma 3.110](#), we have  $H \leq_{\text{m.c.}} G$  as desired.  $\square$

**Lemma 3.112.** Let  $\mathcal{H}$  be a family of  $(\Sigma, \preceq)$ -labeled 2-rooted connected graphs, let  $J$  be a  $(\Sigma, \preceq)$ -labeled 2-rooted graph, and let  $\mathcal{H}_J$  be the class of  $(\Sigma, \preceq)$ -labeled 2-rooted graphs that can be constructed by attaching a graph  $H \in \mathcal{H}$  to  $(u, v)$  for every  $u, v \in V(J)$ . If  $(\mathcal{H}, \leq_{\text{m.c.}})$  is a wqo, then so is  $(\mathcal{H}_J, \leq_{\text{m.c.}})$ .

*Proof.* Let  $(u_1, v_1), \dots, (u_l, v_l)$  be an enumeration of all the pairs of distinct vertices of  $J$ . In this proof, we will design an epi that constructs graphs of  $\mathcal{H}_J$  from a tuple of  $l$  graphs of  $\mathcal{H}$ . Let  $f: (\mathcal{H}^l, \leq_{\text{m.c.}}^l) \rightarrow (\mathcal{H}_J, \leq_{\text{m.c.}})$  be the function that, given a tuple  $(H_1, \dots, H_l)$  of  $l$  graphs of  $\mathcal{H}$ , returns the graph constructed from  $J$  attaching  $H_i$  to  $(u_i, v_i)$  for every  $i \in \llbracket 1, l \rrbracket$ . This function is clearly surjective. Let us show that it is monotone.

Let  $(G_1, \dots, G_l), (H_1, \dots, H_l) \in \mathcal{H}^l$  be two tuples such that the following holds:

$$(H_1, \dots, H_l) \leq_{\text{m.c.}}^l (G_1, \dots, G_l).$$

According to [Remark 3.29](#), it is enough to deal with the cases where these two sequences differ only in one coordinate. Since all parameters of  $f$  play a similar role, we only look at the case where  $H_1 \leq_{\text{m.c.}} G_1$  and  $\forall i \in \llbracket 2, l \rrbracket, H_i = G_i$ . Let  $J'$  be the graph obtained from  $J$  by attaching  $G_i$  to  $(u_i, v_i)$ , for every  $i \in \llbracket 2, l \rrbracket$ . Remark that  $f(H_1, \dots, H_l)$  (respectively  $f(G_1, \dots, G_l)$ ) can be obtained by attaching  $H_1$  (respectively  $G_1$ ) to  $(u_1, v_1)$  in  $J'$ . By [Lemma 3.110](#) and since  $H_1 \leq_{\text{m.c.}} G_1$ , we have  $J \oplus_{u_1}^{v_1} H_1 \leq_{\text{m.c.}} J \oplus_{u_1}^{v_1} G_1$  and thus  $f(H_1, \dots, H_l) \leq_{\text{m.c.}} f(G_1, \dots, G_l)$ . Consequently,  $f$  is monotone and surjective:  $f$  is an epi. In order to show that  $\mathcal{H}_J$  is a wqo, it suffices to prove that the domain of  $f$  is a wqo (cf. [Remark 3.28](#)). As a finite Cartesian product of wqos,  $(\mathcal{H}^l, \leq_{\text{m.c.}}^l)$  is a wqo by [Lemma 3.26](#). This concludes the proof.  $\square$

**Lemma 3.113.** Let  $\mathcal{H}$  be a family of  $(\Sigma, \preceq)$ -labeled 2-rooted connected graphs and let  $\mathcal{H}_\circ$  be the class of  $(\Sigma, \preceq)$ -labeled graphs that can be constructed from a cycle by attaching a graph of  $\mathcal{H}$  to either  $(u, v)$  or  $(v, u)$  for every edge  $\{u, v\}$ , after deleting the edge  $\{u, v\}$ . If  $(\mathcal{H}, \leq_{\text{m.c.}})$  is a wqo, then so is  $(\mathcal{H}_\circ, \leq_{\text{m.c.}})$ .

*Proof.* Again, this proof relies on the property of epimorphisms to send wqos on wqos: we will present an epi that maps sequences of graphs of  $(\mathcal{H}, \leq_{\text{m.c.}})$  to graphs of  $(\mathcal{H}_\circ, \leq_{\text{m.c.}})$ . Let  $\mathcal{H}' = \mathcal{H} \cup \{(H, s, r), (H, r, s) \in \mathcal{H}\}$ , i.e.  $\mathcal{H}'$  contains graphs of  $\mathcal{H}$  with the roots possibly swapped. As the union of two wqos,  $(\mathcal{H}', \leq_{\text{m.c.}})$  is a wqo

**(Remark 3.25).** We consider the function  $f: (\mathcal{H}^*, \leq_{\text{m.c.}}^*) \rightarrow (\mathcal{H}_o, \leq_{\text{m.c.}})$  that, given a sequence  $\langle H_1, \dots, H_k \rangle$  of graphs of  $(\mathcal{H}', \leq_{\text{m.c.}})$  (for some integer  $k \geq 2$ ), returns the graph obtained from the cycle on vertices  $v_0, \dots, v_{k-1}$  (in this order) by deleting the edge  $\{v_i, v_{(i+1) \bmod k}\}$  and attaching  $H_i$  to  $(v_i, v_{(i+1) \bmod k})$ , for all  $i \in \llbracket 1, k \rrbracket$ . Observe that by definition of  $\mathcal{H}_o$  and  $\mathcal{H}'$ , the function  $f$  is surjective. We now show that  $f$  is monotone. Let  $G = \langle G_0, \dots, G_{k-1} \rangle$  and  $H = \langle H_0, \dots, H_{l-1} \rangle \in \mathcal{H}^*$  be two sequences such that  $G \leq_{\text{m.c.}}^* H$ . For the sake of readability, we will refer to the vertices of  $f(G)$  (respectively  $f(H)$ ) and to the graphs of  $G$  (respectively  $H$ ) by the same names. By definition of the relation  $\leq_{\text{m.c.}}^*$ , there is an increasing function  $\rho: \llbracket 0, k-1 \rrbracket \rightarrow \llbracket 0, l-1 \rrbracket$  such that for every  $i \in \llbracket 0, k-1 \rrbracket$ , we have  $G_i \leq_{\text{m.c.}} H_{\rho(i)}$ .

A crucial remark here is that since the graphs of  $\mathcal{H}'$  are connected, each of them can be contracted to an edge between its two roots. Therefore, for every graph  $H_i$  of the sequence  $H$  (for some  $i \in \llbracket 0, l-1 \rrbracket$ ) we can first contract  $H_i$  to an edge in  $f(H)$  and then contract this edge. That way we obtain a graph similar to  $f(H)$  except that  $H_i$  has been deleted and its roots merged: this is the graph  $f(\langle H_0, \dots, H_{i-1}, H_{i+1}, \dots, H_{l-1} \rangle)$ . By applying this operation on every subgraph of  $f(H)$  belonging to  $\{H_i, i \in \llbracket 1, l \rrbracket \setminus \rho(\llbracket 0, k \rrbracket)\}$ , we obtain the graph  $f(\langle H_{\rho(i)} \rangle_{i \in \llbracket 1, k \rrbracket})$  and we thus have  $f(\langle H_{\rho(i)} \rangle_{i \in \llbracket 1, k \rrbracket}) \leq_{\text{m.c.}} f(H)$ . Now, recall that the function  $\rho$  is such that for every  $i \in \llbracket 0, k-1 \rrbracket$ , we have  $G_i \leq_{\text{m.c.}} H_{\rho(i)}$ . Furthermore, the graphs  $f(G)$  and  $f(\langle H_{\rho(i)} \rangle_{i \in \llbracket 1, k \rrbracket})$  are both constructed by attaching graphs to the same graph (a cycle on  $k$  vertices). By **Corollary 3.111**, we therefore have  $f(G) \leq_{\text{m.c.}} f(\langle H_{\rho(i)} \rangle_{i \in \llbracket 1, k \rrbracket})$ , hence  $f(G) \leq_{\text{m.c.}} f(H)$  by transitivity of  $\leq_{\text{m.c.}}$ . We just proved that  $f$  is an epi. The domain of  $f$  is a wqo (as a set of finite sequences from a wqo, cf. **Theorem 3.1**), so its codomain  $(\mathcal{H}_o, \leq_{\text{m.c.}})$  is a wqo as well according to **Remark 3.28** and this concludes the proof.  $\square$

**Lemma 3.114.** Let  $k \in \mathbb{N}$  and let  $\mathcal{H}$  be a class of 2-rooted graphs, none of which having more than  $k$  edges between the two roots. Let  $\mathcal{H}^-$  be the class of graphs of  $\mathcal{H}$  where all edges between the two roots have been removed. If  $(\mathcal{H}, \leq_{\text{m.c.}})$  is a wqo, then so is  $(\mathcal{H}^-, \leq_{\text{m.c.}})$ .

*Proof.* Let us assume that  $(\mathcal{H}, \leq_{\text{m.c.}})$  is a wqo. For every  $i \in \llbracket 0, k \rrbracket$ , let  $\mathcal{H}_i$  be the subclass of graphs of  $\mathcal{H}$  having exactly  $i$  edges between the two roots. Each class  $\mathcal{H}_i$  ( $i \in \llbracket 0, k \rrbracket$ ) is a subclass of  $\mathcal{H}$  which is well-quasi-ordered by  $\leq_{\text{m.c.}}$ , therefore it is well-quasi-ordered by  $\leq_{\text{m.c.}}$  as well. Let  $f$  be the function that, given a 2-rooted graph  $G$ , returns a copy of  $G$  where all edges between the roots have been deleted. The rest of the proof draws upon the following remark.

**Remark 3.115.** Let  $G, H$  be two edge-rooted graphs where the edge between the roots has the same multiplicity. Then  $H \leq_{\text{m.c.}} G \Leftrightarrow f(H) \leq_{\text{m.c.}} f(G)$  (every model of  $H$  in  $G$  is also a contraction model of  $f(H)$  in  $f(G)$ , and vice-versa).

Let  $i \in \llbracket 0, k \rrbracket$ , let  $\mathcal{H}_i^- = \{f(H), H \in \mathcal{H}_i\}$ , and let  $\langle f(G_i) \rangle_{i \in \mathbb{N}}$  be an infinite sequence of  $\mathcal{H}_i^-$ . By an observation above,  $(\mathcal{H}_i, \leq_{\text{m.c.}})$  is a wqo, hence  $\langle G_i \rangle_{i \in \mathbb{N}}$  has a good pair  $(G_i, G_j)$  (with  $i, j \in \mathbb{N}$ ,  $i < j$ ). According to **Remark 3.115**,  $(f(G_i), f(G_j))$  is a good pair of  $\langle f(G_i) \rangle_{i \in \mathbb{N}}$ . Every infinite sequence of  $(\mathcal{H}_i^-, \leq_{\text{m.c.}})$  has a good pair, therefore this qoset is a wqo. Remark that  $(\mathcal{H}^-, \leq_{\text{m.c.}})$  is the union of the  $k+1$  wqos  $\{(\mathcal{H}_i^-, \leq_{\text{m.c.}})\}_{i \in \llbracket 0, k \rrbracket}$ , therefore it is a wqo as well (cf. **Remark 3.25**) and this concludes the proof.  $\square$

**Proposition 3.108** provides an interesting description of the structure of 2-connected simple graphs. The two following easy lemmas show that it can easily be adapted to multigraphs.

**Lemma 3.116.** *Let  $G$  be a graph and let  $G'$  be its underlying simple graph. The graph  $G$  is 2-connected iff  $G'$  is 2-connected or  $G = \theta_k$  for some integer  $k \geq 2$ .*

*Proof.* It is clear that  $G$  is 2-connected whenever  $G'$  is. Let us now assume that  $G$  is 2-connected but  $G'$  is not, and let  $u, v \in V(G')$  be two distinct vertices of  $G'$  such that there is no pair of internally disjoint paths from  $u$  to  $v$  in  $G'$ . Since  $G$  is 2-connected, there are two internally disjoint paths  $P$  and  $Q$  in  $G$  linking  $u$  to  $v$ . Remark that if  $P$  and  $Q$  are edge-disjoint, then the corresponding paths in  $G'$  are internally disjoint and link  $u$  to  $v$ , a contradiction with the choice of these two vertices. Therefore  $P$  and  $Q$  share an edge (which has multiplicity at least two). Since these paths are internally disjoint, their ends must be the ends of the edge that they share:  $\{u, v\}$  is an edge with multiplicity at least two. Removing (all multiplicities of) the edge  $\{u, v\}$  in  $G$  yields two connected components, one,  $G_u$ , containing  $u$  and the other,  $G_v$ , containing  $v$ . Since every path from vertices of  $G_u$  to vertices of  $G_v$  in  $G$  contains  $u$ , the graph  $G_u$  contains only the vertex  $u$  (otherwise  $G$  is not 2-connected) and by symmetry  $V(G_v) = \{v\}$ . Therefore  $G = \theta_k$ , for some integer  $k \geq 2$ , as required.  $\square$

**Lemma 3.117** (extension of **Proposition 3.108** to graphs). *Every 2-connected graph has a tree-decomposition  $(T, \mathcal{X})$  such that  $|X_t \cap X_{t'}| = 2$  for every edge  $\{t, t'\} \in T$  and where every torso is either 3-connected or a cycle.*

*Proof.* Let  $G$  be a 2-connected graph and  $G'$  be its underlying simple graph. If  $G'$  is 2-connected, then by **Proposition 3.108** it has a tree-decomposition  $(T, \mathcal{X})$  such that  $|X_t \cap X_{t'}| = 2$  for every edge  $\{t, t'\} \in T$  and where every torso is either 3-connected, or a cycle. Noticing that  $(T, \mathcal{X})$  is also a tree-decomposition of  $G$  concludes this case. If  $G'$  is not 2-connected, then by **Lemma 3.116** we have  $G = \theta_k$  for some integer  $k \geq 2$ . If  $k = 2$  the graph  $G$  is a cycle, and if  $k > 2$  it is 3-connected, therefore it has a trivial tree-decomposition with one bag, which satisfies the properties required in the statement of the lemma.  $\square$

We call a tree decomposition as in **Lemma 3.117** a *Tutte decomposition*.

### 3.8.2 Well-quasi-ordering graphs without big bonds

The main result is proved in three steps. First, we show that for every  $k \in \mathbb{N}$ , the class of labeled 2-connected graphs of  $\mathcal{G}_{1,k}$  is well-quasi-ordered by  $\leq_{\text{m.c.}}$ . Then, we use **Lemma 3.32** to extend this result to all graphs of  $\mathcal{G}_{1,k}$ , i.e. all connected graphs not containing a bond of size more than  $k$ . The result for disconnected graphs then follows by the application of **Lemma 3.34**.

**Lemma 3.118.** *For every  $k \in \mathbb{N}$ , and for every wqo  $(\Sigma, \preceq)$ , the qoset*

$$(\text{lab}_{(\Sigma, \preceq)}(\mathcal{G}_{1,k}^{(2)}), \leq_{\text{m.c.}})$$

*is a wqo.*

*Proof.* Let  $k \in \mathbb{N}$ , and let  $(\Sigma, \preceq)$  be a wqo. By contradiction, let us assume that  $(\text{lab}_{(\Sigma, \preceq)}(\mathcal{G}_{1,k}^{(2)}), \leq_{\text{m.c.}})$  is not a wqo. We consider the edge-rooted closure  $\mathcal{H}$  of  $\text{lab}_{(\Sigma, \preceq)}(\mathcal{G}_{1,k}^{(2)})$ , i.e. the class of all edge-rooted graphs whose underlying non-rooted graphs belongs to  $\text{lab}_{(\Sigma, \preceq)}(\mathcal{G}_{1,k}^{(2)})$ . Clearly,  $(\mathcal{H}, \leq_{\text{m.c.}})$  is not a wqo, as a consequence of our initial assumption. We will show that this leads to a contradiction.

Let  $\{A_i\}_{i \in \mathbb{N}}$  be an infinite minimal (wrt.  $\leq_{\text{m.c.}}$ ) bad sequence of  $(\mathcal{H}, \leq_{\text{m.c.}})$ : for every  $i \in \mathbb{N}$ ,  $A_i$  is a minimal graph (wrt.  $\leq_{\text{m.c.}}$ ) such that there is an infinite bad sequence starting with  $A_0, \dots, A_i$ . For every  $i \in \mathbb{N}$ ,  $A_i$  has a Tutte decomposition (**Lemma 3.117**) which has a bag containing the endpoints of the edge  $\{A_i.r, A_i.s\}$  (because it is a tree decomposition). Let  $A_i.X$  be the torso of some (arbitrarily chosen) bag in such a decomposition which contains  $A_i.r$  and  $A_i.s$ .

For every edge  $x, y \in V(A_i.X)$ , let  $A_i.V_{x,y}$  be the vertex set of the (unique) block which contains both  $x$  and  $y$  in the graph obtained from  $A_i$  by deleting vertices  $V(A_i.X) \setminus \{x, y\}$  and adding the edge  $\{x, y\}$  with multiplicity 2.

Let us consider graphs obtained by contracting all the edges of  $A_i$  that does not have both endpoints in  $A_i.V_{x,y}$  in a way such that  $A_i.r$  gets contracted to  $x$  and  $A_i.s$  gets contracted to  $y$ . Remark that for fixed  $i$  and  $(x, y)$ , these graphs differ only by the multiplicity of the edge between the two roots  $x$  and  $y$ . For every  $i \in \mathbb{N}$  and  $x, y \in V(A_i.X)$ , we denote by  $A_i.C_{x,y}$  an arbitrarily chosen such graph. Eventually, we set  $A_i.C = \{A_i.C_{x,y}, x, y \in V(A_i.X)\}$ . Remark that every graph of  $A_i.C$  belongs to  $\mathcal{G}_{1,k}^{(2)}$  and is a contraction of  $A_i$ .

*Claim 3.119.*  $\mathcal{C} = \cup_{i \in \mathbb{N}} A_i.C$  is wqo by  $\leq_{\text{m.c.}}$ .

*Proof.* By contradiction, assume that  $(\mathcal{C}, \leq_{\text{m.c.}})$  has an infinite bad sequence  $\{B_i\}_{i \in \mathbb{N}}$ . By definition of  $\mathcal{C}$ , for every  $i \in \mathbb{N}$  there is a  $j = \varphi(i) \in \mathbb{N}$  such that  $B_i \leq_{\text{m.c.}} A_j$ . Let  $i_0 \in \mathbb{N}$  be an integer with  $\varphi(i_0)$  minimum. Let us consider the following infinite sequence:

$$A_0, \dots, A_{\varphi(i_0)-1}, B_{i_0}, B_{i_0+1}, \dots$$

Remark that this sequence cannot have a good pair of the form  $A_i \leq_{\text{m.c.}} A_j$ ,  $0 \leq i < j < \varphi(i_0)$  (respectively  $B_i \leq_{\text{m.c.}} B_j$ ,  $i_0 \leq i < j$ ) since  $\{A_i\}_{i \in \mathbb{N}}$  (respectively  $\{B_i\}_{i \in \mathbb{N}}$ ) is an antichain. Let us assume that there is a good pair of the form  $A_i \leq_{\text{m.c.}} B_j$ , for some  $i \in \llbracket 0, \varphi(i_0) - 1 \rrbracket$ ,  $j \geq i_0$ . Then we have  $A_i \leq_{\text{m.c.}} B_j \leq_{\text{m.c.}} A_{\varphi(j)}$ . By the choice of  $i_0$  we have  $\varphi(i_0) \leq \varphi(j)$ , hence  $i < \varphi(j)$  so  $(A_i, A_{\varphi(j)})$  is a good pair of  $\{A_i\}_{i \in \mathbb{N}}$ , a contradiction. Therefore, this sequence is an infinite bad sequence of  $(\mathcal{H}, \leq_{\text{m.c.}})$  and we have  $B_{i_0} \leq_{\text{m.c.}} A_{\varphi(i_0)}$  and  $B_{i_0} \not\leq_{\text{m.c.}} A_{\varphi(i_0)}$ . This contradicts the minimality of  $\{A_i\}_{i \in \mathbb{N}}$ , therefore  $(\mathcal{C}, \leq_{\text{m.c.}})$  is a wqo.  $\square$

Let  $\mathcal{C}^-$  be the class of 2-rooted graphs obtained from graphs of  $\mathcal{C}$  by deleting the edge between the roots. We set  $\mathcal{C}^+ = \{H \oplus_{H.r}^{H.s} \theta_i, i \in \llbracket 0, k \rrbracket, H \in \mathcal{C}^-\}$ . In other words  $\mathcal{C}^+$  is the class of graphs that can be constructed by possibly replacing the edge at the root of a graph of  $\mathcal{C}$  by an edge of multiplicity  $i$ , for any  $i \in \llbracket 1, k \rrbracket$ .

*Remark 3.120.* It follows from **Lemma 3.114** that both  $(\mathcal{C}^-, \leq_{\text{m.c.}})$  and  $(\mathcal{C}^+, \leq_{\text{m.c.}})$  are wqos.

Notice that for every  $i \in \mathbb{N}$  and  $\{x, y\} \in E(A_i.X)$ , the graph  $A_i[A_i.V_{x,y}]$  rooted in  $(x, y)$  belongs to  $\mathcal{C}^+$ . As explained thereafter, this property enables us to see  $A_i$  as a graph built from graphs of  $\mathcal{C}^+$ .

According to [Lemma 3.117](#), for every  $i \in \mathbb{N}$ , the graph  $A_i.X$  (which is the torso of a bag of a Tutte decomposition) is either a 3-connected graph (and thus  $|V(A_i.X)| < \zeta_k$  by [Proposition 3.109](#)), or a cycle (of any length). Therefore we can partition  $\{A_i\}_{i \in \mathbb{N}}$  into at most  $\zeta_k$  subsequences depending on the *type* of  $A_i.X$ , where this type can be either “cycle” or one type for each possible value of  $|V(A_i.X)|$  when  $A_i.X$  is 3-connected. Let us show that each of these subsequences is finite.

*First case:*  $\{A_i\}_{i \in \mathbb{N}}$  has an infinite subsequence  $\{D_i\}_{i \in \mathbb{N}}$  such that for every  $i \in \mathbb{N}$ ,  $D_i.X$  is a cycle. Then each graph of  $\{D_i\}_{i \in \mathbb{N}}$  can be constructed by attaching a graph of the wqo  $(\mathcal{C}^+, \leq_{\text{m.c.}})$  to each edge of a cycle after deleting this edge. By [Lemma 3.113](#), these graphs are wqo by  $\leq_{\text{m.c.}}$ , a contradiction.

*Second case:* for some positive integer  $n < \zeta_k$ ,  $\{A_i\}_{i \in \mathbb{N}}$  has an infinite subsequence  $\{D_i\}_{i \in \mathbb{N}}$  such that for every  $i \in \mathbb{N}$ ,  $|V(D_i.X)| = n$ . Then every graph of  $\{D_i\}_{i \in \mathbb{N}}$  can be constructed by attaching a graph of the wqo  $(\mathcal{C}^+, \leq_{\text{m.c.}})$  to each pair of distinct vertices of  $\bar{K}_n$ . By [Lemma 3.112](#),  $\{D_i\}_{i \in \mathbb{N}}$  has a good pair, which is contradictory since it is a bad sequence.

We just proved that  $\{A_i\}_{i \in \mathbb{N}}$  can be partitioned into a finite number of subsequences each of which is finite. Hence  $\{A_i\}_{i \in \mathbb{N}}$  is finite as well, a contradiction. Therefore our initial assumption is false and  $(\text{lab}_{(\Sigma, \preceq)}(\mathcal{G}_{1,k}^{(2)}), \leq_{\text{m.c.}})$  is a wqo.  $\square$

**Corollary 3.121.** *For every  $k \in \mathbb{N}$ , the class  $\mathcal{G}_{1,k}$  is well-quasi-ordered by  $\leq_{\text{m.c.}}$ .*

*Proof.* According to [Lemma 3.118](#), for every wqo  $(\Sigma, \preceq)$ , the class of  $\Sigma$ -labeled 2-connected graphs of  $\mathcal{G}$  are wqo by  $\leq_{\text{m.c.}}$ . By [Lemma 3.32](#), this implies that  $(\mathcal{G}_{1,k}, \leq_{\text{m.c.}})$  is a wqo and we are done.  $\square$

The proof of [Theorem 3.107](#) now follows from the combination of [Corollary 3.121](#) and [Lemma 3.34](#).

### 3.8.3 Canonical antichains and multigraph contractions

This section is devoted to the proof of the two results related to canonical antichains of  $\leq_{\text{m.c.}}$ .

*Remark 3.122.* Every canonical antichain of  $\leq_{\text{m.c.}}$  is infinite.

Recall that  $\mathcal{A}_\theta$  is the class of all connected multigraphs on two vertices, and that  $\mathcal{A}_{\bar{K}}$  is the class of all edgeless graphs, as depicted on [Figure 3.25](#).

Let us remind the statement of [Theorem 3.22](#) before we prove it.

**Theorem 3.22.** *Every antichain  $\mathcal{A}$  of  $\leq_{\text{m.c.}}$  is canonical iff each of the following sets are finite:*

$$\mathcal{A}_\theta \setminus \mathcal{A}; \quad \mathcal{A}_{\bar{K}} \setminus \mathcal{A}; \quad \text{and} \quad \mathcal{A} \setminus \{\mathcal{A}_\theta \cup \mathcal{A}_{\bar{K}}\}.$$




Figure 3.25: Two infinite antichains for multigraph contractions: multiedges and edgeless graphs.

*Proof.* “ $\Rightarrow$ ”: Let  $\mathcal{A}$  be a canonical antichain of  $\leq_{\text{m.c.}}$  and let us assume for contradiction that  $\mathcal{B} = \mathcal{A}_\theta \setminus \mathcal{A}$  (respectively  $\mathcal{B} = \mathcal{A}_{\overline{K}} \setminus \mathcal{A}$ ) is infinite. Let  $\mathcal{B}^+$  be the closure of  $\mathcal{B}$  and remark that  $\mathcal{B}^+ = \mathcal{B} \cup \{K_1\}$  (respectively  $\mathcal{B}^+ = \mathcal{B}$ ). Then the contraction downwards-closed class  $\mathcal{B}^+$  has finite intersection with  $\mathcal{A}$  whereas it contains the infinite antichain  $\mathcal{B}$ . This is a contradiction with the fact that  $\mathcal{A}$  is canonical, hence both  $\mathcal{A}_\theta \setminus \mathcal{A}$  and  $\mathcal{A}_{\overline{K}} \setminus \mathcal{A}$  are finite.

Let us now assume that  $\mathcal{C} = \mathcal{A} \setminus \{\mathcal{A}_\theta \cup \mathcal{A}_{\overline{K}}\}$  is infinite and let  $\mathcal{C}^+$  be the closure of  $\mathcal{C}$ . Being a subset of an antichain,  $\mathcal{C}$  is an antichain as well and consequently  $\mathcal{C}^+$  is a contraction downwards-closed class that is not well-quasi-ordered. By [Theorem 3.14](#),  $\mathcal{C}^+$  contains infinitely many elements of  $\mathcal{A}_\theta \cup \mathcal{A}_{\overline{K}}$ . Notice that besides being infinite,  $\mathcal{C}^+ \cap (\mathcal{A}_\theta \cup \mathcal{A}_{\overline{K}})$  is also disjoint from  $\mathcal{A} \cap (\mathcal{A}_\theta \cup \mathcal{A}_{\overline{K}})$ , otherwise  $\mathcal{A}$  would contain an element from  $\mathcal{C}$  contractible to an element of  $\mathcal{A} \cap (\mathcal{A}_\theta \cup \mathcal{A}_{\overline{K}})$ . But then one of  $\mathcal{A}_\theta \setminus \mathcal{A}$  and  $\mathcal{A}_{\overline{K}} \setminus \mathcal{A}$  is infinite, a contradiction with our previous conclusion. Therefore  $\mathcal{C}$  is finite.

“ $\Leftarrow$ ”: Let  $\mathcal{A}$  be an antichain such that each of  $\mathcal{A}_\theta \setminus \mathcal{A}$ ,  $\mathcal{A}_{\overline{K}} \setminus \mathcal{A}$ , and  $\mathcal{A} \setminus \{\mathcal{A}_\theta \cup \mathcal{A}_{\overline{K}}\}$  is finite, and let us show that  $\mathcal{A}$  is canonical. Let  $\mathcal{F}$  be a contraction downwards-closed class of graphs. If  $\mathcal{F} \cap \mathcal{A}$  is infinite, then  $\mathcal{F}$  trivially contains the infinite antichain  $\mathcal{F} \cap \mathcal{A}$ . On the other hand, if  $\mathcal{F} \cap \mathcal{A}$  is finite then by [Theorem 3.14](#) the class  $\mathcal{F}$  is well-quasi-ordered, hence by definition it does not contain an infinite antichain. Consequently,  $\mathcal{A}$  is canonical, as required.  $\square$

We can now prove [Corollary 3.23](#), that has already been stated in the introduction of this chapter.

**Corollary 3.23.** *Every canonical antichain of  $\leq_{\text{m.c.}}$  is fundamental<sup>5</sup>.*

*Proof.* Let  $\mathcal{A}$  be a canonical antichain of  $\leq_{\text{m.c.}}$ . Observe that we have the following:

$$\text{Incl}_{\leq_{\text{m.c.}}}(\mathcal{A}) = \text{Incl}_{\leq_{\text{m.c.}}}(\mathcal{A} \cap \mathcal{A}_\theta) \cup \text{Incl}_{\leq_{\text{m.c.}}}(\mathcal{A} \cap \mathcal{A}_{\overline{K}}) \cup \text{Incl}_{\leq_{\text{m.c.}}}(\mathcal{A} \setminus (\mathcal{A}_\theta \cup \mathcal{A}_{\overline{K}})).$$

Now, it is easy to notice that:

<sup>5</sup>The definition of the concept of *fundamental antichain* is given in [Section 3.2](#).

- $\text{Incl}_{\leq \text{m.c.}}(\mathcal{A} \cap \mathcal{A}_\theta) \subseteq \text{Incl}_{\leq \text{m.c.}}(\mathcal{A}_\theta) = \{K_1\}$ ;
- $\text{Incl}_{\leq \text{m.c.}}(\mathcal{A} \cap \mathcal{A}_{\overline{K}}) \subseteq \text{Incl}_{\leq \text{m.c.}}(\mathcal{A}_{\overline{K}}) = \emptyset$ ;
- $\text{Incl}_{\leq \text{m.c.}}(\mathcal{A} \setminus (\mathcal{A}_\theta \cup \mathcal{A}_{\overline{K}}))$  is finite, because  $\mathcal{A} \setminus (\mathcal{A}_\theta \cup \mathcal{A}_{\overline{K}})$  is finite by **Theorem 3.22** and since  $\mathcal{A}$  is canonical.

Therefore,  $\text{Incl}_{\leq \text{m.c.}}(\mathcal{A})$  is finite as well and hence cannot contain an infinite antichain; this proves that  $\mathcal{A}$  is fundamental.  $\square$

# CHAPTER 4

---

## EXCLUSION THEOREMS

One of the most celebrated results from the Graph Minors series of Robertson and Seymour is the following result, also known as the *Grid Exclusion Theorem*.

**Theorem 4.1** (Grid Exclusion Theorem, [RS86]). *There exists a function  $f: \mathbb{N} \rightarrow \mathbb{N}$  such that, for every integer  $h$ , every graph that does not contain a minor isomorphic to the  $h \times h$ -grid has treewidth at most  $f(h)$ .*

This result is an *exclusion theorem*: it relates a graph parameter (in this case the treewidth) with the absence of some pattern as a substructure. In this chapter, we present exclusion theorems related to the parameters of girth (Section 4.1), treewidth (Section 4.3), maximum degree (Section 4.2), and tree-cut width (Section 4.4).

Beside their combinatorial value, exclusion theorems has proven useful in order to obtain Erdős–Pósa-type results. This aspect will be illustrated in Chapter 5.

This chapter contains material that previously appeared in the following articles:

- [RT15] *Low polynomial exclusion of planar graph patterns*, co-authored with Dimitrios M. Thilikos, Journal of Graph Theory (2015);
- [RST16] *An edge variant of the Erdős–Pósa property*, co-authored with Ignasi Sau and Dimitrios M. Thilikos, Discrete Mathematics (2016);
- [CRST15c] *Minors in graphs of large  $\theta_r$ -girth*, co-authored with Dimitris Chatzidimitriou, Ignasi Sau, and Dimitrios M. Thilikos, submitted (2015);
- [GKRT16] *Packing and covering immersion models of planar subcubic graphs*, co-authored with Archontia Giannopoulou, O-joung Kwon, and Dimitrios M. Thilikos, WG 2016.

### 4.1 Clique-majors in graphs of large $\theta_r$ -girth

In this section, we introduce the concept of  $\theta_r$ -girth of a graph and show that graphs of sufficiently large minimum degree contain as a minor cliques whose order is an

exponential function of their girth, extending a result of [KO03]. We also show that the minimum degree can be replaced by some connectivity measurement.

### 4.1.1 The quest for large clique-majors

A classic result in graph theory asserts that if a graph has minimum degree  $ck\sqrt{\log k}$ , then it can be transformed to a complete graph of at least  $k$  vertices by applying edge contractions (i.e., it contains a  $k$ -clique minor). This result has been proved by Kostochka in [Kos84] and Thomason in [Tho83a] and a precise estimation of the constant  $c$  has been given by Thomason in [Tho01a]. For recent results related to conditions that force a clique minor see [Mar04, JW13, DHJ<sup>+</sup>13, FKO09, KO04a].

Recall that the *girth* of a graph  $G$  is the minimum length of a cycle in  $G$ . Interestingly, graphs of large minimum degree contain clique-minors whose order is an *exponential* function of their girth. In particular, it follows by the main result of Kühn and Osthus in [KO03] that there is a constant  $c$  such that, if a graph has minimum degree  $d \geq 3$  and girth  $z$ , then it contains as a minor a clique of size  $k$ , where

$$k \geq \frac{d^{cz}}{\sqrt{z \cdot \log d}}.$$

In this section we provide conditions, alternative to the above one, that can force the existence of a clique-minor whose size is exponential.

***H*-girth.** Given two graphs  $G$  and  $H$ , we define the *H-girth* of  $G$  as the minimum number of edges of an  $H$ -major in  $G$ . If  $G$  does not contain  $H$  as a minor, we will say that its  $H$ -girth is equal to infinity. For every  $r \in \mathbb{N}$ , recall that  $\theta_r$  denotes the graph with two vertices and  $r$  parallel edges. Clearly, the girth of a graph is its  $\theta_2$ -girth and, for every  $r_1 \leq r_2$ , the  $\theta_{r_1}$ -girth of a graph is at most its  $\theta_{r_2}$ -girth.

Our first result is the following extension of the result of Kühn and Osthus in [KO03] for the case of  $\theta_r$ -girth.

**Theorem 4.2.** *There is a constant  $c > 0$  such that, for every  $r \geq 2$ ,  $d \geq 3r$ , and  $z \geq r$ , if a graph has minimum degree  $d$  and  $\theta_r$ -girth at least  $z$ , then it contains as a minor a clique of size  $k$ , where*

$$k \geq \frac{\left(\frac{d}{r}\right)^{\frac{cz}{r}}}{\sqrt{\frac{z}{r} \cdot \log d}}.$$

In the formula above, a lower bound to the minimum degree as a function of  $r$  is necessary. An easy computation shows that when applying [Theorem 4.2](#) for  $r = 2$ , we can get the aforementioned formula of Kühn and Osthus, where the constant in the exponent is one fourth of the constant of [Theorem 4.2](#).

Our second finding is that this degree condition can be replaced by some “loose connectivity” requirement.

**Loose connectivity.** For two integers  $\alpha, \beta \in \mathbb{N}$ , a graph  $G$  is called  $(\alpha, \beta)$ -loosely connected if for every  $A, B \subseteq V(G)$  such that  $V(G) = A \cup B$  and  $G$  has no edge between  $A \setminus B$  and  $B \setminus A$ , we have that  $|A \cap B| < \beta \Rightarrow \min(|A \setminus B|, |B \setminus A|) \leq \alpha$ . Intuitively, this means that a *small* separator (i.e., one of less than  $\beta$  vertices) cannot “split” the graph into two *large* parts (that is, with more than  $\alpha$  vertices each).

Our second result indicates that the requirement on the minimum degree in [Theorem 4.2](#) can be replaced by the loose connectivity condition as follows.

**Theorem 4.3.** *There is a constant  $c > 0$  such that, for every  $r \geq 2$ ,  $\alpha \geq 1$ , and  $z \geq 168 \cdot \alpha \cdot r \log r$ , it holds that if a graph has more than  $(\alpha + 1) \cdot (2r - 1)$  vertices, is  $(\alpha, 2r - 1)$ -loosely connected, and has  $\theta_r$ -girth at least  $z$ , then it contains as a minor a clique of size  $k$  where*

$$k \geq \frac{2^{c \cdot \frac{z}{r\alpha}}}{\sqrt{rz}}.$$

Both [Theorem 4.2](#) and [Theorem 4.3](#) are derived from two more general results, namely [Theorem 4.5](#) and [Theorem 4.4](#), respectively. [Theorem 4.5](#) asserts that graphs with large  $\theta_r$ -girth and sufficiently large minimum degree contain as a minor a graph whose minimum degree is exponential in the girth. [Theorem 4.4](#) replaces the minimum degree condition with the absence of sufficiently large “edge-protrusions”, that are roughly tree-like structured subgraphs with small boundary to the rest of the graph (see [Subsection 4.1.2](#) for the detailed definitions). While these theorems are more general, we present [Theorem 4.2](#) and [Theorem 4.3](#) as the main results of this section as they are easier to apply.

As an application of these results, we will in [Subsection 4.3.6](#) optimally improve (up to a constant factor) the upper-bound on the treewidth of graphs excluding  $k \cdot \theta_r$  as a minor.

**Organization of the section.** The main notions used in this section are defined in [Subsection 4.1.2](#). Then, we show in [Subsection 4.1.3](#) that the proofs of [Theorem 4.2](#) and [Theorem 4.3](#) can be derived from [Theorem 4.5](#) and [Theorem 4.4](#), which are proved in [Subsection 4.1.4](#).

## 4.1.2 Definitions specific to this section

In this section, when giving the running time of an algorithm involving some graph  $G$ , we agree that  $n = |V(G)|$  and  $m = |E(G)|$ .

In order to decompose graphs along edge cuts, we introduce the following edge-counterpart of the notion of (*vertex-*)*protrusion* used in [[BFL+09a](#), [BFL+09b](#)] (among others). A subset  $Y \subseteq V(G)$  is a  $t$ -*edge-protrusion* of  $G$  with *extension*  $w$  (for some positive integer  $w$ ) if the graph  $G[Y \cup N_G(Y)]$  has a rooted tree-partition  $\mathcal{D} = (T, s, \mathcal{X})$  of width at most  $t$  and such that  $N_G(Y) = X_s$  and  $|V(T)| \geq w$ . The protrusion  $Y$  is said to be *connected* whenever  $Y \cup N_G(Y)$  induces a connected subgraph in  $G$ .

We recall that rooted tree-partitions, that are mentioned in the following definition, are defined in [Section 2.5](#) ([Definition 2.6](#)).

**Distance-decompositions.** A *distance-decomposition* of a connected graph  $G$  is a rooted tree-partition  $\mathcal{D} = (T, s, \mathcal{X})$  of  $G$ , where the following additional requirements are met (see also [YBdFT99]):

- (i)  $X_s$  contains only one vertex, we shall call it  $u$ , referred to as the *origin* of  $\mathcal{D}$ ;
- (ii) for every  $t \in V(T)$  and every  $x \in X_t$ ,  $\text{dist}_G(x, u) = \text{dist}_T(t, s)$ ;
- (iii) for every  $t \in V(T)$ , the graph  $G_t = G \left[ \bigcup_{t' \in \text{desc}(T, s)(t)} X_{t'} \right]$  is connected; and
- (iv) if  $C$  is the set of children of a vertex  $t \in V(T)$ , then the graphs  $\{G_{t'}\}_{t' \in C}$  are the connected components of  $G_t \setminus X_t$ .

Observe that a distance decomposition is uniquely determined by its origin. An example of distance-decomposition is given in Figure 4.1. For every vertex  $u$  of a graph on  $m$  edges, a distance-decomposition  $(T, s, \mathcal{X})$  with origin  $u$  can be constructed in  $O(m)$  steps by breadth-first search.




Figure 4.1: A graph (left) and a distance-decomposition with origin  $u_5$  of it (right).

Let  $P$  be a path in  $G$  and  $\mathcal{D} = (T, s, \mathcal{X})$  a distance-decomposition of  $P$ . We say that  $P$  is a *straight path* if the heights, in  $(T, s)$ , of the indices of the bags in  $\mathcal{D}$  that contain vertices of  $P$  are pairwise distinct. Obviously, in that case, the sequence of the heights of the bags that contain each subsequent vertex of the path is strictly monotone.

Let us introduce the notion of grouped partition, that extends the construction used in the proof of [Die05, Lemma 7.2.3] to our setting.

**Grouped partitions.** Let  $G$  be a connected graph and let  $d \in \mathbb{N}$ . A *d-grouped partition* of  $G$  is a partition  $\mathcal{R} = \{R_1, \dots, R_l\}$  of  $V(G)$  (for some positive integer  $l$ ) such that for each  $i \in \{1, \dots, l\}$ , the graph  $G[R_i]$  is connected and there is a vertex  $s_i \in R_i$  with the following properties:

- (i)  $\text{ecc}_{G[R_i]}(s_i) \leq 2d$  and

- (ii) for each edge  $e = \{x, y\} \in E(G)$  where  $x \in R_i$  and  $y \in R_j$  for some distinct integers  $i, j \in \{1, \dots, l\}$ , it holds that  $\text{dist}_G(x, s_i) \geq d$  and  $\text{dist}_G(y, s_j) \geq d$ .

A set  $S = \{s_1, \dots, s_l\}$  as above is a *set of centers* of  $\mathcal{R}$  where  $s_i$  is the *center* of  $R_i$  for  $i \in \{1, \dots, l\}$ .

Given a graph  $G$ , we define a  $d$ -*scattered set*  $W$  of  $G$  as follows:

- $W \subseteq V(G)$  and
- $\forall u, v \in W, \text{dist}_G(u, v) > d$ .

If  $W$  is inclusion-maximal, it will be called a *maximal  $d$ -scattered set* of  $G$ .

**Frontiers and ports.** Let  $G$  be a graph, let  $\mathcal{R} = \{R_1, \dots, R_l\}$  be a  $d$ -grouped partition of  $G$ , and let  $S = \{s_1, \dots, s_l\}$  be a set of centers of  $\mathcal{R}$ . For every  $i \in \llbracket 1, l \rrbracket$ , we denote by  $\mathcal{D}_i = (T_i, s_i, \mathcal{X}_i)$  the unique distance-decomposition with origin  $s_i$  of the graph  $G[R_i]$  where  $\mathcal{X}_i = \{X_t^i\}_{t \in V(T_i)}$ . For every  $i \in \llbracket 1, l \rrbracket$  and every  $h \in \llbracket 0, \text{ecc}_{T_i}(s_i) \rrbracket$ , we denote by  $I_i^h$  the vertices of  $(T_i, s_i)$  that are at distance  $h$  from  $s_i$  and we set  $I_i^{<h} = \bigcup_{h'=0}^{h-1} I_{h'}^i$  and  $I_i^{\geq h} = \bigcup_{h'=h}^{\text{ecc}_{T_i}(s_i)} I_{h'}^i$ . We also set

$$V_i^h = \bigcup_{t \in I_i^h} X_t^i, \quad V_i^{<h} = \bigcup_{t \in I_i^{<h}} X_t^i, \quad \text{and} \quad V_i^{\geq h} = \bigcup_{t \in I_i^{\geq h}} X_t^i.$$

The *vertex-frontier*  $F_i$  of  $R_i$  is the set of vertices in  $V_i^{d-1}$  that are connected in  $G$  to a vertex  $x \in V(G) \setminus R_i$  via a path, the internal vertices of which belong to  $V_i^{\geq d}$ . The *node-frontier* of  $T_i$  is

$$N_i = \{t \in V(T_i) \mid F_i \cap X_t^i \neq \emptyset\}. \quad (4.1)$$

A vertex  $t \in I_i^{\geq d-1}$  is called a *port* of  $T_i$  if  $X_t^i$  contains some vertex that is adjacent in  $G$  to a vertex of  $V(G) \setminus R_i$ .

### 4.1.3 Finding small $\theta_r$ -majors

#### Two intermediate results

In this section we show how [Theorem 4.2](#) and [Theorem 4.3](#) can be proved using the two following intermediate results, that will be proved in a subsequent section.

**Theorem 4.4.** *There exists an algorithm that, with input three positive integers  $r, w, z$  and an  $n$ -vertex graph  $G$ , outputs one of the following:*

- a  $\theta_r$ -major in  $G$  with at most  $z$  edges,
- a connected  $(2r - 2)$ -edge-protrusion  $Y$  of  $G$  with extension more than  $w$ , or
- an  $H$ -major in  $G$  for some graph  $H$  where  $\delta(H) \geq \frac{1}{r-1} 2^{\frac{z-5r}{4r(2w+1)}}$ ,

in  $O_r(m)$  steps.

**Theorem 4.5.** *There exists an algorithm that, with input three integers  $r, \delta, z$ , where  $r \geq 2$ ,  $\delta \geq 3r$ , and  $z \geq r$  and an  $n$ -vertex graph  $G$ , outputs one of the following:*

- a  $\theta_r$ -major in  $G$  with at most  $z$  edges,
- a vertex  $v$  of  $G$  of degree less than  $\delta$ , or
- an  $H$ -major in  $G$  for some graph  $H$  where  $\delta(H) \geq \frac{\delta-2r+3}{r-1} \cdot \lfloor \frac{\delta}{r-1} - 1 \rfloor^{\frac{z-r}{4r}}$ ,

in  $O_r(m)$  steps.

The results of Chandran and Subramanian in [CS05] imply that if  $G$  has girth at least  $z$  and minimum degree at least  $\delta$ , then  $\text{tw}(G) \geq \delta^{c \cdot z}$ , for some constant  $c$ . As in the third condition of **Theorem 4.5** it holds that  $\text{tw}(G) \geq \text{tw}(H) \geq \delta(H)$ , **Theorem 4.5** can also be seen as a qualitative extension of the results of [CS05].

The above two results will be used to prove **Theorem 4.2** and **Theorem 4.3**. We will also need the following result of Kostochka [Kos84].

**Proposition 4.6** ([Kos84], see also [Tho83a, Tho01a]). *There exists a constant  $\xi \in \mathbb{R}$  such that for every  $d \in \mathbb{N}$ , every graph of average degree at least  $d$  contains a clique of order  $k$  as a minor, for some integer  $k$  satisfying*

$$k \geq \xi \cdot \frac{d}{\sqrt{\log d}}.$$

Let us restate **Theorem 4.2** before proving it.

**Theorem 4.2.** *There is a constant  $c > 0$  such that, for every  $r \geq 2$ ,  $d \geq 3r$ , and  $z \geq r$ , if a graph has minimum degree  $d$  and  $\theta_r$ -girth at least  $z$ , then it contains as a minor a clique of size  $k$ , where*

$$k \geq \frac{\left(\frac{d}{r}\right)^{\frac{cz}{r}}}{\sqrt{\frac{z}{r} \cdot \log d}}.$$

*Proof.* Observe that since  $G$  has no  $\theta_r$ -major with at most  $z$  edges and  $G$  has minimum degree  $\delta \geq 3r$ , a call to the algorithm of **Theorem 4.5** on  $(r, \delta, z, G)$  should return an  $H$ -major in  $G$ , for some graph  $H$  where  $\delta(H) \geq \frac{\delta-2r+3}{r-1} \cdot \lfloor \frac{\delta}{r-1} - 1 \rfloor^{\frac{z-r}{4r}} =: d$ . Using the fact that  $z - r \geq z/2$ , it is not hard to check that there is a positive constant  $c' \in \mathbb{R}$  such that

$$\xi \cdot \frac{d}{\sqrt{\log d}} \geq \frac{\left(\frac{\delta}{r}\right)^{c' \cdot \frac{z}{r}}}{\sqrt{\frac{z}{r} \cdot \log \delta}}.$$

Hence by **Proposition 4.6**,  $G$  has a clique of the desired order as a minor.  $\square$

We also recall **Theorem 4.3** before providing its proof.

**Theorem 4.3.** *There is a constant  $c > 0$  such that, for every  $r \geq 2$ ,  $\alpha \geq 1$ , and  $z \geq 168 \cdot \alpha \cdot r \log r$ , it holds that if a graph has more than  $(\alpha + 1) \cdot (2r - 1)$  vertices, is  $(\alpha, 2r - 1)$ -loosely connected, and has  $\theta_r$ -girth at least  $z$ , then it contains as a minor a clique of size  $k$  where*

$$k \geq \frac{2^{c \cdot \frac{z}{r\alpha}}}{\sqrt{rz}}.$$

*Proof.* As in the proof of [Theorem 4.2](#), the properties that  $G$  enjoys will force a minor of large minimum degree. Let us call the algorithm of [Theorem 4.4](#) on  $(r, 3\alpha, z, G)$ . We assumed that  $G$  has no  $\theta_r$ -major with  $z$  edges or less, hence the output of the algorithm cannot be such a major. Let us now assume that the algorithm outputs a  $(2r - 2)$ -edge-protrusion  $Y$  of extension more than  $3\alpha$ , and let  $(T, s, \mathcal{X})$  be a rooted tree-partition of  $Y$  of width at most  $2r - 2$  such that  $N_G(Y) = X_s$  and  $|V(T)| > 3\alpha$ . It is known that every tree of order  $n$  has a vertex, the removal of which partitions the tree into components of size at most  $n/2$  each. Hence, there is a vertex  $v \in V(T)$  and a partition  $(Z, Z')$  of  $V(T) \setminus \{v\}$  such that:

- both  $Z \cup \{v\}$  and  $Z' \cup \{v\}$  induce connected subtrees of  $T$ ;
- $\frac{1}{3}|V(T)| \leq |Z|, |Z'| \leq \frac{2}{3}|V(T)|$ ; and
- $X_s \subseteq Z$  or  $v = s$ .

Let  $A = Z' \cup \{X_v\}$  and  $B = V(G) \setminus Z'$ . Notice that  $V(G) = A \cup B$  and that no edge of  $G$  lies between  $A$  and  $B$ . As  $A \cap B = X_v$ , we have  $|A \cap B| < 2r - 1$ . Last,  $Z' \subseteq A \setminus B$  and  $Z \subseteq B \setminus A$  give that  $|A \setminus B|, |B \setminus A| \geq \alpha$ . The existence of  $A$  and  $B$  contradicts the fact that  $G$  is  $(\alpha, 2r - 1)$ -loosely connected. Thus  $G$  has no  $(2r - 2)$ -edge-protrusion  $Y$  of extension more than  $3\alpha$ .

A consequence of this observation is that the only possible output of the algorithm mentioned above is an  $H$ -major of  $G$  for some graph  $H$ , where

$$\delta(H) \geq \frac{1}{r-1} \cdot 2^{\frac{z-5r}{4r(6\alpha+1)}} \geq \frac{1}{r} \cdot 2^{\frac{z}{168 \cdot r\alpha}} =: d.$$

Notice also that  $\log d = \frac{z}{168 \cdot r\alpha}$  which, by the condition of the theorem, is a non-negative number. Moreover,  $\log d \leq z/r$ . Therefore, there is there is a constant  $c'' \in \mathbb{R}$  such that

$$\xi \cdot \frac{d}{\sqrt{\log d}} \geq \frac{2^{c'' \cdot \frac{z}{r\alpha}}}{\sqrt{z \cdot r}}$$

in order to conclude the proof. □

#### 4.1.4 Structural results on graphs with large $\theta_r$ -girth

This section is devoted to the proofs of [Theorem 4.4](#) and [Theorem 4.5](#).

## Preliminary results

Before proving [Theorem 4.4](#) and [Theorem 4.5](#) we need some preliminary results. Let us start with some definitions.

Let  $(T, s)$  be a rooted tree and let  $N$  be a subset of its leaves. We say that a vertex  $u$  of  $T$  is  $N$ -critical if either it belongs to  $N \cup \{s\}$  or there are at least two vertices in  $N$  that are descendants of two distinct children of  $u$ . An  $N$ -unimportant path in  $T$  is a path with at least 2 vertices, with exactly two  $N$ -critical vertices, which are its endpoints (see [Figure 4.2](#) for a picture). As we see in the proof of the following lemma,  $N$ -unimportant paths are the maximal paths with internal vertices of degree 2 that appear if we repeatedly delete leaves that do not belong to  $N \cup \{s\}$ . Notice that an  $N$ -unimportant path in  $T$  cannot have an internal vertex that belongs to some other  $N$ -unimportant path. Also, among the two endpoints of an  $N$ -unimportant path there is always one which is a descendant of the other.




Figure 4.2: An unimportant path (dashed) in a tree. Gray subtrees are those without vertices from  $N$ .

**Lemma 4.7.** *Let  $d, k \in \mathbb{N}$ ,  $k \geq 1$ . Let  $(T, s)$  be a rooted tree and let  $N$  be a set of leaves of  $(T, s)$ , each of which is at distance at least  $d$  from  $s$ . If for some integer  $k$ , every  $N$ -unimportant path in  $T$  has length at most  $k$ , then  $|N| \geq 2^{d/k}$ .*

*Proof.* We consider the subtree  $T'$  of  $T$  obtained by repeatedly deleting leaves that do not belong to  $N \cup \{s\}$ . By construction, every leaf of  $(T', s)$  belongs to  $N$ , hence our goal is then to show that  $(T', s)$  has many leaves. Notice that in  $(T', s)$ , every vertex of degree at least 3 is  $N$ -critical. Therefore, the  $N$ -unimportant paths of  $(T', s)$  are the maximal paths, the internal vertices of which have degree two. By contracting each of these paths into an edge, we obtain a tree  $T''$  where every internal vertex has degree at least 3. Observe that every edge on a root-leaf path of  $T''$  is originated from the contraction of a path on at most  $k$  edges, as we assume that every  $N$ -unimportant path in  $T$  has length at most  $k$ . We deduce that  $T''$  has height at least  $d/k$ , hence it has at least  $2^{d/k}$  leaves. Consequently,  $T'$  has at least  $2^{d/k}$  leaves, and then  $|N| \geq 2^{d/k}$ .  $\square$

If  $(T, s, \mathcal{X})$  is a distance-decomposition of a graph and  $t \in V(T) \setminus \{s\}$ , we call  $E^{(t)}$  the set of edges that have one endpoint in  $X_t$  and the other in  $X_{p(t)}$ .

**Lemma 4.8.** *Let  $G$  be an  $n$ -vertex graph, let  $r$  be a positive integer, let  $\mathcal{D} = (T, s, \mathcal{X})$  be a distance-decomposition of  $G$ , and let  $d > 1$  be the height of  $(T, s)$ . Then either  $G$  contains a  $\theta_r$ -major with at most  $2 \cdot r \cdot d$  edges or for every vertex  $i \in V(T) \setminus s$ , it holds that  $|E^{(i)}| \leq r - 1$ . Moreover there exists an algorithm that, in  $O_r(m)$  steps, either finds such a major, or asserts that  $|E^{(i)}| \leq r - 1$  for every  $i \in V(T) \setminus s$ .*

*Proof.* We consider the non-trivial case where  $r \geq 2$ . Suppose that there exists a node  $t$  of  $(T, s)$  such that  $|E^{(t)}| \geq r$ . Clearly, such a  $t$  can be found in  $O(m)$  steps. We will prove that  $G$  contains a  $\theta_r$ -major. Let  $k$  be the height of  $t$  in  $T$ .

We need first the following claim.

**Claim 4.9.** Given a non-empty proper subset  $U$  of  $X_t$ , we can find in  $G_t$  a path of length at most  $2k$  from a vertex of  $U$  to a vertex of  $X_t \setminus U$ , in  $O(m)$  steps.

*Proof of Claim 4.9.* We can compute a shortest path  $P$  from a vertex of  $U$  to a vertex of  $X_t \setminus U$ , in  $O(m)$  steps using a BFS. Let us show that  $P$  has length at most  $2k$ . Let  $u \in U$  and  $v \in X_t \setminus U$  be the endpoints of  $P$ , and let  $w$  be a vertex of  $P$  of the lowest possible height  $h$  ( $0 \leq h \leq k$ ). Then it holds that  $\text{dist}_{G_t}(v, u) = \text{dist}_{G_t}(U, v)$ . We examine the non-trivial case where  $P$  has more than one edge. By minimality of  $P$  we have  $w \notin X_t$ .

Our next step is to prove that if  $P$  has more than one edge, then both the subpaths of  $P$  from  $u$  to  $w$  and from  $v$  to  $w$  are straight. Suppose now, without loss of generality, that the subpath from  $u$  to  $w$  is not straight and let  $z$  be the first vertex of it (starting from  $u$ ) which is contained in a bag of height greater than or equal to the height of the bag of its predecessor in  $P$ . By definition of a distance-decomposition (in particular items (ii) and (iii)), there is at least one vertex  $x \in X_t$  which is connected by a straight path  $P'$  to  $z$  in  $G$ . Then there are two possibilities:

- either  $x \in U$ , and then the union of the path  $P'$  and the portion of  $P$  between  $z$  and  $v$  is a path that is shorter than  $P$ ;
- or  $x \in X_t \setminus U$ , and in this case the union of the path  $P'$  and the portion of  $P$  between  $u$  and  $z$  is a path that is shorter than  $P$ .

As, in both cases, the occurring paths contradict the construction of  $P$ , we conclude that both the subpath of  $P$  from  $u$  to  $w$  and the one from  $v$  to  $w$  are straight. This implies that  $P$  has length at most  $2 \cdot (k - h) \leq 2 \cdot k$  and the claim follows.  $\diamond$

Our next step is to construct a vertex set  $U$  and a set of paths  $\mathcal{P}$  as follows. We set  $\mathcal{P} = \emptyset$ ,  $U = \emptyset$ , and we start by adding in  $U$  an arbitrarily chosen vertex  $u \in X_t$ . Using the procedure of Claim 4.9, we repeatedly find a path from a vertex of  $U$  to a vertex of  $X_t \setminus U$ , add this second vertex to  $U$  and the path to  $\mathcal{P}$ , until there are at least  $r$  edges in  $E^{(t)}$  that have endpoints in  $U$ .

The construction of  $U$  requires at most  $r$  repetitions of the procedure of Claim 4.9, and therefore  $O(r \cdot m)$  steps in total. Clearly  $|U| \leq r$ , hence  $|\mathcal{P}| \leq r - 1$ . Besides,

every path in  $\mathcal{P}$  has length at most  $2k$  according to [Claim 4.9](#). Notice now that  $\mathbf{UP}$  is a connected subgraph of  $G_t$  with at most  $2k \cdot (r - 1)$  edges.

As there are at least  $r$  edges in  $E^{(t)}$  with endpoints in  $U$  we may consider a subset  $F$  of them where  $|F| = r$ . Since  $\mathcal{D}$  is a distance-decomposition (by item (ii) of the definition), each edge  $e \in F$  is connected to the origin by a path of length  $d - k - 1$  whose edges do not belong to  $G_t$ . Let  $\mathcal{P}'$  be the collection of these paths. Clearly, the paths in  $\mathcal{P}'$  contain, in total, at most  $r \cdot (d - k - 1)$  edges.

If we now contract in  $G$  all edges in  $\mathcal{P}$  and all edges in  $\mathcal{P}'$ , except those in  $F$ , and then remove all edges not in  $F$ , we obtain a graph isomorphic to  $\theta_r$ . Therefore we found in  $G$  a  $\theta_r$ -major with at most

$$\begin{aligned} r \cdot (d - k - 1) + 2 \cdot k \cdot (r - 1) + r &\leq r \cdot (d - k - 1) + 2 \cdot k \cdot r + r \\ &= r \cdot (d + k) \\ &\leq 2 \cdot r \cdot d \end{aligned} \quad (\text{since } d \geq k)$$

edges in  $O(r \cdot m)$  steps. □

The following result is a direct consequence of [Lemma 4.8](#) and item (ii) of the definition of a distance-decomposition.

**Corollary 4.10.** *Let  $G$  be an  $n$ -vertex graph, let  $r$  be a positive integer, let  $\mathcal{D} = (T, s, \mathcal{X})$  be a distance-decomposition of  $G$ , and let  $d > 1$  be the height of  $(T, s)$ . If some bag of  $\mathcal{D}$  contains at least  $r$  vertices, then  $G$  contains a  $\theta_r$ -major with at most  $2 \cdot r \cdot d$  edges, which can be found in  $O_r(m)$  steps. □*

The remaining lemmata will be related to [grouped partitions](#).

**Lemma 4.11.** *For every positive integer  $d$  and every connected graph  $G$  there is a  $d$ -grouped partition of  $G$  that can be constructed in  $O(m)$  steps.*

*Proof.* If  $\text{diam}(G) \leq 2d$ , then  $\{V(G)\}$  is a  $d$ -grouped partition of  $G$ . Otherwise, let  $R = \{s_1, \dots, s_l\}$  be a maximal  $2d$ -scattered set in  $G$ . This set can be constructed in  $O(m)$  steps by breadth-first search. The sets  $\{R_i\}_{i \in \llbracket 1, l \rrbracket}$  are constructed by the following procedure:

1. Set  $k = 0$  and  $R_i^0 = \{s_i\}$  for every  $i \in \llbracket 1, l \rrbracket$ ;
2. For every  $i \in \llbracket 1, l \rrbracket$ , every  $v \in R_i^k$  and every  $u \in N_G(v)$ , if  $u$  has not been considered so far, add  $u$  to  $R_i^{k+1}$ ;
3. If  $k < 2d$ , increment  $k$  by 1 and go to step 2;
4. Let  $R_i = \bigcup_{k=0}^{2d} R_i^k$  for every  $i \in \llbracket 1, l \rrbracket$ .

Let  $\mathcal{R} = \{R_i\}_{i \in \llbracket 1, l \rrbracket}$ . By construction, each set  $R_i$  induces a connected graph in  $G$ . It remains to prove that  $\mathcal{R}$  is a partition of  $V(G)$  and that it has the desired properties.

Notice that in the above construction if a vertex is assigned to the set  $R_i$ , then it is not assigned to  $R_j$ , for every distinct integers  $i, j \in \llbracket 1, l \rrbracket$ . Let  $v \in V(G)$  be a vertex that does not belong to  $R_i$  for any  $i \in \llbracket 1, l \rrbracket$  after the procedure is completed. Then for every  $i \in \llbracket 1, l \rrbracket$  we have  $\text{dist}_G(v, s_i) > 2d$  and  $v \notin R$ , which contradicts the maximality of  $R$ . Therefore  $\mathcal{R}$  is a partition of  $V(G)$ .

Since for each vertex  $v$  in  $R_i$  it holds that  $\text{dist}_G(v, s_i) \leq 2d$ ,  $\mathcal{R}$  obviously satisfies property (i) of the definition.

For property (ii) of the definition, let  $e = \{x, y\}$  be an edge in  $G$  such that  $x \in R_i$ ,  $y \in R_j$ , for some distinct integers  $i, j \in \llbracket 1, l \rrbracket$ . Towards a contradiction, we assume without loss of generality that  $\text{dist}_G(x, s_i) < d$ . This means that during the construction of  $R_i$ , the vertex  $x$  was added to the set  $R_i^k$  for some  $k \leq d - 1$ . Also, since the vertex  $y$  is adjacent to  $x$  but was added to  $R_j^l$  for some  $l \leq 2d$  instead of  $R_i^{k+1}$ , it follows that  $l \leq k + 1$ , which means that  $\text{dist}_G(y, s_j) \leq k + 1$ . Hence  $\text{dist}_G(s_i, s_j) \leq \text{dist}_G(s_i, x) + \text{dist}_G(x, y) + \text{dist}_G(y, s_j) \leq k + 1 + k + 1 \leq 2d$  again is not possible since  $R$  is a  $2d$ -scattered set.

Finally, in the procedure above, each edge of the graph is encountered at most once, hence the whole algorithm will take at most  $O(m)$  time. This concludes the proof of the lemma.  $\square$

**Lemma 4.12.** *Let  $G$  be a graph, let  $\mathcal{R} = \{R_1, \dots, R_l\}$  be a  $d$ -grouped partition of  $G$ , and let  $s_i$  be a center of  $R_i$ , for every  $i \in \llbracket 1, l \rrbracket$ . If for some distinct  $i, j \in \llbracket 1, l \rrbracket$ ,  $G$  has at least  $r$  edges from vertices in  $R_i$  to vertices in  $R_j$  then  $G[R_i \cup R_j]$  contains a  $\theta_r$ -major with at most  $4 \cdot r \cdot d + r$  edges, which can be found in  $O_r(m)$  steps.*

*Proof.* Suppose that for some  $i \in \llbracket 1, l \rrbracket$ ,  $G$  has a set  $F$  of at least  $r$  edges from vertices in  $R_i$  to vertices in  $R_j$ . Let  $R'_i \subseteq R_i$  and  $R'_j \subseteq R_j$  be the sets of the endpoints of those edges. Since  $\mathcal{R}$  is a  $d$ -grouped partition of  $G$ , it holds that, for each  $x \in R'_i$  and  $y \in R'_j$ ,  $\text{dist}_G(x, s_i) \leq 2d$  and  $\text{dist}_G(y, s_j) \leq 2d$ . That directly implies that for every  $h \in \{i, j\}$ , there is a collection  $\mathcal{P}_h$  of  $r$  paths, each of length at most  $2d$  and not necessarily disjoint, in  $G[R_h]$  connecting  $s_h$  with each vertex in  $R'_h$ , which we can find in  $O_r(m)$  steps. It is now easy to observe that the graph  $Q$ , obtained from  $\mathbf{UP}_i \cup \mathbf{UP}_j$  by adding all edges of  $F$ , is the union of  $r$  paths between  $s_i$  and  $s_j$ , each containing at most  $4 \cdot d + 1$  edges. Therefore,  $Q$  is a major of  $\theta_r$  with at most  $4 \cdot r \cdot d + r$  edges, as required. As mentioned earlier the construction of  $\mathcal{P}_i$  and  $\mathcal{P}_j$  takes  $O_r(m)$  steps.  $\square$

**Lemma 4.13.** *Let  $r, d \in \mathbb{N}$ , let  $G$  be a graph with no  $\theta_r$ -major on  $2rd$  edges, let  $\mathcal{R} = \{R_1, \dots, R_l\}$  be a  $d$ -grouped partition of  $G$ , and let  $S = \{s_1, \dots, s_l\}$  be a set of centers of  $\mathcal{R}$ . For every  $i \in \llbracket 1, l \rrbracket$ , let  $\mathcal{D}_i = (T_i, r_i, \mathcal{X}_i)$  be the distance-decomposition with origin  $s_i$  of the graph  $G[R_i]$ .*

*If for some  $i \in \llbracket 1, l \rrbracket$  and  $w \in \mathbb{N}$ , the tree  $T_i$ , with node-frontier  $N_i$ , has an  $N_i$ -unimportant path of length at least  $2(w + 1)$ , then  $G$  has a connected  $(2r - 2)$ -edge-protrusion  $Y$  with extension more than  $w$ , which can be constructed in  $O_r(m)$  steps.*

*Proof.* Let  $P = t_0 \dots t_p$  be a  $N_i$ -unimportant path of length  $p \geq 2(w + 1)$  in  $T_i$ . We assume without loss of generality that  $t_p \in \text{desc}_{(T_i, r_i)}(t_0)$ . Due to the definition of distance-decompositions, the vertices in  $X_{t_0}^i$  or  $X_{t_p}^i$  form a vertex-separator of  $G$ . Let

$Z \subseteq E(G)$  be the set containing all edges between  $X_{t_0}^i$  and  $X_{t_1}^i$  and all edges between  $X_{t_{p-1}}^i$  and  $X_{t_p}^i$  in  $G$ . Let  $T'_i$  be the subtree of  $T_i$  that we obtain if we remove the descendants of  $t_p$  and any vertex that is not a descendant of  $t_1$ . Notice that  $Z$  is an edge-separator of  $G$ : it separates the vertices in bags of  $T'_i$  from the rest of the graph. Furthermore,  $Z$  has at most  $2r - 2$  edges as two bags of  $\mathcal{D}_i$  are never connected by more than  $r - 1$  vertices, according to [Lemma 4.8](#) applied to the fact that  $G$  has no  $\theta_r$ -major on  $2rd$  edges or less. Let  $Y = \bigcup_{t \in V(T'_i) \setminus \{t_0, t_p\}} X_t^i$ . In other words,  $Y$  consists of the vertices in the bags of  $T'_i$  excluding  $X_{t_0}^i$  and  $X_{t_p}^i$ . Obviously,  $N_G(Y) = X_{t_0} \cup X_{t_p}$ .

We will now construct a rooted tree-partition  $\mathcal{F} = (T_{\mathcal{F}}, r_{\mathcal{F}}, \mathcal{X}_{\mathcal{F}})$  of  $G[Y \cup N_G(Y)]$  of width at most  $2r - 2$  and such that  $|V(T_{\mathcal{F}})| > w$ . Let  $T_{\mathcal{F}}$  be the tree obtained from  $T'_i$  by identifying, for every  $j \in \llbracket 0, \lfloor (p-1)/2 \rfloor \rrbracket$ , the vertex  $t_j$  with the vertex  $t_{p-j}$ . If multiple edges are created during this identification, we replace them with simple ones. We also delete loops that may be created. Let us define the elements of  $\mathcal{X}_{\mathcal{F}} = \{X_t^{\mathcal{F}}\}_{t \in V(T_{\mathcal{F}})}$  as follows. If  $t \in V(T_{\mathcal{F}})$  is the result of the identification of  $t_j$  and  $t_{p-j}$  for some  $j \in \llbracket 0, \lfloor (p-1)/2 \rfloor \rrbracket$ , then we set  $X_t^{\mathcal{F}} = X_{t_j} \cup X_{t_{p-j}}$ . On the other hand, if  $t \in V(T_{\mathcal{F}})$  is a vertex of  $T'_i$  that has not been identified with some other vertex, then  $X_t^{\mathcal{F}} = X_t$ . The construction of  $\mathcal{F}$  is completed by setting  $r_{\mathcal{F}}$  to be the result of the identification of  $t_0$  and  $t_p$ , the endpoints of  $P$ .

It is easy to verify that  $\mathcal{F}$  is a rooted tree-partition of  $G[Y \cup N_G(Y)]$  of width at most  $2r - 2$ . Notice also that the identification of the antipodal vertices of the path  $P$  creates a path in  $T_{\mathcal{F}}$  of length  $\lfloor (p-1)/2 \rfloor$ . This implies that the extension of  $\mathcal{F}$  is at least  $\lfloor (p-1)/2 \rfloor \geq w + 1$ . Besides, all the operations performed to construct  $\mathcal{F}$  can be implemented in  $O_r(m)$  steps. This completes the proof.  $\square$

We continue with two easy lemmata related to **ports and frontiers**.

**Lemma 4.14.** *Let  $G$  be a graph, let  $\mathcal{R} = \{R_1, \dots, R_l\}$  be a  $d$ -grouped partition of  $G$ , and let  $S = \{s_1, \dots, s_l\}$  be a set of centers of  $\mathcal{R}$ . For every  $i \in \llbracket 1, l \rrbracket$ , let  $\mathcal{D}_i = (T_i, r_i, \mathcal{X}_i)$  be the distance-decomposition with origin  $s_i$  of the graph  $G[R_i]$ , and let  $N_i$  be the node-frontier of  $T_i$ . Then, for every  $i \in \llbracket 1, l \rrbracket$ , there are at least  $|N_i|$  ports in  $T_i$ .*

*Proof.* Let  $i \in \llbracket 1, l \rrbracket$ . We will show that every vertex in the node-frontier of  $T_i$  has a descendant which is a port. For every vertex  $t \in N_i \subseteq V(T_i)$ , there is, by definition, a path from  $t$  to a vertex in  $G \setminus R_i$ , the internal vertices of which belong to  $V_i^{\geq d}$ . Let  $v$  be the last vertex of this path (starting from  $t$ ) which belongs to  $R_i$  and let  $t' \in V(T)$  be the vertex such that  $v \in X_{t'}^i$ . Then  $t'$  is a port of  $T_i$ . Observe that  $t'$  cannot be the descendant of any other vertex of  $N_i$ . Therefore there are at least  $|N_i|$  ports in  $T_i$ .  $\square$

**Corollary 4.15.** *Let  $G$  be a graph, let  $\mathcal{R} = \{R_1, \dots, R_l\}$  be a  $d$ -grouped partition of  $G$ , and let  $S = \{s_1, \dots, s_l\}$  be a set of centers of  $\mathcal{R}$ . For every  $i \in \llbracket 1, l \rrbracket$ , let  $\mathcal{D}_i = (T_i, r_i, \mathcal{X}_i)$  be the distance-decomposition with origin  $s_i$  of the graph  $G[R_i]$ , and let  $N_i$  be the node-frontier of  $T_i$ . If for some integer  $k$ , every  $N_i$ -unimportant path in  $T_i$  has length at most  $k$ , then  $T_i$  contains at least  $2^{d/k}$  ports.*  $\square$

*Proof.* Let  $i \in \llbracket 1, l \rrbracket$ . From [Lemma 4.14](#), it is enough to prove that  $|N_i| \geq 2^{d/k}$ . Then the result follows by applying [Lemma 4.7](#) for  $(T_i, s_i)$ ,  $d$ ,  $N_i$ , and  $k$ .  $\square$

### Proof of Theorem 4.4

*Proof.* Let  $d = \frac{z-r}{4r}$ . According to Lemma 4.11, we can construct in  $O(m)$  steps a  $d$ -grouped partition  $\mathcal{R} = \{R_1, \dots, R_l\}$  of  $V(G)$ , with a set of centers  $S = \{s_1, \dots, s_l\}$ , and also, for every  $i \in \llbracket 1, l \rrbracket$ , the distance-decompositions  $\mathcal{D}_i = (T_i, r_i, \mathcal{X}_i)$  with origins  $s_i$  of the graphs  $G[R_i]$ . For every  $i \in \llbracket 1, l \rrbracket$ , we use the notation  $\mathcal{X}_i = \{X_t^i\}_{t \in V(T_i)}$  and denote by  $N_i$  the node-frontiers of  $T_i$ .

Similarly, by applying the algorithm of Lemma 4.8, in  $O_r(m)$  steps we either find a  $\theta_r$ -major in  $G$  with at most  $2 \cdot r \cdot d \leq z$  edges or we know that for every  $i \in \llbracket 1, l \rrbracket$  and every  $t \in V(T_i)$ , the bag  $X_t^i$  contains at most  $r - 1$  vertices.

Using the algorithm of Lemma 4.13, in  $O_r(m)$  steps we either find a  $(2r - 2)$ -edge-protrusion with extension more than  $w$ , or we know that for every  $i \in \llbracket 1, l \rrbracket$ , all  $N_i$ -unimportant paths of  $T_i$  have length at most  $2w + 1$ .

We may now assume that none of the above algorithms provided a  $\theta_r$ -major with  $z$  edges, or a  $(2r - 2)$ -edge-protrusion.

From Corollary 4.15, for every  $i \in \llbracket 1, l \rrbracket$  the tree  $T_i$  contains at least  $2^{\frac{d-1}{2w+1}} = 2^{\frac{z-5r}{4r \cdot (2w+1)}}$  ports, which by definition means that there are at least  $2^{\frac{z-5r}{4r \cdot (2w+1)}}$  edges in  $G$  with one endpoint in  $R_i$  and the other in  $V(G) \setminus R_i$ .

By applying the algorithm of Lemma 4.12, in  $O_r(m)$  steps, we will either find a  $\theta_r$ -major in  $G$  with at most  $z = 4 \cdot r \cdot d + r$  edges or we know that for every two distinct  $i, j \in \llbracket 1, l \rrbracket$  there are at most  $r - 1$  edges of  $G$  with one endpoint in  $R_i$  and one in  $R_j$ . By Lemma 4.12, for every distinct integers  $i, j \in \llbracket 1, l \rrbracket$  there are at most  $r - 1$  edges with one endpoint in  $R_i$  and the other in  $R_j$ . As a consequence of the two previous implications, for every  $i \in \llbracket 1, l \rrbracket$  there is a set  $Z_i \subseteq \llbracket 1, l \rrbracket \setminus \{i\}$ , where  $|Z_i| \geq \frac{1}{r-1} 2^{\frac{z-5r}{4r(2w+1)}}$ , such that for every  $j \in Z_i$  there exists an edge with one endpoint in  $R_i$  and the other in  $R_j$ . Consequently, if we now contract all edges in  $G[R_i]$  for every  $i \in \llbracket 1, l \rrbracket$ , the resulting graph  $H$  is a minor of  $G$  of minimum degree at least  $\frac{1}{r-1} 2^{\frac{z-5r}{4r(2w+1)}}$ . Therefore, we output  $G$ , which is an  $H$ -major, as required in this case.  $\square$

### Proof of Theorem 4.5

*Proof.* The proof is quite similar to the one of Theorem 4.4. If  $G$  contains a vertex  $v$  of degree less than  $\delta$ , we can easily find it in  $O_r(m)$  steps. Hence, from now on we can assume that every vertex has degree at least  $\delta$ .

Let  $d = \frac{z-r}{4r}$ . From Lemma 4.11, in  $O(m)$  steps, we can construct a  $d$ -grouped partition  $\mathcal{R} = \{R_1, \dots, R_l\}$  of  $G$ , with a set of centers  $S = \{s_1, \dots, s_l\}$ , and also the distance-decomposition  $\mathcal{D}_i = (T_i, r_i, \mathcal{X}_i)$  with origins  $s_i$  of the graphs  $G[R_i]$ , for every  $i \in \llbracket 1, l \rrbracket$ . We use again the notation  $\mathcal{X}_i = \{X_t^i\}_{t \in V(T_i)}$ .

As in the proof of Theorem 4.4, in  $O_r(m)$  steps, we can either find a  $\theta_r$ -major in  $G$  with at most  $z = 4 \cdot r \cdot d + r$  edges or we know that for every distinct integers  $i, j \in \llbracket 1, l \rrbracket$  there are at most  $r - 1$  edges of  $G$  with one endpoint in  $R_i$  and one in  $R_j$  (cf. Lemma 4.12).

Using Corollary 4.10, we can in  $O_r(m)$  steps either find a  $\theta_r$ -major in  $G$  with at most  $z$  edges or we know that every bag of  $\mathcal{D}_i$  has less than  $r$  vertices, for every  $i \in$

$\llbracket 1, l \rrbracket$ . Let  $i \in \llbracket 1, l \rrbracket$  and let  $u \in R_i$  be a vertex at distance less than  $d$  from  $s_i$ . As  $u$  has degree at least  $3r$ , it must have neighbors in at least 3 different bags of  $\mathcal{D}_i$ , apart from the one containing it. This means that every vertex in  $T_i$  of distance less than  $d$  from  $r_i$  has degree at least  $\lfloor \frac{\delta}{r-1} \rfloor \geq 3$  and therefore  $T_i$  has at least  $\lfloor \frac{\delta}{r-1} - 1 \rfloor^d$  leaves. Notice also that if  $t$  is a leaf of  $T_i$ , then each vertex in  $X_t^i$  can have at most  $r - 1$  neighbors in  $X_{p(t)}^i$  and at most  $r - 2$  neighbors in  $X_t^i$ . Therefore there are at least  $\delta - (r - 1) - (r - 2) = \delta - 2r + 3$  edges in  $G$  with one endpoint in  $X_t^i$  and the other in  $V(G) \setminus R_i$ . This means that for every  $i \in \llbracket 1, l \rrbracket$  there are at least  $(\delta - 2r + 3) \cdot \lfloor \frac{\delta}{r-1} - 1 \rfloor^d$  edges with one endpoint in  $R_i$  and the other  $V(G) \setminus R_i$ .

Similarly to the proof of [Theorem 4.4](#), we deduce that, for each  $i \in \llbracket 1, l \rrbracket$ , there is a set  $Z_i \subseteq \llbracket 1, l \rrbracket \setminus \{i\}$  where  $|Z_i| \geq \frac{\delta - 2r + 3}{r - 1} \cdot \lfloor \frac{\delta}{r-1} - 1 \rfloor^d$  such that, for every  $j \in Z_i$ , there exists an edge with one endpoint in  $R_i$  and the other in  $R_j$ . This implies the existence of an  $H$ -major in  $G$  for some  $H$  with  $\delta(H) \geq \frac{\delta - 2r + 3}{r - 1} \cdot \lfloor \frac{\delta}{r-1} - 1 \rfloor^{\frac{z-r}{4r}}$ . We then output  $G$ , which, in this case, is an  $H$ -major.  $\square$

### 4.1.5 Remarks on the $H$ -girth

The results presented in this section are centered around the concept of  $\theta_r$ -girth which generalizes the usual notion of girth of a graph. A natural line of research is to investigate the  $H$ -girth parameter for different instantiations of  $H$ . An interesting problem in this direction could be to characterize the graphs  $H$  for which our results ([Theorem 4.2](#) and [Theorem 4.3](#)) can be extended.

From its definition, the  $H$ -girth is related to the minor relation. An other direction of research would be to extend the parameter of  $H$ -girth to other containment relations. One could consider, for a fixed graph  $H$ , the minimum size of an induced subgraph that can be contracted to  $H$ , or the minimum size of a subdivision of  $H$  in a graph. The first one of these parameters is related to induced minors and the second one to topological minors.

As the usual notion of girth appears in various contexts in graph theory, we wonder for which graphs  $H$  the results related to girth can be extended to the  $H$ -girth or to the two aforementioned variants.

## 4.2 Degree and $\theta_r$ -packings

In this section we show how lower bounds on the degree of the vertices of a graph can be used to prove the existence of a packing of  $\theta_r$ -majors.

### 4.2.1 On maximum degree and edge-disjoint packings

The main result of this section is the following. It relates the maximum degree of a 2-connected graph with the presence of an edge-disjoint union of  $\theta_r$ -majors.

**Lemma 4.16.** *Let  $k > 0, r > 0$  be two integers, and let  $G$  be a 2-connected graph with  $\Delta(G) \geq 2kr$ . Then  $G$  contains a subgraph that is the disjoint union of  $k$  edge-disjoint  $\theta_r$ -majors.*

For the purpose of the proof, we deal with graphs in which some vertices are marked. If  $G$  is a graph and  $m: V(G) \rightarrow \{0, 1\}$  is a function, we say that  $(G, m)$  is a graph *marked* by  $m$ . A vertex  $v$  of  $G$  such that  $m(v) = 1$  is said to be *marked*. We denote by  $\mu$  the function that, given a graph, returns its number of marked vertices. We now define an  $r$ -good partition. Given a positive integer  $r$ , a marked tree  $(T, m)$  is said to have an  $r$ -good partition of root  $v$  if there is a pair  $((T_1, m_1), (T_2, m_2))$  of marked trees such that:

- (i)  $T_1$  and  $T_2$  are subtrees of  $T$  such that  $(E(T_1), E(T_2))$  is a partition of  $E(T)$ ;
- (ii)  $r \leq \mu((T_1, m_1)) \leq 2r$ ;
- (iii)  $v \in V(T_2)$ ; and
- (iv) every vertex that is marked in  $(T, m)$  is either marked in  $(T_1, m_1)$  or marked in  $(T_2, m_2)$ , but not in both. In other words, for every  $u \in V(T)$ ,
  - if  $v \in V(T_1) \cap V(T_2)$  then  $m(v) = 1 \Leftrightarrow m_1(v) = 1$  or  $m_2(v) = 1$  but not both;
  - otherwise, let  $i \in \{1, 2\}$  be the integer such that  $v \in V(T_i)$ . Then we have  $m(v) = m_i(v)$ .

We remark that because of (iv),  $\mu(T) = \mu(T_1) + \mu(T_2)$ . If for every  $v \in V(T)$ ,  $(T, m)$  has an  $r$ -good partition of root  $v$ , then  $T$  is said to have an  $r$ -good partition.

**Lemma 4.17.** *For every integer  $r > 0$  and every marked tree  $(T, m)$ , if  $\mu(T) \geq 2r$  then  $(T, m)$  has an  $r$ -good partition.*

*Proof.* Let  $r > 0$  be an integer. We prove this lemma by induction on the size of the tree.

*Base case:*  $|V(T)| = 0$ . Since  $2r \geq 2 > |V(T)|$ ,  $T$  does not have  $2r$  marked vertices and we are done.

*Induction step:* Assume that for every integer  $n' < n$ , every marked tree  $(T', m')$  on  $n'$  vertices and satisfying  $\mu((T', m')) \geq 2r$  has an  $r$ -good partition (induction hypothesis).

Let us prove that every marked tree on  $n$  vertices has a  $r$ -good partition if it has at least  $2r$  marked vertices. Let  $(T, m)$  be a tree on  $n$  vertices and let  $v$  be a vertex of  $T$ . We assume that  $\mu((T, m)) \geq 2r$ . We distinguish two cases.

- $\mu((T, m)) = 2r$ :

Let  $T_1 = T$ , let  $m_1 = m$ , let  $T_2 = (\{v\}, \emptyset)$ , and let  $m_2: V(T_2) \rightarrow \{0, 1\}$  be the function equal to 0 on every vertex of  $T_2$ . Remark that  $(E(T_1), E(T_2)) = (E(T), \emptyset)$  is a partition of  $E(T)$ ,  $T_2$  contains  $v$ , and as  $(T, m)$  contains (exactly)  $2r$  marked vertices, so does  $(T_1, m)$ . Consequently  $((T_1, m_1), (T_2, m_2))$  is an  $r$ -good partition of  $(T, m)$ .

- $\mu((T, m)) > 2r$ :

We distinguish different cases depending on the degree of the root  $v$  in  $T$ .

Case 1:  $\deg(v) = 1$ .

Let  $u$  be the neighbor of  $v$  in  $T$ , let  $T' = T \setminus \{v\}$ , and  $m' = m|_{V(T')}$ . Remark that  $\mu((T', m')) \geq 2r$  and  $|V(T')| = |V(T)| - 1$ . By induction hypothesis,  $(T', m')$  has an  $r$ -good partition  $((T'_1, m_1), (T'_2, m_1))$  of root  $u$ . We extend it to  $T$  by setting  $T_1 = T'_1$  and  $T_2 = (V(T'_2) \cup \{v\}, E(T'_2) \cup \{v, u\})$ . Notice that  $T_2$  contains  $v$ . As the subtree  $T'_1$  contains at least  $r$  and at most  $2r$  marked vertices (induction hypothesis), so does  $T_1$ . Also, remark that  $(E(T_1), E(T_2))$  is a partition of  $E(T)$  and that since  $u \in T'_2$ , the graph  $T_2$  is connected. Therefore the pair  $(T_1, T_2)$  is an  $r$ -good partition of  $T$ .

Case 2:  $\deg(v) = d > 1$ .

Let  $u_1, \dots, u_d$  be the neighbors of  $v$  in  $T$  and for every  $i \in \llbracket 1, d \rrbracket$ , let  $C_i$  be the connected component of  $T \setminus \{v\}$  that contains  $u_i$ . We also define, for every  $i \in \llbracket 1, d \rrbracket$ , the restricted marking function  $w_i = m|_{V(C_i)}$ .

Subcase (a): there exists  $i \in \llbracket 1, d \rrbracket$  such that  $\mu((C_i, w_i)) > 2r$ .

Let  $T' = (V(C_i) \cup \{v\}, E(C_i) \cup \{u, v\})$  and let  $m' = m|_{V(T')}$ . Remark that  $|V(T')| < |V(T)|$  and  $\mu((T', m')) > 2r$ . According to the induction hypothesis,  $(T', m')$  has an  $r$ -good partition  $((T_1, m_1), (T_2, m_2))$  of root  $u_i$ . Similarly as before, we can extend it into an  $r$ -good partition  $((T_1, m_1), (T_2, m_2))$  of  $(T, m)$ . This is done by setting:

$$\begin{aligned} T_1 &= T'_1, \\ m_1 &= m'_1, \\ T_2 &= (V(T'_2) \cup \{v\}, E(T'_2) \cup \{v, u_i\}), \text{ and} \\ m_2 &: \begin{cases} v \mapsto 0 \\ u \in V(T_2) \setminus \{v\} \mapsto m'_2(u) \end{cases} \end{aligned}$$

Subcase (b): there exists  $i \in \llbracket 1, d \rrbracket$  such that  $r \leq \mu((C_i, w_i)) \leq 2r$ .

Let  $T_1 = C_i$  and  $T_2 = T[E(T) \setminus E(T_1)]$ . In this case,  $(E(T_1), E(T_2))$  is a partition of  $E(T)$  and  $T_2$  is connected since it contains  $v$ , the vertex which is adjacent to the  $C_j$ 's. Thus, if we set  $m_1 = m|_{V(T_1)}$  and  $m_2 = m|_{V(T_2)}$ ,  $((T_1, m_1), (T_2, m_2))$  is an  $r$ -good partition of  $(T, m)$ .

Subcase (c): for all  $i \in \llbracket 1, d \rrbracket$ ,  $\mu((C_i, w_i)) < r$ .

Let  $j = \min \left\{ j \in \llbracket 2, d \rrbracket, \sum_{i=1}^j \mu((C_i, w_i)) \geq r \right\}$ . We set:

$$\begin{aligned} T_1 &= (\cup_{i \in \llbracket 1, j \rrbracket} V(C_i) \cup \{v\}, \cup_{i \in \llbracket 1, j \rrbracket} (E(C_i) \cup \{v, u_i\})), \\ m_1 &: \begin{cases} v \mapsto 0 \\ u \in VT_1 \setminus \{v\} \mapsto m(u) \end{cases} \text{ ,} \\ T_2 &= T[E(T) \setminus E(T_1)], \text{ and} \\ m_2 &= m|_{V(T_2)}. \end{aligned}$$

By definition of  $j$ ,  $\mu((T_1, m_1)) \geq r$  and as for every  $i \in \llbracket 1, d \rrbracket$ ,  $\mu((C_i, w_i)) < r$  we also have  $\mu((T_1, m_1)) < 2r$ . As before, the pair  $((T_1, m_1), (T_2, m_2))$  is an  $r$ -good partition of  $(T, m)$ .

In conclusion, we proved by induction that for every integer  $r$ , every tree having at least  $2r$  marked vertices has an  $r$ -good partition.  $\square$

We are now ready to give the proof of [Lemma 4.16](#).

*Proof of Lemma 4.16.* As  $G$  is 2-connected, the removal of a vertex  $v$  of maximum degree gives a connected graph. Let  $T$  be a minimal tree of  $G \setminus \{v\}$  spanning the neighborhood  $N_G(v)$  of  $v$ . We mark the vertices of  $T$  that are elements of  $N_G(v)$ : this gives the marking function  $m$  for  $T$ . Let us prove by induction on  $k$  that  $(T, m)$  has  $k$  edge-disjoint marked subtrees  $(T_1, m_1), \dots, (T_k, m_k)$ , each containing at least  $r$  marked vertices. If we do so, then we are done because  $\{\{v\}, T_i\}_{i \in \llbracket 1, k \rrbracket}$  is a collection of  $k$  edge-disjoint  $\theta_r$  majors. In fact, as for every  $i \in \llbracket 1, k \rrbracket$ ,  $T_i$  contains  $r' \geq r$  vertices adjacent to  $v$  in  $G$ , contracting the edges of  $T_i$  in  $G[\{v\} \cup V(T_i)]$  gives the graph  $\theta_{r'}$ . Let  $r > 0$  be an integer.

*Base case  $k = 1$ :* Clear.

*Induction step  $k > 1$ :* Assume that for every  $k' < k$ , every tree with at least  $2k'r$  vertices marked has  $k'$  edge-disjoint subtrees, each with at least  $r$  marked vertices. Let  $(T, m)$  be a marked tree such that  $\mu((T, m)) \geq 2kr$ . According to [Lemma 4.17](#),  $(T, m)$  has an  $r$ -good partition  $((T_1, m_1), (T'_1, m'_1))$  such that  $r \leq \mu((T_1, m_1)) \leq 2r$  and  $\mu((T'_1, m'_1)) = \mu((T, m)) - \mu((T_1, m_1)) \geq 2(k-1)r$ . By induction hypothesis,  $(T'_1, m'_1)$  has  $k-1$  edge-disjoint marked subtrees  $(T_2, m_2), \dots, (T_k, m_k)$  each containing at least  $r$  marked vertices. Remark that as all these trees are subgraphs of  $T'_1$ , which is edge-disjoint from  $T_1$  in  $T$ , they are edge-disjoint from  $T_1$  as well. Consequently,  $(T_1, m_1), (T_2, m_2), \dots, (T_k, m_k)$  is the family of edge-disjoint subtrees we were looking for.  $\square$

*Remark 4.18.* Let  $r > 0$  be an integer. For every positive integer  $k$ , the graph  $\theta_{kr-1}$  does not contain  $k$  edge-disjoint  $\theta_r$ -majors. Therefore the bound provided by [Lemma 4.16](#) is tight up to a constant factor.

## 4.2.2 On the minimum degree and vertex-disjoint packings

In this subsection we show that every graph of large minimum degree contains  $k \cdot \theta_r$  as minor, which can be found in polynomial time. Our proof relies on the following result.

**Theorem 4.19** (Theorem 12 of [\[BTV07\]](#)). *Given  $k, r \in \mathbb{N}_{\geq 1}$  and an input graph  $G$  such that  $\delta(G) \geq k(r+1) - 1$ , a partition  $(V_1, \dots, V_k)$  of  $V(G)$  satisfying  $\forall i \in \llbracket 1, k \rrbracket, \delta(G[V_i]) \geq r$  can be found in  $O(n^c)$  steps, for some  $c \in \mathbb{N}$ .*

[Theorem 4.19](#) is the algorithmic version of the next older result.

**Theorem 4.20** ([\[Sti96, Corollary 3\]](#)). *For every  $k, r \in \mathbb{N}_{\geq 1}$ , every graph  $G$  with  $\delta(G) \geq k(r+1) - 1$  has a partition  $(V_1, \dots, V_k)$  of its vertex set satisfying  $\delta(G[V_i]) \geq r$  for every  $i \in \llbracket 1, k \rrbracket$ .*

**Theorem 4.21.** *There is an algorithm that, given  $k, r \in \mathbb{N}_{\geq 1}$  and a graph  $G$  with  $\delta(G) \geq kr$ , returns a subgraph of  $G$  containing an edge-disjoint union of  $k$  copies of a  $\theta_r$ -major, in  $O(m)$  steps.*

*Proof.* Starting from any vertex  $u$ , we grow a maximal (non induced) path  $P$  in  $G$  by iteratively adding to  $P$  a vertex that is adjacent to the previously added vertex but does not belong to  $P$ . Since  $\delta(G) \geq kr$ , any such path will have length at least  $kr + 1$ . At the end, all the neighbors of the last vertex  $v$  of  $P$  belong to  $P$  (otherwise  $P$  could be extended). Since  $v$  has degree at least  $kr$ ,  $v$  has at least  $kr$  neighbors in  $P$ . Let  $w_0, \dots, w_{kr-1}$  be an enumeration of the  $kr$  first neighbors of  $v$  in the order given by  $P$ , starting from  $u$ . For every  $i \in \llbracket 0, k-1 \rrbracket$ , let  $S_i$  be the subgraph of  $G$  induced by  $v$  and the subpath of  $P$  starting at  $w_{ir}$  and ending at  $w_{(i+1)r-1}$ . Observe that for every  $i \in \llbracket 0, k-1 \rrbracket$ ,  $S_i$  contains a  $\theta_r$ -major and that the intersection of every pair of graphs from  $\{S_i\}_{i \in \llbracket 0, k-1 \rrbracket}$  is  $\{v\}$ . Hence  $P$  contains a  $\mathcal{E}\text{-}\mathcal{M}(\theta_r)$ -packing of  $G$  of size  $k$ , as desired. Every edge of  $G$  is considered at most once in this algorithm, yielding to a running time of  $O(m)$  steps.  $\square$

**Corollary 4.22.** *There is an algorithm that, given  $r \in \mathbb{N}_{\geq 1}$  and a graph  $G$  with  $\delta(G) \geq r$ , returns a  $\theta_r$ -major of  $G$  in  $O(m)$ -steps.*

Observe that the previous lemma only deals with edge-disjoint packings. An analogue of **Theorem 4.21** for vertex-disjoint packings can be proved using **Theorem 4.19** to the price of a worse time complexity.

**Theorem 4.23.** *There is an algorithm that, given  $k, r \in \mathbb{N}_{\geq 1}$  and a graph  $G$  with  $\delta(G) \geq k(r+1) - 1$ , outputs a  $(k \cdot \theta_r)$ -major of  $G$  in  $O(n^c + m)$  steps, where  $c$  is the constant of **Theorem 4.19**.*

*Proof.* After applying the algorithm of **Theorem 4.19** on  $G$  to obtain in  $O(n^c)$ -time  $k$  graphs  $G[V_1], \dots, G[V_k]$ , we extract a  $\theta_r$ -major from each of them using **Corollary 4.22**.  $\square$

### 4.3 Treewidth and excluded majors

In this section, we show upper-bounds on the treewidth of graphs not containing a major of some fixed pattern, among which: the wheel, the double wheel, any graph of pathwidth at most 2, the yurt graph, and the disjoint union of copies of the graph  $\theta_r$ .

In [**RST94a**], Robertson, Seymour, and Thomas proved that every planar graph is a minor of a large enough grid.

**Lemma 4.24** ([**RST94a**, (1.5)]). *For every positive integer  $h$ , the  $h \times h$ -grid contains as a minor any planar graph  $H$  satisfying  $2|V(H)| + 4|E(H)| \leq h$ .*

Together with the Grid Exclusion Theorem which upper-bounds the treewidth of graph excluding a grid (**Theorem 4.1**), the above lemma implies the next result.

**Theorem 4.25.** *There is a function  $f: \mathbb{N} \rightarrow \mathbb{N}$  such that, for every for every planar graph  $H$  on  $h$  vertices, every graph  $G$  that does not contain a minor isomorphic to  $H$  has treewidth at most  $f(h)$ .*

The original proof of **Theorem 4.1** in [RS86] does not provide any explicit estimation for the function  $f$ . Later, in [RST94a], Robertson, Seymour, and Thomas proved the same result for  $f(h) = 2^{O(h^5)}$ , while a less complicated proof appeared in [DJGT99a]. The bound  $f(h) \leq h - 2$  was also obtained in [BRST91] in the case where  $H$  is required to be a forest. **Theorem 4.25** has several applications in algorithms and a lot of research has been devoted to optimizing the function  $f$  in general or for specific instantiations of  $H$  (see [RST94b, DJGT99b]).

For a long time, whether **Theorem 4.25** can be proved for a polynomial  $f$  was an open problem. In [RST94a], an  $\Omega(h^2 \cdot \log h)$  lower bound was provided for the best possible estimation of  $f$  and was also conjectured that the optimal estimation should not be far away from this lower bound. In fact, a more precise variant of the same conjecture was given by Demaine, Hajiaghayi, and Kawarabayashi in [DHK09] where they conjectured that **Theorem 4.25** holds for  $f(h) = O(h^3)$ . The bounds of [RST94a] were then improved by Kawarabayashi and Kobayashi [iKK12], where they show that **Theorem 4.25** holds for  $f(h) = 2^{O(h \cdot \log h)}$ . The same bounds were obtained by Leaf and Seymour [LS15]. Until recently, this was the best known estimation of the function  $f$ .

In a breakthrough result [CC14], Chekuri and Chuzhoy proved that **Theorem 4.1** holds for  $f(h) = O(h^{228})$ . Chuzhoy recently improved this bound.

**Theorem 4.26** ([Chu16], see also [CC14, Chu15]). *There exists a function  $f_1: \mathbb{N} \rightarrow \mathbb{N}$  with  $f_1(h) = O(h^{19} \text{polylog } h)$  such that, for every integer  $h$ , every graph that does not contain a minor isomorphic to the  $h \times h$ -grid has treewidth at most  $f_1(h)$ .*

The remaining open question is whether the degree of this polynomial bound can be substantially reduced in general. In this direction, one may still consider restrictions either on the graph  $G$  or on the graph  $H$  that yield a low polynomial dependence between the treewidth and the size of the excluded minor. In the first direction, Demaine and Hajiaghayi proved in [DH08] that, when  $G$  is restricted to belong to some graph class excluding some fixed graph as a minor, then **Theorem 4.25** (optimally) holds for  $f(h) = O(h)$ . Similar results have been proved by Fomin, Saurabh, and Lokshantov, in [FLS12], for the case where  $G$  is either a unit disk graph or a map graph that does not contain a clique as a subgraph.

In a second direction, one may consider  $H$  to be some specific planar graph and find a good upper bound for the treewidth of the graphs that exclude it as a minor. More generally, we can consider a parametrized class of planar graphs  $\mathcal{H}_k$  where each graph in  $\mathcal{H}_k$  has size (or a given parameter) bounded by a polynomial in  $k$ , and prove that the following fragment of **Theorem 4.25** holds for some low degree polynomial function  $f: \mathbb{N} \rightarrow \mathbb{N}$ :

$$\forall k \geq 0 \forall H \in \mathcal{H}_k, \text{ if } H \not\leq_m G \text{ then } \text{tw}(G) \leq f(k). \quad (4.2)$$

The question can be stated as follows: find pairs  $(\mathcal{H}_k, g(k))$  for which (4.2) holds for some  $f(k) = O(g(k))$ , where  $\mathcal{H}_k$  is as general as possible and  $g$  is as small as possible (and certainly polynomial). It is known, for example, that (4.2) holds for the pair  $(\{C_k\}, k)$ , where  $C_k$  is the cycle or a path of  $k$  vertices (see e.g. [Bod93, FL94]), and for the pair  $(\{K_{2,k}\}, k)$ , (see [BvLTT97b]). Two more results in the same direction that appeared in the last decade are the following: according to the result of Birmelé, Bondy, and Reed in [BBR07a], (4.2) holds for the pair  $(\mathcal{P}_k, k^2)$  where  $\mathcal{P}_k$  contains all minors of  $K_2 \times C_k$  (we denote by  $K_2 \times C_k$  the Cartesian product of  $K_2$  and the cycle of  $k$  vertices, also known as the  $k$ -prism). Finally, one of the consequences of the recent results of Leaf and Seymour in [LS15], implies that (4.2) holds for the pair  $(\mathcal{F}_k, k)$ , where  $\mathcal{F}_k$  contains every graph on  $k$  vertices that contains a vertex that meets all its cycles.

**Results presented in this section.** In this section we provide new exclusion theorems by proving that (4.2) holds for the pairs:

- $(\mathcal{H}_k^0, k^2)$ , where  $\mathcal{H}_k^0$  contains all simple graphs  $H$  on  $k$  vertices and of pathwidth at most 2;
- $(\mathcal{H}_k^1, k)$ , where  $\mathcal{H}_k^1$  contains all minors of a wheel on  $k+1$  vertices – see Figure 2.3;
- $(\mathcal{H}_k^2, k^2 \log^2 k)$ , where  $\mathcal{H}_k^2$  contains all minors of a double wheel on  $k+2$  vertices – see Figure 2.3;
- $(\mathcal{H}_k^3, k^4)$ , where  $\mathcal{H}_k^3$  contains all minors of the yurt graph on  $2k+1$  vertices (i.e. the graph obtained if we take a  $(2 \times k)$ -grid and add a new vertex adjacent with all the vertices of its “upper layer” – see Figure 4.6); and
- $(\mathcal{H}_{k,r}^4, k \log k)$ , where  $\mathcal{H}_{k,r}^4$  contains all minors of the graph  $k \cdot \theta_r$ .

Notice that none of the classes  $\mathcal{H}_k^0$ ,  $\mathcal{H}_k^1$ ,  $\mathcal{H}_k^2$ , and  $\mathcal{H}_k^3$  is minor comparable with the classes  $\mathcal{P}_k$  and  $\mathcal{F}_k$  treated in [BBR07a] and [LS15], whereas  $\mathcal{H}_{k,r}^4 \subsetneq \mathcal{P}_{kr}$ . Moreover,  $\mathcal{H}_k^1 \subsetneq \mathcal{H}_k^2 \subsetneq \mathcal{H}_k^3$ , while  $\mathcal{H}_{k,r}^3 \subsetneq \mathcal{H}_k^0$ . The above results are presented thereafter in detail, without the  $O$ -notation.

**Theorem 4.27.** *Let  $k > 0$  be an integer and  $G$  be a graph. If  $\text{tw}(G) \geq 36k - 2$ , then  $G$  contains a wheel of order  $k$  as minor.*

**Theorem 4.28.** *Let  $k > 0$  be an integer and  $G$  be a graph. If  $\text{tw}(G) \geq 12(8k \log(8k) + 2)^2 - 4$ , then  $G$  contains a double wheel of order at least  $k$  as minor.*

**Theorem 4.29.** *Let  $G$  be a graph, let  $H$  be a simple graph on  $k$  vertices such that  $\text{pw}(H) \leq 2$ . If  $\text{tw}(G) \geq 3(k-2)^2 - 1$  then  $G$  contains  $H$  as a minor.*

**Theorem 4.29** can be extended to the setting of graphs that are not simple as follows.

**Corollary 4.30.** *Let  $G$  be a graph, let  $H$  be a graph such that  $\text{pw}(H) \leq 2$  and let  $k = |V(H)| + |E(H)|$ . If  $\text{tw}(G) \geq 3(k-2)^2 - 1$  then  $G$  contains  $H$  as a minor.*

**Theorem 4.31.** *Let  $k > 0$  be an integer and  $G$  be a graph. If  $\text{tw}(G) \geq 6k^4 - 24k^3 + 48k^2 - 48k + 23$ , then  $G$  contains the yurt graph of order  $k$  as minor.*

**Theorem 4.32.** *Let  $k > 0$  and  $r \geq 2$  be two integers and let  $G$  be a graph. If  $\text{tw}(G) \geq 2^{6r} \cdot k \cdot \log(k+1)$ , then  $G$  contains  $k \cdot \theta_r$  as a minor.*

The aforementioned results will we proved in the forthcoming sections, in this order. The proofs of the four first results use as a departure point the results of Leaf and Seymour in [LS15], whereas the last one uses the results of Geelen, Gerards, Robertson, and Whittle on the excluded minors for the matroids of branch-width  $k$  [GGRW03] together with Theorem 4.4. In Subsection 4.3.1, we introduce notions that we will use and we prove two lemmas, for later use. We give in Subsection 4.3.7 lower bounds on the best function  $f$  of (4.2) that one can expect for the classes studied here and we discuss the tightness of our results.

### 4.3.1 Our tools

This section contains some definitions, as well as two lemmas that will be useful later.

**Definition 4.33** (linked set). Let  $G$  be a graph and  $S \subseteq V(G)$ . The set  $S$  is said to be *linked* in  $G$  if for every two subsets  $X_1, X_2$  of  $S$  (not necessarily disjoint) such that  $|X_1| = |X_2|$ , there is a set  $Q$  of  $|X_1|$  (vertex-)disjoint paths between  $X_1$  and  $X_2$  in  $G$  whose length is not one (but can be null) and whose endpoints only are in  $S$ .

**Definition 4.34** (left-contains, [LS15]). Let  $H$  be a graph on  $r$  vertices,  $G$  a graph and  $(A, B)$  a separation of order  $r$  in  $G$ . We say that  $(A, B)$  *left-contains*  $H$  if there is a  $H$ -minor model  $\mu$  in  $G[A]$  such that  $|\mu(v) \cap B| = 1$  for every  $v \in V(H)$ .

In this section, we denote by  $B_h$  the complete binary tree of height  $h$ , for every integer  $h > 0$ . Given a tree  $T$ , we denote by  $L(T)$  the set of its leaves.

**Proposition 4.35** ([LS15, (4.3)]). *Let  $k > 0$  be an integer, let  $F$  be a forest on  $k$  vertices and let  $G$  be a graph. If  $\text{tw}(G) \geq \frac{3}{2}k - 1$ , then  $G$  has a separation  $(A, B)$  of order  $k$  such that*

- $G[B \setminus A]$  is connected;
- $A \cap B$  is linked in  $G[B]$ ;
- $(A, B)$  left-contains  $F$ .

**Proposition 4.36** (Erdős–Szekeres Theorem, [ES87]). *Let  $k$  and  $\ell$  be two positive integers. Then any sequence of  $(\ell - 1)(k - 1) + 1$  distinct integers contains either an increasing subsequence of length  $k$  or a decreasing subsequence of length  $\ell$ .*

Recall that if  $T$  is a tree and  $x, y \in V(T)$ , we denote by  $xTy$  the unique path of  $T$  linking  $x$  to  $y$ , and by  $xT\overset{\circ}{y}$  the same path were  $y$  has been deleted.

**Lemma 4.37.** *For every tree  $T$ ,  $|V(T)| \leq \frac{|L(T)| \cdot \text{diam}(T)}{2} + 1$ .*

*Proof.* Root  $T$  to a vertex  $r \in V(T)$  that is halfway of a longest path of  $T$ . For each leaf  $x \in L(T)$ , we know that  $|V(xT\overset{\circ}{r})| \leq \left\lfloor \frac{\text{diam}(T)}{2} \right\rfloor$ . Observe that  $V(T) = \{r\} \cup \bigcup_{x \in L(T)} V(xT\overset{\circ}{r})$ . Therefore,

$$\begin{aligned} |V(T)| &\leq \sum_{x \in L(T)} |V(xT\overset{\circ}{r})| + 1 \\ |V(T)| &\leq |L(T)| \cdot \left\lfloor \frac{\text{diam}(T)}{2} \right\rfloor + 1. \end{aligned}$$

Notice that equality holds for the subdivided star (obtained from  $K_{1,n}$  by subdividing  $k$  times every edge, for some  $n, k \in \mathbb{N}$ ).  $\square$

**Definition 4.38** (The set  $\Lambda(T)$ ). Let  $T$  be a tree. We denote by  $\Lambda(T)$  the set containing every graph obtained as follows: take the disjoint union of  $T$ , a path  $P$  where  $|V(P)| \geq \sqrt{|L(T)|}$ , and an extra vertex  $v_{\text{new}}$ , and add edges such that

- (i) there is an edge between  $v_{\text{new}}$  and every vertex of  $P$ ;
- (ii) there are  $|V(P)|$  disjoint edges between  $P$  and  $L(T)$ ;
- (iii) there are no more edges than the edges of  $T$  and  $P$  and the edges mentioned in (i) and (ii).

**Lemma 4.39.** *Let  $n \geq 1$  be an integer,  $T$  be a tree on  $n$  vertices and let  $G$  be a graph. If  $\text{tw}(G) \geq 3n - 1$ , then  $H \leq_m G$  for some  $H \in \Lambda(T)$ .*

*Proof.* Let  $n$ ,  $T$ , and  $G$  be as in the statement of the lemma. Let  $l$  be the number of leaves of  $T$ , and let  $J$  be a path on  $l$  vertices. We consider the disjoint union of  $J$  and  $T$ .

The graph  $G$  has treewidth at least  $\frac{3}{2}(n + l) - 1$ , then by [Proposition 4.35](#),  $G$  has a separation  $(A, B)$  of order  $n + l$  such that

- (i)  $G[B \setminus A]$  is connected;
- (ii)  $A \cap B$  is linked in  $G[B]$ ;
- (iii)  $(A, B)$  left-contains the graph  $J \cup T$ .

Let  $\varphi$  be a  $J \cup T$ -minor model in  $G[A]$  that witnesses (iii). We call the vertices of  $A \cap B$  that belong to  $\varphi(v)$  for some  $v \in V(J)$  the  $J$ -part, and vertices that belong to  $\varphi(v)$  for some  $v \in L(T)$  forms the  $L(T)$ -part. Notice that two distinct vertices of the  $J$ -part (respectively  $L(T)$ -part) belong to the images by  $\mu$  of two distinct vertices of  $J$  (respectively  $L$ ) by the definition of the left-containment.

Let  $\mathcal{P}$  a set of  $l$  disjoint paths with the one endpoint in the  $J$ -part and the other in the  $L(T)$ -part, and whose interior belongs to  $B \setminus A$ . The existence of such paths is given by (ii). For each  $P \in \mathcal{P}$ , we arbitrarily choose a vertex  $v_P$  of the interior of  $P$ , that is,  $v_P \in V(P) \setminus A$ . By (i),  $G[B \setminus A]$  is connected: let  $Y$  be a smallest tree spanning the vertices  $\{v_P\}_{P \in \mathcal{P}}$ . Let  $s = \sqrt{|L(T)|}$ , and let  $Y^*$  be the tree obtained from  $Y$  by dissolving every vertex of degree two that is not  $v_P$  for some  $P \in \mathcal{P}$ . We are now facing two possible situations.

*Case 1:*  $Y^*$  has a path of length  $s$ . Let  $Q$  be the path of  $Y$  corresponding to a path of length  $s$  in  $Y^*$  and let  $S$  be the set of vertices  $u \in V(Q)$  that are not of degree two or that are  $v_P$  for some  $P \in \mathcal{P}$ . Observe that from every  $u \in S$ , there is a path  $J_u$  to the  $L(T)$ -part and a path  $J'_u$  to the  $J$ -part. Indeed, if  $u = v_P$  for some  $P \in \mathcal{P}$ , then  $u$  is a vertex of  $P$  linking (by definition) a vertex of the  $L(T)$ -part to a vertex of the  $J$ -part. Otherwise,  $u$  is of degree at least 3 in  $Y$  and every leaf of the subtrees of  $Y \setminus Q$  (at least one of which is adjacent to  $u$ ), is a  $v_P$  for some  $P \in \mathcal{P}$  (by minimality of  $Y$ ), so is connected to the  $L(T)$ -part and the  $J$ -part as explained above. Furthermore, for every two distinct  $u, v \in S$ , the aforementioned paths are disjoint.

Let us now summarize.  $G[\cup_{v \in V(J)} \varphi(v)]$  is a connected subgraph of  $G$ , which is connected by the  $s$  disjoint paths  $J'_{uu \in S}$  to the path  $Y$ . All the endpoints of the paths  $J'_{uu \in S}$  on  $Y$  are connected by  $s$  disjoint paths  $J_{uu \in S}$  to the  $L(T)$ -part, which correspond to the leaves in a major of  $T$ . Therefore this graph contains a member of  $\Lambda(T)$  as a minor, as required.

*Case 2:*  $\text{diam}(Y^*) < s$ . From Lemma 4.37,  $|L(Y)| = |L(Y^*)| \geq s$ . Observe that  $L(Y) \subseteq \{v_P\}_{P \in \mathcal{P}}$  (this follows by the minimality of  $Y$ ). Let  $S = V(Y) \setminus L(Y)$ . We consider the minor of  $G$  obtained by contracting, for every  $P \in \mathcal{P}$  such that  $v_P \in L(Y)$ , every edge of the subpath connecting the  $J$ -part to a leaf of  $Y$ . In this graph,  $S$  induces a connected subgraph adjacent to at least  $s$  distinct vertices of the  $J$ -part. All these  $s$  vertices of the  $J$ -part are connected by  $s$  disjoint paths to distinct vertices of the  $L(T)$ -part. Thus this contains a member of  $\Lambda(T)$  as a minor, and so do  $G$ . □

The proof of Theorem 4.32 requires additional definitions, that we introduce in the corresponding section as they are not used in the proofs of the other results.

### 4.3.2 Excluding a wheel

In this section we prove Theorem 4.27. For every integer  $r \geq 3$ , we denote by  $W_r$  the wheel of order  $r$  where every vertex of the cycle is adjacent to the center. An example is given in Figure 4.3.

Recall that in this section,  $B_h$  is the complete binary tree of height  $h$ , for every integer  $h > 0$ .

**Lemma 4.40.** *Let  $h > 2$  be an integer. Let  $G$  be a graph obtained from the union of the tree  $T = B_h$  and a path  $P$  by adding the edges  $\{l, \psi(l)\} \in E(G)$  for every  $l \in L(T)$ , where  $\psi: L(T) \rightarrow V(P)$  is a bijection. Then  $G$  contains a wheel of order  $2^{h-2} + 1$  as a minor.*




Figure 4.3: A wheel of order six (left) and a double wheel of order 6 (right).

*Proof.* Let  $h, \psi, T, P = p_1 \dots p_{2^h}$  and  $G$  be as above. Let  $r$  be the root of  $T$ .

In the arguments to follow, if  $t \in V(T)$ , we denote by  $T_t$  the subtree of  $T$  rooted at  $t$  (i.e. the subtree of  $T$  whose vertices are all the vertices  $t' \in V(T)$  such that the path  $t'Tr$  contains  $t$ ).

We consider the vertices  $u = \psi^{-1}(p_1) \in L(T)$  and  $v = \psi^{-1}(p_{2^h}) \in L(T)$ .

Let  $\tau$  be a largest subtree of  $T$  which is disjoint from  $uTv$ . Let  $L_\tau = L(\tau) \cap L(T)$  and let  $Q = \psi(L_\tau) \subseteq P$ . It is not hard to see that  $G$  contains  $W_{|Q|+1}$  as a minor. Indeed, the paths  $P$  and  $uTv$  together with the edges  $\{p_1, u\}$  and  $\{p_{2^h}, v\}$  form a cycle in  $G$ . Besides, the tree  $\tau$ , which is disjoint from this cycle, has at least  $|Q| + 1$  vertices that are adjacent to distinct vertices of  $P$ :  $|Q|$  of them are the elements of  $Q$ , and the other one is the (only) vertex of  $\tau$  adjacent to  $uTv$  (which exists by maximality of  $\tau$ ). In the subgraph of  $G$  induced by  $V(P) \cup V(uTv) \cup V(\tau)$ , contracting  $\tau$  to a vertex gives a vertex adjacent to at least  $|Q| + 1$  vertices of a (non necessarily induced) cycle, a graph containing  $W_{|Q|+1}$  as subgraph.

Depending on  $G$ ,  $|Q|$  may take different values. However, we show that it is never less than  $2^{h-2}$ . Remember,  $|Q|$  is the number of leaves that a largest subtree of  $T$  that is disjoint from  $uTv$  shares with  $T$ . The root  $r$  of  $T$  has two children  $r_1$  and  $r_2$ , inducing two subtrees  $T_{r_1}$  and  $T_{r_2}$  of  $T$ . Let  $w = \text{lca } T(u, v) \in V(T) \setminus L(T)$ .

*Case 1.*  $w \neq r$ . As  $w \neq r$ ,  $w$  is a vertex of one of  $\{T_{r_1}, T_{r_2}\}$ , say  $T_{r_1}$ , which contains also  $u$  and  $v$ , and thus the path  $uTv$ . The other subtree  $T_{r_2}$  is then disjoint from  $uTv$ , it has height  $h - 1$  and is complete so it has  $2^{h-1}$  leaves. Consequently, in this case  $|Q| \geq 2^{h-1}$ .

*Case 2.*  $w = r$ . In this case, the path  $uTv$  contains  $r$  (and  $r \neq u, r \neq v$  as  $u$  and  $v$  are leaves) so  $u$  and  $v$  are not in the same subtree of  $\{T_{r_1}, T_{r_2}\}$  and  $uTv$  contains the two edges  $\{r, r_1\}$  and  $\{r, r_2\}$ . For every  $i \in \{1, 2\}$ , we denote by  $r_{i,1}$  and  $r_{i,2}$  the two children of  $r_i$  in  $T$ . We assume without loss of generality that  $u \in V(T_{r_{1,1}})$  and  $v \in V(T_{r_{2,1}})$  (if not, we just rename the  $r_i$ 's and  $r_{i,j}$ 's). Notice that the path  $uTv$  is the concatenation of the paths  $uT_{r_1}r_1, r_1T_{r_2}, r_2T_{r_2}v$ . Since the tree  $T_{r_{1,2}}$  is disjoint from  $uTv$ , is complete and is of height  $h - 2$ , it has  $2^{h-2}$  leaves. Therefore we have

$$|Q| \geq 2^{h-2}.$$

In both cases,  $|Q| \geq 2^{h-2}$  and according to what we proved before,  $G$  contains a major of  $W_{|Q|+1}$ . As every wheel contains as a minor every smaller wheel, we proved that  $G$  contains a wheel of order at least  $2^{h-2}$ .  $\square$

We are now ready to prove **Theorem 4.27**. Let us first recall its statement.

**Theorem 4.27.** *Let  $k > 0$  be an integer and  $G$  be a graph. If  $\text{tw}(G) \geq 36k - 2$ , then  $G$  contains a wheel of order  $k$  as minor.*

*Proof.* Let  $k > 0$  be an integer,  $G$  be a graph such that  $\text{tw}(G) \geq 36k - 2$ , and let  $h = \lceil \log 4k \rceil$ . Since there is a major of a wheel in every bigger wheel, we have

$$\begin{aligned} W_k &\leq_m W_{2^{\lceil \log k \rceil} + 1} \\ &\leq_m W_{2^{\lceil (\log 4k) - 2 \rceil} + 1} \\ &\leq_m W_{2^{h-2} + 1} \end{aligned}$$

Therefore, if we prove that there is a  $W_{2^{h-2}+1}$ -major in  $G$ , then we are done because the minor relation is transitive. Let  $Y_h^-$  be the graph of the following form: the disjoint union of the complete binary tree  $B_h$  of height  $h$  with leaves set  $Y_L$  and of the path  $Y_P$  on  $2^h$  vertices, and let  $\mathcal{Y}_h$  be the set of graphs of the same form, but with the extra edges  $\{\{l, \phi(l)\}\}_{l \in Y_L}$ , where  $\phi : Y_L \rightarrow V(Y_P)$  is a bijection. As we proved in **Lemma 4.40** that every graph in  $\mathcal{Y}_h$  contains the wheel of order  $2^{h-2} + 1$  as minor, showing that  $G$  contains an element of  $\mathcal{Y}_h$  as minor suffices to prove this lemma. That is what we will do.

From our initial assumption, we deduce the following.

$$\begin{aligned} \text{tw}(G) &\geq 36k - \frac{5}{2} \\ &\geq \frac{3}{2}(3 \cdot 2^{\log 8k} - 1) - 1 \\ &\geq \frac{3}{2}(3 \cdot 2^{\lceil \log 4k \rceil + 1} - 1) - 1 \\ \text{tw}(G) &\geq \frac{3}{2}(3 \cdot 2^h - 1) - 1 \end{aligned}$$

According to **Proposition 4.35**,  $G$  has a separation  $(A, B)$  of order  $3 \cdot 2^h - 1$  such that

- (i)  $G[B \setminus A]$  is connected;
- (ii)  $A \cap B$  is linked in  $G[B]$ ;
- (iii)  $(A, B)$  left-contains the graph  $Y_h^-$ .

By definition of *left-contains*, there is an  $Y_h^-$ -minor model in  $G[A]$  such that every element of  $\mathcal{M}^- := \{\varphi^-(v), v \in Y_h^-\}$  contains exactly one element of  $A \cap B$ . For every  $x \in A \cap B$ , we denote by  $M_x^-$  the element of  $\mathcal{M}^-$  that contains  $x$ . Let  $L$  (respectively  $R$ ) be the subset of  $A \cap B$  of vertices that belong to an element of  $\mathcal{M}$  related to the leaves of  $B_h$  in  $Y_h^-$  (respectively to the path  $P$ ). We remark that these sets are both of cardinality  $2^h$ .

Since  $A \cap B$  is linked in  $G[B]$  (see (ii)), there is a set  $\mathcal{P}$  of  $2^h$  disjoint paths between the vertices of  $L$  and the elements of  $R$ . Let  $\psi: L \rightarrow V(P)$  be the function that match each element  $l$  of  $L$  with the (unique) element of  $R$  it is linked to by a path (that we call  $\mathcal{P}_l$ ) of  $\mathcal{P}$ . Observe that  $\psi$  is a bijection. We set

$$\begin{aligned} \forall l \in L, M_l &= M_l^- \cup V(l\mathcal{P}_l\psi(l)) \\ \forall r \in (A \cap B) \setminus L, M_r &= M_r^- \end{aligned}$$

$$\mathcal{M} = \bigcup_{x \in A \cup B} M_x.$$

Let us show that  $\mathcal{M}$  allows us to define a minor model of some  $H \in \mathcal{Y}_h$ . Let us consider the following mapping.

$$\varphi: \begin{cases} V(Y_h^-) & \rightarrow \mathcal{M} \\ x & \mapsto M_x \end{cases}$$

We claim that  $\varphi$  is an  $H$ -minor model, for some  $H \in \mathcal{Y}_h$ . This is a consequence of the following remarks.

*Remark 4.41.* Every element of  $\mathcal{M}$  is either an element of  $\mathcal{M}^-$ , or the union of a element  $M$  of  $\mathcal{M}^-$  and of the vertices of a path that start in  $M$ , thus every element of  $\mathcal{M}$  induces a connected subgraph of  $G$ .

*Remark 4.42.* The paths of  $\mathcal{P}$  are all disjoint and are disjoint from the elements of  $\mathcal{M}^-$ . Every interior of path of  $\mathcal{P}$  is in but one element of  $\mathcal{M}$ , therefore the elements of  $\mathcal{M}$  are disjoint.

*Remark 4.43.* The elements  $\{m_l\}_{l \in L}$  are in bijection with the elements of  $\{m_r\}_{r \in R}$  (thanks to the function  $\psi$ ) and every two vertices  $l \in L$  and  $\psi(l) \in R$  are such that there is an edge between  $m_l$  and  $m_{\psi(l)}$  (by definition of  $\mathcal{M}^+$ ).

We just proved that  $\varphi$  is a minor model of a graph of  $\mathcal{Y}_h$  in  $G$ . Finally, we apply [Lemma 4.40](#) to find a minor model of the wheel of order  $2^{h-2} + 1 = 2^{\lceil \log k \rceil - 2} + 1 \geq k$  in  $G$  and this concludes the proof.  $\square$

### 4.3.3 Excluding a double wheel

This section is devoted to the proof of [Theorem 4.28](#). Recall that for every integer  $n \geq 3$ , a double wheel of order  $n$  is obtained from a cycle of order  $n$  by adding two non-adjacent, each connected to at least three vertices of the cycle. We denote by  $W_n^2$  the double wheel of order  $n$  where the two extra vertices are adjacent to every vertex of the cycle.

**Lemma 4.44.** *Let  $G$  be a graph and  $h > 0$  be an integer. If  $\text{tw}(G) \geq 6 \cdot 2^h - 4$ , then  $G$  contains as minor a double wheel of order at least  $\frac{2^{\frac{h}{2}} - 2}{2^{h-3}}$ .*

*Proof.* Let  $h$  and  $G$  be as above. Observe that  $\text{tw}(G) \geq 3(2^{h+1} - 1) - 1$ . As the binary tree  $T = B_h$  has  $2^{h+1} - 1$  vertices,  $G$  contains a graph  $H \in \Lambda(B_h)$  as minor (by **Lemma 4.39**). Let us show that any graph  $H \in \Lambda(B_h)$  contains a double wheel of order at least  $\frac{2^{\frac{h}{2}} - 2}{2^{h-3}}$  as minor.

Let  $P$  be the path of length at least  $2^{\frac{h}{2}}$  in the definition of  $H$ . Let  $L$  be the set, of size at least  $2^{\frac{h}{2}}$ , of the leaves of  $T$  that are adjacent to  $P$  in  $H$ . Such a set exists by definition of  $\Lambda(B_h)$ . We also define  $u$  (respectively  $u'$ ) as the vertex of  $L(T)$  that is adjacent to one end of  $P$  (respectively to the other end of  $P$ ) and  $Q = uTu'$ .

As  $T$  is a binary tree of height  $h$ ,  $Q$  has at most  $2h - 1$  vertices. Each vertex of  $Q$  is of degree at most 3 in  $T$  except the two ends which are of degree 1. Consequently,  $T \setminus Q$  has at most  $2h - 3$  connected components that are subtrees of  $T$ . Notice that every vertex of the  $2^{\frac{h}{2}}$  elements of  $L$  is either a leaf of one of these  $2h - 3$  subtrees, or one of the two ends of  $Q$ . By the pigeonhole principle, one of these subtrees, which we call  $T_1$ , has at least  $\frac{2^{\frac{h}{2}} - 2}{2^{h-3}}$  leaves that are elements of  $L$ .

Let  $M_{o_1}$  be the set of vertices of this subtree  $T_1$ . We also set  $M_{o_2} = \{v_{\text{new}}\}$  (cf. **Definition 4.38** for a definition of  $v_{\text{new}}$ ). Let us consider the cycle  $C$  made by the concatenation of the paths  $uPu'$  and  $u'Tu$  in  $H$ .

By definition of  $M_{o_1}$ , there are at least  $\frac{2^{\frac{h}{2}} - 2}{2^{h-3}}$  vertices of  $C$  adjacent to vertices of  $M_{o_1}$ . Let  $J = \{j_1, \dots, j_{|J|}\}$  be the set of such vertices of  $C$ , in the same order as they appear in  $C$  (we then have  $|J| \geq \frac{2^{\frac{h}{2}} - 2}{2^{h-3}}$ ).

We arbitrarily choose an orientation of  $C$  and define the sets of vertices  $M_1, M_2, \dots, M_{|J|}$  as follows.

$$\begin{aligned} \forall i \in \llbracket 1, |J| - 1 \rrbracket, M_i &= V(j_i C j_{i+1}^\circ) \\ M_{|J|} &= V(j_{|J|} C j_1^\circ) \end{aligned}$$

Let  $\mathcal{M} = \{M_1, \dots, M_{|J|}, M_{o_1}, M_{o_2}\}$  and  $\psi: V(W_{|J|}^2) \rightarrow \mathcal{M}$  be the function defined by

$$\begin{aligned} \forall i \in \llbracket 1, |J| \rrbracket, \psi(w_i) &= M_i \\ \psi(o_1) &= M_{o_1} \\ \psi(o_2) &= M_{o_2} \end{aligned}$$

Notice that  $\psi$  maps the vertices of  $W_{|J|}^2$  to connected subgraphs of  $H$  such that  $\forall (v, w) \in E(W_{|J|}^2)$ , there is a vertex of  $\psi(v)$  adjacent in  $H$  to a vertex of  $\psi(w)$ . Therefore,  $\psi$  is a  $W_{|J|}^2$ -minor model in  $H$ .

Since  $|J| \geq \frac{2^{\frac{h}{2}} - 2}{2^{h-3}}$ ,  $H$  contains a double wheel of order at least  $\frac{2^{\frac{h}{2}} - 2}{2^{h-3}}$ , which is what we wanted to show.  $\square$

**Corollary 4.45.** *Let  $l > 0$  be an integer and  $G$  be a graph. If  $\text{tw}(G) \geq 12l - 4$  then  $G$  contains a double wheel of order at least  $\frac{\sqrt{l}-2}{2\log l-5}$  as minor.*

*Proof.* Let  $l$  and  $G$  be as above. First remark that

$$\lceil \log l \rceil - 1 \leq \log l \leq \lceil \log l \rceil \quad (4.3)$$

Our initial assumption on  $\text{tw}(G)$  gives the following.

$$\begin{aligned} \text{tw}(G) &\geq 12l - 4 \\ &\geq 6 \cdot 2^{\log(2l)} - 4 \\ &\geq 6 \cdot 2^{\log l + 1} - 4 \\ &\geq 6 \cdot 2^{\lceil \log l \rceil} - 4 \end{aligned} \quad \text{by (4.3)}$$

By **Lemma 4.44**,  $G$  contains a double wheel of order at least

$$\begin{aligned} q &= \frac{2^{\frac{\lceil \log l \rceil}{2}} - 2}{2 \lceil \log l \rceil - 3} \\ &\geq \frac{2^{\frac{1}{2} \log l} - 2}{2(\log l - 1) - 3} \quad \text{by (4.3)} \\ &\geq \frac{\sqrt{l} - 2}{2 \log l - 5} \end{aligned}$$

Therefore,  $G$  contains a double wheel of order  $q \geq \frac{\sqrt{l}-2}{2\log l-5}$ , as required.  $\square$

We can now deduce **Theorem 4.28**, that we recall hereafter, from **Corollary 4.45**.

**Theorem 4.28.** *Let  $k > 0$  be an integer and  $G$  be a graph. If  $\text{tw}(G) \geq 12(8k \log(8k) + 2)^2 - 4$ , then  $G$  contains a double wheel of order at least  $k$  as minor.*

*Proof.* Applying **Corollary 4.45** for  $l = (8k \log(8k) + 2)^2$  yields that  $G$  contains a double wheel of order at least

$$\begin{aligned} q &\geq \frac{\sqrt{l} - 2}{2 \log l - 5} \\ &\geq \frac{8k \log(8k)}{4 \log(8k \log(8k) + 2) - 5} \\ &\geq \frac{8k \log(8k)}{4 \log(8k \log(8k)) - 1} \\ &\geq \frac{8k \log(8k)}{4(\log(8k) + \log \log(8k)) - 1} \\ &\geq \frac{8k \log(8k)}{8 \log(8k) - 1} \\ &\geq k \end{aligned}$$

Consequently  $G$  contains a double wheel of order  $q \geq k$  and we are done.  $\square$

### 4.3.4 Excluding a graph of pathwidth at most 2

This sections contains the proofs of [Theorem 4.29](#) and [Corollary 4.30](#). We define the graph  $\Xi_r$  as the graph of the following form (see [Figure 4.4](#)).

$$\begin{cases} V(G) = \{x_0, \dots, x_{r-1}, y_0, \dots, y_{r-1}, z_0, \dots, z_{r-1}\} \\ E(G) = \{\{x_i, x_{i+1}\}, \{z_i, z_{i+1}\}\}_{i \in \llbracket 1, r-1 \rrbracket} \cup \{\{x_i, y_i\}, \{y_i, z_i\}\}_{i \in \llbracket 0, r-1 \rrbracket} \end{cases}$$




Figure 4.4: The graph  $\Xi_5$ .

#### Graphs of pathwidth 2 in $\Xi_r$

Instead of proving that a treewidth quadratic in  $|V(H)| + |E(H)|$  forces an  $H$ -minor for every graph  $H$  of pathwidth at most 2, we prove that a treewidth quadratic in  $r$  forces an  $\Xi_r$ -major and then that every graph  $H$  of pathwidth at most 2 with  $|V(H)| + |E(H)| \leq r$  is a minor of  $\Xi_r$ . Recall that every graph has an optimal path decomposition which is nice. Let us state some observations on path decompositions.

*Remark 4.46.* Let  $G$  be a graph and let  $(p_1 p_2 \dots p_k, \mathcal{X})$ ,  $\mathcal{X} = \{X_{p_i}\}_{i \in \llbracket 1, k \rrbracket}$  be a nice (non necessarily optimal) path decomposition of  $G$ . Let  $w$  be the width of this decomposition.

For every  $i \in \llbracket 2, k-1 \rrbracket$ , and if  $p_i$  is a forget node,  $|X_{p_i}| \leq w-1$  and  $p_{i+1}$  is an introduce node, then by setting

$$\begin{aligned} X'_{p_i} &= X_{p_{i-1}} \cup X_{p_{i+1}} \\ \forall j \in \llbracket 1, k \rrbracket, j \neq i, X'_{p_j} &= X_{p_j} \\ \mathcal{X}' &= \left\{ X'_{p_j} \right\}_{j \in \llbracket 1, k \rrbracket} \end{aligned}$$

we create from  $(p_1 p_2 \dots p_k, \mathcal{X}')$  a valid path decomposition of  $G$ , where  $p_i$  is now an introduce node and  $p_{i+1}$  a forget node. Observe that  $|X'_{p_i}| \leq |X_{p_i}| + 2 = w+1$ . Therefore the new path decomposition has the same width as the original one. Note that the condition  $|X_{p_i}| \leq w-1$  holds, for instance, when  $p_{i-1}$  is required to be a forget node too (for  $i \in \llbracket 3, k-1 \rrbracket$ ).

*Remark 4.47.* Let  $G$  be a graph and  $P = (p_1 p_2 \dots p_k, \mathcal{X})$  be a nice path decomposition of  $G$ . For every  $i \in \llbracket 1, k \rrbracket$ , the path  $p_1 \dots p_i$  contains at most as many forget nodes as introduce nodes and the difference between these two numbers is at most  $w+1$  where  $w$  is the width of  $P$ .

**Lemma 4.48.** *Let  $G$  be a graph on  $n$  vertices. Then  $G$  has an optimal path decomposition  $P$  such that*

- (i) *every bag of  $P$  has size  $\text{pw}(G) + 1$ ;*
- (ii) *every two adjacent bags differs by exactly one element, i.e. for every two adjacent vertices  $u$  and  $v$  of  $P$ ,  $|X_u \setminus X_v| = |X_v \setminus X_u| = 1$ .*

*Proof.* Let  $P = (p_1 p_2 \dots p_{2k}, \mathcal{X})$  with  $\mathcal{X} = \{X_{p_i}\}_{i \in \llbracket 1, 2k \rrbracket}$  be a nice optimal path decomposition of  $G$  with as many introduce nodes (respectively forget nodes) as there are vertices in  $G$ .

Let  $s = \text{pw}(G) + 1$ . According to [Remark 4.46](#) and [Remark 4.47](#),  $P$  can be modified into a path decomposition of  $G$  of the same width and such that

- (a) the  $s$  first vertices of  $P$  are introduce nodes and  $p_{s+1}$  is a forget node;
- (b) the  $s$  last vertices of  $P$  are forget nodes and  $p_{2k-s}$  is an introduce node;
- (c) for every  $i \in \llbracket s, 2k - s \rrbracket$ ,  $p_i$  and  $p_{i+1}$  are nodes of different type.

In the arguments to follow, we assume that  $P$  satisfies this property.

*Remark 4.49.* Introduce nodes all have bags of cardinality  $s$ .

*Remark 4.50.* For every  $i \in \llbracket 0, k - s \rrbracket$ , the node  $p_{s+2i}$  is an introduce node and the node  $p_{s+2i+1}$  is a forget node, which implies  $X_{p_{s+2i}} \subsetneq X_{p_{s+2i+1}}$ . Also note that for every  $i \in \llbracket 1, s - 1 \rrbracket$ ,  $X_{p_i} \subsetneq X_{p_s}$  and for every  $i \in \llbracket 2k - s + 1, 2k \rrbracket$ ,  $X_{p_i} \subsetneq X_{p_{2k-s}}$ .

Intuitively, every bag  $X$  that is included in one of its adjacent bags  $X'$  contains no more information than what  $X'$  already contains, so we will just remove it.

We thus define  $P' = p_s p_{s+2} \dots p_{s+2i} \dots p_{2k-s}$  (a path made of all introduce nodes of  $P$ ). Clearly,  $P$  and  $P'$  have the same width and as we deleted only redundant nodes,  $P'$  is still a valid path decomposition of  $G$ .

Since every two adjacent nodes of  $P'$  were introduce nodes separated by a forget node in  $P$ , they only differ by one element. According to [Remark 4.49](#) and since every node of  $P'$  was an introduce node in  $P$ , every bag of  $P'$  have size  $\text{pw}(G) + 1$ . Consequently,  $P'$  is an optimal path decomposition that satisfies the conditions of the lemma statement.  $\square$

*Remark 4.51.* The path decomposition of [Lemma 4.48](#) has length  $V(G) - \text{pw}(G)$ .

*Proof.* Let  $(P, \mathcal{X})$  be such a path decomposition. Remember that the first node of  $P$  has a bag of size  $\text{pw}(G) + 1$  and that every two adjacent nodes of  $P$  have bags which differs by exactly one element. Since every vertex of  $G$  is in a bag of  $P$ , in addition to the first bag containing  $\text{pw}(G) + 1$  vertices of  $G$ ,  $P$  must have  $V(G) - \text{pw}(G) - 1$  other bags in order to contain all vertices of  $G$ . Therefore  $P$  has length  $V(G) - \text{pw}(G)$ .  $\square$

A proof of a slightly weaker version of the following lemma previously appeared [[Pro89](#)].

**Lemma 4.52.** *For every simple graph  $G$  on  $n$  vertices and of pathwidth at most 2, there is a  $G$ -major in  $\Xi_{n-1}$ .*

*Proof.* Let  $G$  be as in the statement of the lemma. We assume that  $\text{pw}(G) = 2$  (if this is not the case we add edges to  $G$  in order to obtain a graph of pathwidth 2 which contains  $G$  as a minor). Let  $r = V(G) - \text{pw}(G) = n - 2$ .

Let  $P = (p_1 \dots p_r, \{X_{p_1}, \dots, X_{p_r}\})$  be an optimal path decomposition of  $G$  satisfying the properties of [Lemma 4.48](#), of length  $r$ . Such decomposition exists according to [Lemma 4.48](#) and [Remark 4.51](#).

Using this decomposition, we will now define a labeling  $\lambda$  of the vertices of  $\Xi_{r+1}$ . When dealing with the vertices of  $\Xi_{r+1}$  we will use the notations given in the definition of this graph. Let  $\lambda: V(\Xi_{r+1}) \rightarrow V(G)$  be the function defined as follows:

(a)  $\lambda(x_0)$  and  $\lambda(y_0)$  are both equal to one (arbitrarily chosen) element of the set  $X_{p_1} \cap X_{p_2}$ ;

(b)  $\lambda(z_0)$  is equal to the only element of the set  $X_{p_1} \cap X_{p_2} \setminus \{\lambda(x_1)\}$ ;

(c)  $\forall i \in \llbracket 2, r \rrbracket$ ,  $\lambda(y_i) = X_{p_i} \setminus X_{p_{i-1}}$  and we consider two cases:

Case 1:  $X_{p_{i-1}} \cap X_{p_i} = X_{p_i} \cap X_{p_{i+1}}$   
 $\lambda(x_i) = \lambda(x_{i-1})$  and  $\lambda(z_i) = \lambda(z_{i-1})$ ;

Case 2:  $X_{p_{i-1}} \cap X_{p_i} \neq X_{p_i} \cap X_{p_{i+1}}$   
if  $X_{p_{i-1}} \cap X_{p_i} \cap X_{p_{i+1}} = \lambda(x_{i-1})$ ,  
then  $\lambda(x_i) = \lambda(x_{i-1})$  and  $\lambda(z_i) = X_{p_i} \setminus X_{p_{i-1}}$ ;  
else  $\lambda(x_i) = X_{p_i} \setminus X_{p_{i-1}}$  and  $\lambda(z_i) = \lambda(z_{i-1})$ .

Thanks to this labeling, we are now able to present a  $G$ -minor model in  $\Xi_{r+1}$  :

$$\forall v \in V(G), M_v = \{u \in V(\Xi_{r+1}), \lambda(u) = v\}$$

$$\mathcal{M} = \{M_v\}_{v \in V(G)}$$

$$\varphi: \begin{cases} V(G) & \rightarrow \mathcal{M} \\ u & \mapsto M_u \end{cases}$$

To show that  $\varphi$  is a  $G$ -minor model in  $\Xi_{r+1}$ , we now check if it matches the definition.

By definition, every element of  $\mathcal{M}$  is a subset of  $V(\Xi_{r+1})$ . To show that every element of  $\mathcal{M}$  induces a connected subgraph in  $G$ , it suffices to show that nodes of  $\Xi_{r+1}$  which have the same label induces a connected subgraph in  $G$  (by construction of the elements of  $\mathcal{M}$ ). This can easily be seen by remarking that for every  $i \in \llbracket 2, r \rrbracket$ , every vertex  $y_i$  of  $\Xi_{r+1}$  gets a new label and that every vertex  $x_i$  (respectively  $z_i$ ) of  $\Xi_{r+1}$  receive either the same label as  $y_i$ , or the same label as  $x_{i-1}$  (respectively  $z_{i-1}$ ).

Let us show that this labeling ensure that if two vertices  $u$  and  $v$  of  $G$  are in the same bag of  $P$ , there are two adjacent vertices of  $\Xi_{r+1}$  that respectively gets labels  $u$  and  $v$ . Let  $u, v$  be two vertices of  $G$  which are in the same bag of  $P$ . Let  $i$  be such that  $X_i$  is the first bag of  $P$  (with respect to the subscripts of the bags of  $P$ ) which contains

both  $u$  and  $v$ . The case  $i = 1$  is trivial so we assume that  $i > 1$ . We also assume without loss of generality that  $X_i \setminus X_{i-1} = \{v\}$ , what gives  $\lambda(y_i) = v$ . Depending on in what case we are, either either  $\lambda(x_i) = u$  (**(c1)**) or  $\lambda(z_i) = u$  (**(c1)** and **(c2)**). In both cases,  $u$  and  $v$  are the labels of two adjacent nodes of  $\Xi_{r+1}$ . By construction of the elements of  $\mathcal{M}$ , this implies that if  $\{u, v\} \in E(G)$ , then there are vertices  $u' \in \varphi(u)$  and  $v' \in \varphi(v)$  such that  $\{u', v'\} \in E(\Xi_{r+1})$ .

Therefore,  $\varphi$  is a  $G$ -minor model in  $\Xi_{n-1}$ , what we wanted to find.  $\square$

Observe that **Lemma 4.52** can be straightforwardly extended to the setting of graphs that are not simple. Indeed, given a graph  $G$ , one can subdivide once every edge in order to obtain a simple graph  $G'$  that contains  $G$  as a minor, and then apply **Lemma 4.52** on  $G'$ , that satisfies  $|V(G')| = |V(G)| + |E(G)|$ .

**Corollary 4.53.** *For every graph  $G$  of pathwidth at most 2, there is a  $G$ -major in  $\Xi_n$ , where  $n = |V(G)| + |E(G)| - 1$ .*

### Exclusion of $\Xi_r$

**Lemma 4.54.** *For any graph, if  $\text{tw}(G) \geq 3\ell - 1$  then  $G$  contains as minor the following graph: a path  $P = p_1 \dots p_{2\ell}$  of length  $2\ell$  and a family  $Q$  of  $\ell$  paths of length 2 such that every vertex of  $P$  is the end of exactly one path of  $Q$  and every path of  $Q$  has one end in  $p_1 \dots p_\ell$  (the first half of  $P$ ) and the other end in  $p_{\ell+1} \dots p_{2\ell}$  (the second half of  $P$ ) (see **Figure 4.5**).*




Figure 4.5: Example for **Lemma 4.54**.

*Proof.* Let  $\ell > 0$  be an integer and  $G$  be a graph of treewidth at least  $3\ell - 1$ . According to **Proposition 4.35**,  $G$  has a separation  $(A, B)$  of order  $2\ell$  such that

- (i)  $G[B \setminus A]$  is connected;
- (ii)  $A \cap B$  is linked in  $G[B]$ ;
- (iii)  $(A, B)$  left-contains a path  $P = p_1 \dots p_{2\ell}$  of length  $2\ell$ .

Let  $\varphi$  be a  $P$ -minor model in  $G[A]$ , and let  $\mathcal{M} = \{\varphi(v), v \in V(P)\}$ . Let us call  $M_1, \dots, M_{2\ell}$  the elements of  $\mathcal{M}$ . We assume without loss of generality that  $\varphi$  maps  $p_i$  to  $M_i$  for every  $i \in \llbracket 1, 2\ell \rrbracket$ .

As  $A \cap B$  is linked in  $G[B]$ , there is a set  $Q$  of  $\ell$  disjoint paths in  $G[B]$  of length at least 2 and such that every path  $q \in Q$  has one end in  $(A \cap B) \cap \bigcup_{i \in \llbracket 1, \ell \rrbracket} M_i$ , the other end in  $(A \cap B) \cap \bigcup_{i \in \llbracket \ell+1, 2\ell \rrbracket} M_i$  and its internal vertices are not in  $A \cap B$ .

Let  $G'$  be the graph obtained from  $G[(\bigcup_{q \in Q} V(q)) \cup (\bigcup_{M \in \mathcal{M}} M)]$  after the following operations.

1. iteratively contract the edges of every path of  $Q$  until it reaches a length of 2. The paths of  $Q$  have length at least 2, so this is always possible.
2. for every  $i \in \llbracket 1, 2\ell \rrbracket$ , contract  $M_i$  to a single vertex. The image of a minor model induce connected subgraphs (by definition) thus this operation can always be performed.

As one can easily check, the graph  $G'$  is the graph we were looking for and it has been obtained by contracting some edges of a subgraph of  $G$ , therefore  $G' \leq_m G$ .  $\square$

Let us now prove [Theorem 4.29](#). First, we restate it.

**Theorem 4.29.** *Let  $G$  be a graph, let  $H$  be a simple graph on  $k$  vertices such that  $\text{pw}(H) \leq 2$ . If  $\text{tw}(G) \geq 3(k-2)^2 - 1$  then  $G$  contains  $H$  as a minor.*

*Proof.* Let  $G, H$  and  $h$  be as in the statement of the Lemma. According to [Lemma 4.52](#), every simple graph  $F$  on  $n$  vertices and of pathwidth at most two is a minor of  $\Xi_{n-1}$ . Therefore, in order to show that  $H \leq_m G$  it is enough to prove that  $\Xi_{k-1} \leq_m G$ . This is what we will do.

According to [Lemma 4.54](#),  $G$  contains as minor two paths  $P = p_1 \dots p_{(k-2)2}$  and  $R = r_1 \dots r_{(k-2)2}$  and a family  $Q$  of  $(k-2)^2$  paths of length 2 such that every vertex of  $P$  or  $R$  is the end of exactly one path of  $Q$  and every path of  $Q$  has one end in  $P$  and the other end in  $R$ . For every  $p \in P$ , we denote by  $\varphi(p)$  the (unique) vertex of  $R$  to which  $p$  is linked to by a path of  $Q$ . Observe that  $\varphi$  is a bijection. By [Proposition 4.36](#), there is a subsequence  $P' = (p'_1, p'_2, \dots, p'_{k-1})$  of the vertices of  $P$  such that the vertices  $\varphi(p'_1), \varphi(p'_2), \dots, \varphi(p'_{k-1})$  appear in  $R$  either in this order or in the reverse order. Let  $R' = (\varphi(p'_1), \varphi(p'_2), \dots, \varphi(p'_{k-1}))$  and  $Q'$  be the set of inner vertices of the paths from  $p'_i$  to  $\varphi(p'_i)$  for all  $i \in \llbracket 1, k-1 \rrbracket$ .

Iteratively contracting in  $G$  the edges of  $P$  (respectively  $R$ ) which have at most one end in  $P'$  (respectively in  $R'$ ) and removing the vertices that are not in  $P', R'$  or  $Q'$  gives the graph  $\Xi_{k-1}$ . The operations used to obtain it are vertices and edge deletions, and edge contractions, thus  $\Xi_{k-1}$  is a minor of  $G$ . This concludes the proof.  $\square$

Observe that [Theorem 4.29](#) can be extended to the setting of graphs that are not simple as we did for [Corollary 4.53](#), what gives [Corollary 4.30](#).

### 4.3.5 Excluding a yurt graph

In this section we prove [Theorem 4.31](#). For every positive integer  $n$ , we denote by  $Y_n$  the Yurt graph of order  $n$  (as a reminder, see [Figure 4.6](#) or refer to [Subsection 2.3.2](#)).




Figure 4.6: The yurt graph of order 5.

For every  $r > 0$ , we define the *comb of order  $r$*  as the tree made from the path  $p_1 p_2 \dots p_r$  and the extra vertices  $v_1, v_2, \dots, v_r$  by adding an edge between  $p_i$  and  $v_i$  for every  $i \in \llbracket 1, r \rrbracket$ .

By using [Lemma 4.39](#) we can immediately prove [Theorem 4.31](#). Let us remind its statement.

**Theorem 4.31.** *Let  $k > 0$  be an integer and  $G$  be a graph. If  $\text{tw}(G) \geq 6k^4 - 24k^3 + 48k^2 - 48k + 23$ , then  $G$  contains the yurt graph of order  $k$  as minor.*

*Proof.* Let  $k > 0$  be an integer and  $G$  be a graph such that  $\text{tw}(G) \geq 6k^4 - 24k^3 + 48k^2 - 48k + 23$ . Let  $C$  be the comb with  $l = k^4 - 4k^3 + 8k^2 - 8k + 4$  teeth. As  $\text{tw}(G) \geq 3|V(C)| - 1$ ,  $G$  contains as a minor some graph of  $\Lambda(C)$  by [Lemma 4.39](#).

Let us prove that every graph of  $\Lambda(C)$  contains the yurt graph of order  $k$ . Let  $H$  be a graph of  $\Lambda(C)$ . We respectively call  $T$ ,  $P$  and  $o$  the tree, path and extra vertex of  $\Lambda(C)$ . Let  $F$  be the subset of edges between  $P$  and the leaves of  $T$

Let  $L = l_0, \dots, l_{k^2-2k+2}$  (respectively  $Q = q_0, \dots, q_{k^2-2k+2}$ ) be the leaves of  $T$  (respectively of  $P$ ) that are the end of an edge of  $F$ . We assume without loss of generality that they appear in this order.

According to [Proposition 4.36](#), there is a subsequence  $Q'$  of  $Q$  of length  $k$  such that the corresponding vertices  $L'$  of  $L$  appear in the same order. As one can easily see, this graph contains the yurt of order  $k$  and we are done.  $\square$

### 4.3.6 Excluding a union of $k$ disjoint copies of $\theta_r$

This section is devoted to the proof of [Theorem 4.32](#). Before we present the proof, we need to introduce some definitions and related results.

#### Preliminaries

Let  $G$  be a graph and  $G_1, G_2$  two non-empty subgraphs of  $G$ . We say that  $(G_1, G_2)$  is a *dissociation* of  $G$  if:

- $V(G_1) \cup V(G_2) = V(G)$ ; and
- $(E(G_1), E(G_2))$  is a partition of  $E(G)$ .

Let  $G$  be a graph. Given a set  $E \subseteq E(G)$ , we define  $V_E$  as the set of all endpoints of the edges in  $E$ . Given a partition  $(E_1, E_2)$  of  $E(G)$  we define  $\delta(E_1, E_2) = |V_{E_1} \cap V_{E_2}|$ .

A *cut*  $C = (X, Y)$  of  $G$  is a partition of  $V(G)$  into two subsets  $X$  and  $Y$ . We define the *cut-set* of  $C$  as  $E_C = \{\{x, y\} \in E(G) \mid x \in X \text{ and } y \in Y\}$  and call  $|E_C|$  the *order* of the cut. Also, given a graph  $G$ , we denote by  $\sigma(G)$  the number of connected components of  $G$ .

**The branchwidth of a graph.** A *branch-decomposition* of a graph  $G$  is a pair  $(T, \tau)$  where  $T$  is a ternary tree and  $\tau$  a bijection from the edges of  $G$  to the leaves of  $T$ . Deleting any edge  $e$  of  $T$  partitions the leaves of  $T$  into two sets, and thus the edges of  $G$  into two subsets  $E_1^e$  and  $E_2^e$ . The *width* of a branch-decomposition  $(T, \tau)$  is equal to  $\max_{e \in E(T)} \{\delta(E_1^e, E_2^e)\}$ . The *branchwidth* of a graph  $G$ , denoted  $\text{bw}(G)$ , is defined as the minimum width over all branch-decompositions of  $G$ .

**The branchwidth of a matroid.** We assume that the reader is familiar with the basic notions of matroid theory. We will use the standard notation from Oxley's book [Oxl92]. The branchwidth of a matroid is defined very similarly to that of a graph. Let  $\mathcal{M}$  be a matroid with finite ground set  $E(\mathcal{M})$  and rank function  $r$ . The *order* of a non-trivial partition  $(E_1, E_2)$  of  $E(\mathcal{M})$  is defined as  $\lambda(E_1, E_2) = r(E_1) + r(E_2) - r(E) + 1$ . A *branch-decomposition* of a matroid  $\mathcal{M}$  is a pair  $(T, \mu)$  where  $T$  is a ternary tree and  $\mu$  is a bijection from the elements of  $E(\mathcal{M})$  to the leaves of  $T$ . Deleting any edge  $e$  of  $T$  partitions the leaves of  $T$  into two sets, and thus the elements of  $E(\mathcal{M})$  into two subsets  $E_1^e$  and  $E_2^e$ . The *width* of a branch-decomposition  $(T, \mu)$  is equal to  $\max_{e \in E(T)} \{\lambda(E_1^e, E_2^e)\}$ . The *branchwidth* of a matroid  $\mathcal{M}$ , denoted  $\text{bw}(\mathcal{M})$ , is again defined as the minimum width over all branch-decompositions of  $\mathcal{M}$ . The *cycle matroid* of a graph  $G$  denoted  $\mathcal{M}_G$ , has ground set  $E(\mathcal{M}_G) = E(G)$  and the cycles of  $G$  as the cycles of  $\mathcal{M}_G$ . Let  $G$  be a graph,  $\mathcal{M}_G$  its cycle matroid and  $(G_1, G_2)$  a dissociation of  $G$ . Then clearly  $(E(G_1), E(G_2))$  is a partition of  $E(\mathcal{M}_G)$ , but to avoid confusion we will henceforth denote it  $(E_1, E_2)$  and we will call it *the partition of  $\mathcal{M}_G$  that corresponds to the dissociation  $(G_1, G_2)$  of  $G$* . Observe that the order of this partition is:

$$\lambda(E_1, E_2) = \delta(E(G_1), E(G_2)) - \sigma(G_1) - \sigma(G_2) + \sigma(G) + 1. \quad (\star)$$

**Minor obstructions.** Let  $\mathcal{G}$  be a graph class. We denote by  $\text{obs}(\mathcal{G})$  the set of all minor-minimal graphs  $H$  such that  $H \notin \mathcal{G}$  and we will call it *the minor obstruction set for  $\mathcal{G}$* . Clearly, if  $\mathcal{G}$  is closed under minors, the minor obstruction set for  $\mathcal{G}$  provides a complete characterization for  $\mathcal{G}$ : a graph  $G$  belongs in  $\mathcal{G}$  if and only if none of the graphs in  $\text{obs}(\mathcal{G})$  is a minor of  $G$ .

Given a class of matroids  $\mathbf{M}$ , *the minor obstruction set for  $\mathbf{M}$* , denoted by  $\text{obs}(\mathbf{M})$ , is defined very similarly to its graph-counterpart: it is simply the set of all minor-minimal matroids  $\mathcal{M}$  such that  $\mathcal{M} \notin \mathbf{M}$ .

We will need the following results.

**Proposition 4.55** ([RS91, Theorem 5.1]). *Let  $G$  be a graph of branchwidth at least 2. Then,  $\text{bw}(G) \leq \text{tw}(G) + 1 \leq \lfloor \frac{3}{2} \text{bw}(G) \rfloor$ .*

**Proposition 4.56** ([BvLTT97a]). Let  $r \in \mathbb{N}_{\geq 1}$  and let  $G$  be a graph. If  $\mathbf{bw}(G) \geq 2r + 1$ , then  $G$  contains a  $\theta_r$ -major.

**Proposition 4.57** ([HJ07, Theorem 4]). Let  $G$  be a graph that contains a cycle and  $\mathcal{M}_G$  be its cycle matroid. Then,  $\mathbf{bw}(G) = \mathbf{bw}(\mathcal{M}_G)$ .

**Proposition 4.58** ([GGRW03, Lemma 4.1]). Let a matroid  $\mathcal{M}$  be a minor obstruction for the class of matroids of branchwidth at most  $k$  and let  $g(n) = (6^{n-1} - 1)/5$ . Then, for every partition  $(X, Y)$  of  $\mathcal{M}$  with  $\lambda(X, Y) \leq k$ , either  $|X| \leq g(\lambda(X, Y))$  or  $|Y| \leq g(\lambda(X, Y))$ .

The following observations are also crucial.

*Observation 4.59.* Let  $\mathcal{G}$  be a graph class that is closed under minors and let  $\mathcal{M}_{\mathcal{G}} = \{\mathcal{M}_G \mid G \in \mathcal{G}\}$ .  $\mathcal{G}$  is minor closed if and only if  $\mathcal{M}_{\mathcal{G}}$  is minor closed. Moreover, for every  $H \in \mathbf{obs}(\mathcal{G})$  it holds that  $\mathcal{M}_H \in \mathbf{obs}(\mathcal{M}_{\mathcal{G}})$ .

The above observation is a direct consequence of the definition of matroid removal/contraction, e.g., see Proposition 4.9 of [Pit14].

*Observation 4.60.* There is a  $c \in \mathbb{R}_{\geq 2}$ , such that for any integer  $k \geq r \geq 2$ , if  $g(n) = (6^{n-1} - 1)/5$ , then  $\frac{1}{r-1} 2^{\frac{c^r \log k - 5r}{4r(2g(2r-2)+1)}} \geq k(r+1) - 1$ . Moreover, this holds for  $c = 2^6 \log_r \frac{2}{3}$ .

Now we are ready to prove [Theorem 4.32](#). Let us recall its statement.

**Theorem 4.32.** Let  $k > 0$  and  $r \geq 2$  be two integers and let  $G$  be a graph. If  $\mathbf{tw}(G) \geq 2^{6r} \cdot k \cdot \log(k+1)$ , then  $G$  contains  $k \cdot \theta_r$  as a minor.

*Proof.* For every  $r \in \mathbb{N}$ , we define  $f(r) = \frac{2}{3} 2^{6r}$ . By [Proposition 4.55](#), it is enough to prove that if  $\mathbf{bw}(G) \geq f(r) \cdot k \cdot \log(k+1)$ , then  $G$  contains  $k \cdot \theta_r$  as a minor. To prove this we use induction on  $k$ .

The case where  $k = 1$  follows from [Proposition 4.56](#) and the fact that  $f(r) \geq 2r + 1$ . We now examine the case where  $k > 1$ , assuming that the proposition holds for smaller values of  $k$ . As  $\mathbf{bw}(G) \geq f(r) \cdot k \cdot \log(k+1)$ ,  $G$  contains a minor obstruction for the class of graphs of branchwidth at most  $f(r) \cdot k \cdot \log(k+1) - 1$ .

*Claim 4.61.* Any  $(2r - 2)$ -edge-protrusion of  $G$  has extension at most  $g(2r - 2)$ .

*Proof of Claim 4.61.* Let  $C = (X, Y)$  be a cut in  $G$  of order at most  $2r - 2$  (one exists as soon as  $G$  has a  $(2r - 2)$ -edge-protrusion) and let  $G_X$  be the subgraph of  $G$  with  $V(G_X) = X \cup N_G(X)$  and let  $E(G_X) = E(G[X]) \cup E_C$ . Clearly the pair  $(G_X, G[Y])$  is a dissociation of  $G$ . Let  $\mathcal{M}_G$  be the cycle matroid of  $G$  and  $(E_X, E_Y)$  be the partition of  $\mathcal{M}_G$  that corresponds to the aforementioned dissociation. By [Proposition 4.57](#),  $\mathbf{bw}(\mathcal{M}_G) = \mathbf{bw}(G) \geq f(r) \cdot k \cdot \log(k+1)$ . Therefore, by [Observation 4.59](#),  $\mathcal{M}_G$  is a minor obstruction for the class of matroids of branchwidth  $f(r) \cdot k \cdot \log(k+1) - 1$ . We set  $\lambda = \lambda(E_X, E_Y)$ . From [\(\\*\)](#), we have:

$$\begin{aligned} \lambda &= r(E_X) + r(E_Y) - r(\mathcal{M}_G) + 1 \\ &= \delta(E(G_X), E(G[Y])) - \sigma(G_X) - \sigma(G[Y]) + \sigma(G) + 1 \\ &\leq \delta(E(G_X), E(G[Y])) \\ &\leq |E_C| = 2r - 2 \\ &\leq f(r) \cdot k \cdot \log(k+1) - 1. \end{aligned}$$

Thus, by [Proposition 4.58](#), either  $|E_X| \leq g(\lambda)$  or  $|E_Y| \leq g(\lambda)$ . Since  $g$  is non-decreasing, either  $|E(G_X)| \leq g(2r-2)$  or  $|E(G[Y])| \leq g(2r-2)$ . This directly implies that for any  $(2r-2)$ -edge-protrusion  $Z$  of  $G$ ,  $G[Z \cup N_G(Z)]$  has at most  $g(2r-2)$  edges. Therefore the extension of  $Z$  is at most  $g(2r-2)$  and the claim follows.  $\diamond$

Combining the above claim, [Observation 4.60](#), and [Theorem 4.4](#), we infer that either  $G$  contains a  $\theta_r$ -major  $M$  with at most  $f(r) \cdot \log k$  edges, or it contains a minor with minimum degree at least  $\frac{1}{r-1} \cdot 2^{\frac{f(r) \log k - 5r}{4r(2g(2r-2)+1)}} \geq k(r+1) - 1$ . If the second case is true, then by [Theorem 4.23](#),  $G$  contains  $k \cdot \theta_r$  as a minor, which proves the inductive step. We now consider the first case. Because  $M$  is 2-connected, we obtain that  $|V(M)| \leq |E(M)|$ . Therefore,  $|V(M)| \leq |E(M)| \leq f(r) \cdot \log k$  and we can bound the treewidth of the graph  $G' = G \setminus V(M)$  as follows:

$$\begin{aligned} \text{tw}(G') &\geq \text{tw}(G) - |V(M)| \\ &\geq f(r) \cdot k \cdot \log(k+1) - f(r) \cdot \log k \\ &\geq f(r) \cdot k \cdot \log k - f(r) \cdot \log k \\ &= f(r) \cdot (k-1) \cdot \log k. \end{aligned}$$

Then, from the induction hypothesis,  $G'$  contains a  $(k-1) \cdot \theta_r$ -major  $M'$  and obviously  $M \cup M'$  is a  $k \cdot \theta_r$ -major in  $G$ , which concludes our proof.  $\square$

[Theorem 4.32](#) implies that for every fixed  $r$ , it holds that every graph excluding  $k \cdot \theta_r$  as a minor has treewidth  $O(k \cdot \log k)$ . We give in [Subsection 4.3.7](#) a lemma indicating that this bound is tight up to the constants hidden in the  $O$ -notation.

### 4.3.7 Lower bounds

An natural question is whether the aforementioned results for the classes  $\{\mathcal{H}_k^i\}_{i \in [1,4]}$  are tight. In this section we provide lower bounds on the function upper-bounding the treewidth of graphs excluding members of these classes.

**Lemma 4.62.** *There is a sequence  $(G_k)_{k \in \mathbb{N}_{\geq 3}}$  of graphs such that:*

- $G_k$  does not contain  $W_k^2$  as a minor, for every  $k \in \mathbb{N}_{\geq 3}$ ; and
- $\text{tw}(G_k) = \Omega(k)$ .

*Proof.* The sequence  $(K_{k-1})_{k \in \mathbb{N}_{\geq 3}}$  satisfies the above properties.  $\square$

As  $W_k \leq_m W_k^2$  for every  $k \in \mathbb{N}_{\geq 3}$ , [Lemma 4.62](#) implies that [\(4.2\)](#) does not hold for the classes  $\mathcal{H}_k^1$  and  $\mathcal{H}_k^2$  with a function  $f$  such that  $f(k) = o(k)$ .

**Corollary 4.63.** *The bound given in [Theorem 4.27](#) is tight up to a constant factor.*

**Lemma 4.64.** *There is a sequence  $(G_i)_{i \in \mathbb{N}_{\geq 1}}$  of graphs and an increasing sequence  $(k_i)_{i \in \mathbb{N}_{\geq 1}}$  of integers such that:*

- $G_i$  does not contain  $k_i \cdot K_3$  as a minor, for every  $i \in \mathbb{N}_{\geq 1}$ ; and

- $\text{tw}(G_i) = \Omega(k_i \log k_i)$ .

*Proof.* According to [Mor94, Theorem 5.13], there is an infinite family  $\{G_i\}_{i \in \mathbb{N}}$  of 3-regular Ramanujan graphs  $G_i$  such that  $i \mapsto |G_i|$  is an increasing function. Furthermore, for every  $i \in \mathbb{N}$ , the graph  $G_i$  has girth at least  $\frac{2}{3} \log |V(G_i)|$  ([Mor94, Theorem 5.13]) and satisfies  $\text{tw}(G_i) = \Omega(|V(G_i)|)$  (see [BEM<sup>+</sup>04, Corollary 1]). For every  $i \in \mathbb{N}$ , let  $k_i$  be the minimum integer such that  $|V(G_i)| < k_i \cdot \frac{2}{3} \log |V(G_i)|$ . Observe that  $(k_i)_{i \in \mathbb{N}}$  is increasing. Notice that  $|V(G_i)| = \Omega(k_i \cdot \log k_i)$ , and thus  $\text{tw}(G_i) = \Omega(k_i \cdot \log k_i)$ . We will show that  $G_i$  does not contain  $k_i$  vertex-disjoint cycles, which implies that  $k_i \cdot \theta_r$  is not a minor of  $G_i$ , for every  $r \in \mathbb{N}_{\geq 2}$ . Suppose for contradiction that  $G_i$  contains  $k_i$  vertex-disjoint cycles. As the girth of  $G_i$  is at least  $\frac{2}{3} \log |V(G_i)|$ , each of these cycles has at least  $\frac{2}{3} \log |V(G_i)|$  vertices. Therefore  $G$  should contain at least  $k \cdot \frac{2}{3} \log |V(G_i)|$  vertices. This implies that  $|V(G)| \geq k \cdot \frac{2}{3} \log |V(G_i)| > |V(G_i)|$ , a contradiction. Therefore  $(k_i)_{i \in \mathbb{N}}$  and  $(G_i)_{i \in \mathbb{N}}$  satisfy the required properties.  $\square$

As every graph of  $\mathcal{H}_{2k}^0 \cup \mathcal{H}_{2k}^3 \cup \mathcal{H}_{k,r}^4$  contains  $k \cdot K_3$  as a minor (for every  $r \in \mathbb{N}_{\geq 2}$ ), we deduce that (4.2) does not hold for the classes  $\mathcal{H}_k^0$ ,  $\mathcal{H}_k^3$ , and  $\mathcal{H}_k^4$  with a function  $f$  such that  $f(k) = o(k \log k)$ .

**Corollary 4.65.** *For every fixed  $r \in \mathbb{N}_{\geq 2}$ , the bound given in Theorem 4.32 is tight up to a constant factor.*

## 4.4 Immersions of planar subcubic graphs in graphs of large tree-cut width

This section contains the proof of the following result.

**Theorem 4.66.** *There is a function  $f : \mathbb{N} \rightarrow \mathbb{N}$  where  $f(h) = O(h^{29} \text{polylog}(h))$  such that for every planar subcubic graph  $H$  with  $h$  edges and every graph  $G$ , if  $\text{tcw}(G) \geq f(h)$  then  $H$  is an immersion of  $G$ .*

This theorem is a consequence of the polynomial bounds of the Grid Exclusion Theorem of Chekuri and Chuzhoy (Theorem 4.26) and the following exclusion theorem for walls. We present it here as we will use it in Chapter 5.

**Theorem 4.67** ([Wol15, Theorem 7]). *Let  $G$  be a graph and  $r \geq 1$  be a positive integer. Let  $f$  be a function for which Theorem 4.1 (the grid exclusion theorem) holds. If  $G$  has tree-cut width at least  $4r^{10} f(r)$ , then  $G$  admits an immersion of the  $r$ -wall.*

We will also use the following ingredient.

**Lemma 4.68** ([Kan96]). *Every simple planar subcubic graph of  $n$  vertices is a topological minor of the  $\lfloor \frac{n}{2} \rfloor$ -grid.*

According to Theorem 4.67, graphs of large enough tree-cut width contain a large wall as an immersion. Therefore we only need to show that every planar subcubic graph is an immersion of a (large enough) wall in order to prove Theorem 4.66. For

this we use a supergraph of the wall defined as follows from the wall  $W_k$ . Let  $E = \{e, e \in E(P_j^{(v)}) \cap E(P_i^{(h)}), j \in \{1, 2, \dots, k\}, i \in \{1, 2, \dots, k + 1\}\}$ . We obtain  $\widehat{W}_k$  from  $W_k$  by adding a second copy of every edge in  $E$ . (For an example, see Figure 4.7.)




Figure 4.7: The graph  $\widehat{W}_5$ .

The next observation is a formal statement of what is depicted on Figure 4.8:  $\widehat{W}_n$  contains the  $(n \times n)$  grid  $\Gamma_n$  as a strong immersion. Branch vertices are depicted by white nodes and horizontal (respectively vertical) paths use the color green (respectively red).




Figure 4.8: Finding  $\Gamma_5$  as a strong immersion in  $\widehat{W}_5$ .

*Observation 4.69.* Let  $k \geq 2$  be an integer. If we define  $\phi$  and  $\psi$  with domains  $V(\Gamma_n)$  and  $E(\Gamma_n)$ , respectively, as follows:

$$\begin{aligned} \phi((i, j)) &= (i, 2j - 1) \\ \psi(\{(i, j), (i, j + 1)\}) &= (i, 2j - 1)(i, 2j)(i, 2(j + 1) - 1) \\ \psi(\{(i, j), (i + 1, j)\}) &= (i, 2j - 1)(i + 1, 2j - 1) \text{ for odd } i \\ \psi(\{(i, j), (i + 1, j)\}) &= (i, 2j - 1)(i, 2j)(i + 1, 2j)(i + 1, 2j - 1) \text{ for even } i, \end{aligned}$$

then  $(\phi, \psi)$  is a  $\Gamma_k$ -strong-immersion model in  $\widehat{W}_k$  (where we assume that  $\Gamma_k$  has vertex set  $\llbracket 1, k \rrbracket^2$ ).

The next result is mentioned in [Tho88] but not proof is provided.

**Lemma 4.70.** *Every planar subcubic graph on  $n$ -vertices is a topological minor of the wall  $W_n$ .*

*Proof.* Let  $H$  be a planar subcubic graph on  $n$  vertices. The proof goes as follows: we first construct a subdivision  $H'$  of  $H$  that is a simple graph. Then we prove that  $H'$  is a strong-immersion of  $\widehat{W}_n$  and obtain the following ordering:

$$H \leq_{\text{t.m.}} H' \leq_{\text{t.m.}} \Gamma_n \leq_{\text{s.im.}} \widehat{W}_n. \quad (4.4)$$

Finally, we construct a subdivision of  $H'$  in  $\widehat{W}_n$ . This is simple, hence it is a subgraph of  $W_n$ , as required.

Let  $H$  be a planar subcubic graph and let  $H'$  be the simple subcubic planar graph obtained from  $H$  by subdividing all but one edges of every multiedge. Notice that the first inequality of equation (4.4) is satisfied. Let us count how many vertices are added during the construction of  $H'$ . As  $H$  is subcubic, among the edges incident to a given vertex, at most two are being subdivided. That way we count each subdivided edge twice (once for each of its endpoints), hence we get:

$$|V(H')| \leq 2|V(H)|.$$

According to Lemma 4.68,  $H'$  is a topological minor of  $\Gamma_n$ : this gives the second inequality of the equation. Observation 4.69 gives the third inequality.

Let  $(\phi_1, \psi_1)$  be an  $H'$ -topological minor model in  $\Gamma_n$  and let  $(\phi_2, \psi_2)$  be the  $\Gamma_n$ -strong-immersion model in  $\widehat{W}_n$  given by Observation 4.69. These two models can be used to construct an  $H'$ -strong immersion model  $(\phi, \psi)$  in  $\widehat{W}_n$ , as the composition of  $(\phi_1, \psi_1)$  and  $(\phi_2, \psi_2)$ : for every  $v \in V(H')$ ,  $\phi(v) = \phi_2(\phi_1(v))$  and for every  $e \in E(H')$ ,  $\psi(e)$  is the concatenation of the paths obtained by applying  $\psi_2$  to the edges of the path  $\psi_1(e)$  (taken in the same order as they appear in this path). Let us call *expansion* of an immersion (respectively topological minor) model  $(\mu, \mu')$  of a graph  $J$  the graph  $(\bigcup_{e \in E(J)} V(\mu'(e)), \bigcup_{e \in E(J)} E(\mu'(e)))$ . Intuitively, this is the graph described by the model. Observe that the model  $(\phi, \psi)$  satisfies the following properties:

- the expansion of  $(\phi, \psi)$  is a subgraph of the expansion of  $(\phi_2, \psi_2)$ ; and
- the branch vertices of  $(\phi, \psi)$  are branch vertices of  $(\phi_2, \psi_2)$ .

We provide the following diagram to recall the roles of the different models we use (topological minor models are indicated by double arrows and strong immersion models by simple ones).

$$\begin{array}{ccc}
H' & \xrightarrow{(\phi_1, \psi_1)} & \Gamma_n \\
& \searrow (\phi, \psi) & \downarrow (\phi_2, \psi_2) \\
& & \widehat{W}_n
\end{array}$$

Let us show the following claim.

*Claim 4.71.* Let  $e, f \in E(H')$ . If  $v$  is an internal vertex of both  $\psi(e)$  and  $\psi(f)$ , then these paths also share an endpoint, which is adjacent to  $v$ .

If  $\psi(e)$  and  $\psi(f)$  share an internal vertex  $v$ , there are two edges  $a \in \psi_1(e)$  and  $b \in \psi_1(f)$  such that both  $\psi_2(a)$  and  $\psi_2(b)$  contain  $v$ . By definition of  $(\phi_2, \psi_2)$ , such a situation occurs only if  $a = \{(i, j), (i + 1, j)\}$  (for even  $i$ ) and  $b = \{(i, j), (i, j + 1)\}$  or  $b = \{(i + 1, j), (i + 1, j + 1)\}$ , for some even  $i \in \llbracket 1, n \rrbracket$  and some  $j \in \llbracket 1, n \rrbracket$  (see [Figure 4.8](#)). Observe that in both cases  $a$  and  $b$  share an endpoint. As  $(\phi_1, \psi_1)$  is a topological minor model,  $\psi_1(e)$  and  $\psi_1(f)$  may meet on endpoints only. Therefore the common endpoint of  $a$  and  $b$  is an endpoint of both  $\psi_1(e)$  and  $\psi_1(f)$ , hence  $\psi(e)$  and  $\psi(f)$  have a common endpoint. This proves the first part of the claim. The second part is now clear from the definition of  $(\phi_2, \psi_2)$ , as we know that the paths  $\psi_1(e)$  and  $\psi_1(f)$  start from the same vertex, one with a “vertical” edge, the other with a “horizontal” edge (see [Figure 4.8](#)).  $\diamond$

If  $(\phi, \psi)$ , which is a strong immersion model, is a topological minor model, then we can directly jump to the next step. Otherwise, according to [Claim 4.71](#), there are two edges  $e = \{u, v\}, f = \{u, w\}$  of  $H$  and vertices  $x, y \in \widehat{W}_n$  such that  $\psi(e)$  and  $\psi(f)$  both start with  $x = \phi(u)$  followed by  $y$ . Hence  $\{x, y\}$  is a double edge of  $\widehat{W}_n$ . As  $(\phi, \psi)$  is a strong immersion model of a subcubic graph,  $x$  has degree at most three in the expansion of  $(\phi, \psi)$ . We can therefore modify  $(\phi, \psi)$  as follows: we set  $\phi(u) = x$  and we shorten  $\psi(\{u, v\})$  and  $\psi(\{u, w\})$  by removing the edge  $\{x, y\}$  from each of them. In the case where there is a third vertex  $t \in V(H) \setminus \{v, w\}$  adjacent to  $u$ , we also extend the path  $\psi(\{t, u\})$  by adding the edge  $\{x, y\}$ . See [Figure 4.9](#) for an example.




Figure 4.9: Swapping branch vertices.

It is easy to see that by applying these changes we still get an  $H'$ -strong immersion model, with less crossings of certifying paths. By repeatedly applying these steps we eventually obtain a  $H'$ -topological minor model in  $\widehat{W}_n^+$ . Notice that its expansion is a simple graph, as  $H'$  is a simple graph. Therefore this expansion is also a subgraph of  $W_n$ . We proved that  $H'$  is a topological minor of  $W_n$ . It follows that the same holds for  $H$  and we are done.  $\square$

The proof of [Theorem 4.66](#) is now straightforward: given that  $\text{tcw}(G) \geq 4h^{10} f_1(h)$  (recall that  $f_1$  is the function of the Grid Exclusion Theorem, [Theorem 4.26](#)),  $G$  contains  $W_h$  as an immersion ([Theorem 4.67](#)). As we just proved with [Lemma 4.70](#), every planar subcubic graph on  $h$  vertices is an immersion of  $W_n$ , so it is an immersion of  $G$ . Also,  $4h^{10} f_1(h) = O(h^{29} \text{polylog}(h))$ .

# CHAPTER 5

---

## THE ERDŐS-PÓSA PROPERTY

In this chapter, we present general techniques for proving that a class of graphs have the Erdős–Pósa property. The first one is based on tree-like decompositions. Initially used with tree-decompositions, it can in fact also be applied to other decompositions as tree-partitions and tree-cut decompositions. The second technique is related to the notion of girth and its extensions. We then apply these techniques to obtain new Erdős–Pósa-type results. In particular, we shed some light on the edge variant of the Erdős–Pósa property. In the remaining sections, we provide negative results on the Erdős–Pósa property of classes defined by containment relations and we present a collection of tables listing the known results on the Erdős–Pósa property.

- [RST16] *An edge variant of the Erdős–Pósa property*, co-authored with Ignasi Sau and Dimitrios M. Thilikos, *Discrete Mathematics* (2016);
- [CRST15b] *An  $O(\log \text{OPT})$ -approximation for covering/packing minor models of  $\theta_r$* , co-authored with Dimitris Chatzidimitriou, Ignasi Sau, and Dimitrios M. Thilikos, *WAOA* 2015;
- [GKRT16] *Packing and covering immersion models of planar subcubic graphs*, co-authored with Archontia Giannopoulou, O-joung Kwon, and Dimitrios M. Thilikos, *WG* 2016;
- [RT16] *Recent techniques and results on the Erdős–Pósa property*, co-authored with Dimitrios M. Thilikos, submitted (2016).

### 5.1 Introduction

A considerable part of Combinatorics has been developed around min-max theorems. Min-max theorems usually identify dualities between certain objects in graphs, hypergraphs, and other combinatorial structures. The target is to prove that the absence of the primal object implies the presence of the dual one and vice versa.

A classic example of such a duality is Menger’s theorem: the primal concept is the existence of  $k$  internally disjoint paths between two vertex sets  $S$  and  $T$  of a graph

$G$ , while the dual concept is a collection of  $k$  vertices that intersect all  $(S, T)$ -paths. Another example is König's theorem where the primal notion is the existence of a matching of  $k$  vertices in a bipartite graph and the dual one is the existence of a vertex cover of size  $k$ . It is also known that, in case of general graphs, this duality becomes an approximate one, i.e., a vertex cover of size  $2k$ . In both aforementioned examples, the duality is between the notions of *packing* and *covering* of a collection  $\mathcal{C}$  of combinatorial objects of a graph. In Menger's theorem  $\mathcal{C}$  consists of all  $(S, T)$ -paths of  $G$  while in König's theorem  $\mathcal{C}$  is the set of all edges of  $G$ . That way, both aforementioned min-max theorems can be stated, for some class of graphs  $\mathcal{G}$  (called *host class*) and some gap function  $f : \mathbb{N} \rightarrow \mathbb{N}$ , as follows:

*For every graph  $G$  in  $\mathcal{G}$ , either  $G$  contains  $k$ -vertex disjoint objects in  $\mathcal{C}$  or it contains  $f(k)$  vertices intersecting all objects in  $\mathcal{C}$  that appear in  $G$ .*

Clearly, for the case of Menger's theorem the host class is the class of all graphs while in the case of König's theorem the host class is restricted to the class of bipartite graphs. In both cases the derived duality is an *exact* one in the sense that  $f$  is the identity function. However, this is not the case if we want to extend the duality of König's theorem in the case of all graphs, where we can consider  $f : k \mapsto 2k$  (i.e., we have an *approximate* duality).

One of the most celebrated results about packing/covering dualities was obtained by Paul Erdős and Lajos Pósa in 1965 where the object to cover and pack was the set of all cycles of  $G$  [EP65]. In this case the host class contains all graphs, while  $f : k \mapsto O(k \cdot \log k)$ . Moreover, Erdős and Pósa proved that this gap is optimal in the sense that it cannot be improved to a function  $f : k \mapsto o(k \cdot \log k)$ . This result motivated a long line of research for min-max dualities that are not necessarily exact. Since then, a multitude of results on Erdős–Pósa properties have appeared for several combinatorial objects, including extensions to digraphs [LY78, Sey96, RRST96b, HM13, GT10], rooted graphs [KKK12, PW12, Joo14, BJS14], labeled graphs [KW05], signed graphs [HNP06, ADG04], hypergraphs [Alo02, Bou13, BT15], matroids [GK09], and other combinatorial structures [GL69] (see [Ree97] for a survey on this topic). Also it is worth stressing that Erdős–Pósa dualities have been useful in more applied domains. For example, in bioinformatics where they were useful for upper-bounding the number of fixed-points of a boolean networks [Ara08, ADG04, ARS16].

The chapter is organized as follows. We first describe some recent techniques for proving Erdős–Pósa properties, mainly based on techniques related to tree-like decompositions of graphs (Subsection 5.3.1 and Subsection 5.3.2). We focused our presentation to the description of general frameworks that, we believe, might be useful for further investigations. We then present several results related to the Erdős–Pósa property. Lastly, in Subsection 5.6.2, we provide an extensive update of results on Erdős–Pósa properties, reflecting the current progress on this vibrant area of graph theory. Most of the notions used in this chapter are defined in Section 5.2.

## 5.2 Notions related to the Erdős–Pósa property

Let us introduce definitions related to the topic of this chapter and that will also be used in the next chapter.

Most of the definitions we give here have two variants: one is related to vertices and the other one is related to edges. In order to make the definitions more concise, we use symbols  $v$  and  $e$  in order to distinguish the vertex and the edge variants of the properties/parameters that we are dealing with. For instance, if  $x \in \{v, e\}$ , and  $G$  is a graph, we set  $A_x(G) = V(G)$  if  $x = v$  and  $A_x(G) = E(G)$  if  $x = e$ . Similarly,  $x$ -disjoint stands for *vertex-disjoint* when  $x = v$  and for *edge-disjoint* when  $x = e$ .

**Packing and covering.** Let  $\mathcal{H}$  be a family of graphs and let  $x \in \{v, e\}$ . An  $x$ - $\mathcal{H}$ -cover of  $G$  is a set  $C \subseteq A_x(G)$  such that  $G \setminus C$  does not contain any subgraph isomorphic to a member of  $\mathcal{H}$ . An  $x$ - $\mathcal{H}$ -packing in  $G$  is a collection of  $x$ -disjoint subgraphs of  $G$ , each being isomorphic to some graph of  $\mathcal{H}$ .

We denote by  $x$ - $\text{pack}_{\mathcal{H}}(G)$  the maximum size of an  $x$ - $\mathcal{H}$ -packing, which we call *packing number* (with respect to  $\mathcal{H}$ ) and by  $x$ - $\text{cover}_{\mathcal{H}}(G)$  the minimum size of an  $x$ - $\mathcal{H}$ -covering in  $G$ , also referred to as *covering number* (also defined relatively to  $\mathcal{H}$ ).

There is an easy inequality between these two parameters.

*Remark 5.1.* For every  $x \in \{v, e\}$ , for every graph class  $\mathcal{H}$ , and every graph  $G$ , the following holds:

$$x\text{-pack}_{\mathcal{H}}(G) \leq x\text{-cover}_{\mathcal{H}}(G).$$

Indeed, any  $x$ - $\mathcal{H}$ -cover must contain at least one vertex (if  $x = v$ ) or edge (if  $x = e$ ) of each element of an  $x$ - $\mathcal{H}$ -packing of maximum size.

The Erdős–Pósa property is concerned with the other direction, that is, bounding the covering number in terms of the packing number.

### The Erdős–Pósa property.

**Definition 5.2.** Let  $\mathcal{G}$  and  $\mathcal{H}$  be two graph classes and let  $x \in \{v, e\}$ . We say that  $\mathcal{H}$  has the  $x$ -Erdős–Pósa property for  $\mathcal{G}$  if there is a function  $f: \mathbb{N} \rightarrow \mathbb{N}$  such that the following holds:

$$\forall G \in \mathcal{G}, x\text{-cover}_{\mathcal{H}}(G) \leq f(x\text{-pack}_{\mathcal{H}}(G)).$$

Any function  $f$  satisfying the above inequality is called a *gap* of the  $x$ -Erdős–Pósa property of  $\mathcal{H}$  for  $\mathcal{G}$ . When a class of graphs has the  $x$ -Erdős–Pósa-property for the class of finite graphs, we simply say that it has the  $x$ -Erdős–Pósa-property. We usually refer to  $\mathcal{G}$  as the *host* graph class and by  $\mathcal{H}$  as the *guest* graph class.

## 5.3 General tools for proving Erdős–Pósa type results

In this section we present general techniques for proving Erdős–Pósa type results. They are either based on tree-like decompositions of graphs, that allow to find small separators, or on variants of the girth, which certify that the graph contains a small subgraph of the desired type.

### 5.3.1 The Erdős–Pósa property from graph decompositions

Let  $\mathcal{H}$  be a graph class,  $\mathbf{p}$  be a graph parameter, and  $x \in \{v, e\}$ . We say that a function  $f: \mathbb{N} \rightarrow \mathbb{N}$  is a *ceiling* for the triple  $(\mathbf{p}, \mathcal{H}, x)$  if for every graph  $G$ ,  $\mathbf{p}(G) \leq f(x\text{-pack}_{\mathcal{H}}(G))$ . Intuitively, there is a ceiling for the triple  $(\mathbf{p}, \mathcal{H}, x)$  if a large value of  $\mathbf{p}$  on a graph forces a large  $x$ -packing of elements of  $\mathcal{H}$ . Notice that every ceiling for  $(\mathbf{p}, \mathcal{H}, v)$  is a ceiling for  $(\mathbf{p}, \mathcal{H}, e)$ , since a vertex-disjoint packing is a special case of an edge-disjoint packing.

Given a graph parameter  $\mathbf{p}$  and an integer  $k$ , we denote

$$\mathcal{G}_{\mathbf{p} \leq k} = \{G, \mathbf{p}(G) \leq k\}.$$

**Theorem 5.3.** *Let  $\mathcal{H}$  be a class of graphs,  $x \in \{v, e\}$ ,  $\mathbf{p}$  be a graph parameter, let  $f: \mathbb{N} \rightarrow \mathbb{N}$  be a function and let  $h_r: \mathbb{N} \rightarrow \mathbb{N}$  be a function, for every  $r \in \mathbb{N}$ . Suppose that the following two conditions hold:*

- A.  *$f$  is a ceiling for the triple  $(\mathbf{p}, \mathcal{H}, x)$ ;*
- B. *for every  $r \in \mathbb{N}$ ,  $\mathcal{H}$  has the  $x$ -Erdős–Pósa property for  $\mathcal{G}_{\mathbf{p} \leq r}$  with gap  $h_r$ ;*

*then  $\mathcal{H}$  has the  $x$ -Erdős–Pósa property with gap  $k \mapsto h_{f(k)}(k)$ .*

*Proof.* Let  $G$  be a graph and let  $k = x\text{-pack}_{\mathcal{H}}(G)$ . We have  $\mathbf{p}(G) \leq f(k)$ , by definition of a ceiling. Therefore,  $G \in \mathcal{G}_{\mathbf{p} \leq f(k)}$ , and thus  $x\text{-cover}_{\mathcal{H}}(G) \leq h_{f(k)}(k)$ .  $\square$

**Theorem 5.3** will be used as a master theorem for the results of this section.

#### Vertex version and tree decompositions

In a breakthrough paper [CC13], Chekuri and Chuzhoy proved that every graph of large treewidth can be partitioned into several subgraphs of large treewidth, with a polynomial dependency between the treewidth of the original graph, the one of the subgraphs, and the number of subgraphs. In fact they proved the two next results.

**Theorem 5.4** ([CC13, Theorem 1.1]). *There is a non-decreasing function  $f_2: \mathbb{R} \rightarrow \mathbb{R}$  with  $f_2(t) = \text{polylog } t$  such that, for every graph  $G$  and every positive integers  $h$  and  $p$ , if*

$$hp^2 \leq \frac{\text{tw}(G)}{f_2(\text{tw}(G))},$$

*then there is a partition  $G_1, \dots, G_h$  of  $G$  into vertex-disjoint subgraphs such that  $\text{tw}(G_i) \geq p$  for every  $i \in \llbracket 1, h \rrbracket$ .*

**Theorem 5.5** ([CC13, Theorem 1.2]). *There is a non-decreasing function  $f_3: \mathbb{R} \rightarrow \mathbb{R}$  with  $f_3(t) = \text{polylog } t$  such that, for every graph  $G$  and every positive integers  $h$  and  $p$ , if*

$$h^3 p \leq \frac{\text{tw}(G)}{f_3(\text{tw}(G))},$$

*then there is a partition  $G_1, \dots, G_h$  of  $G$  into vertex-disjoint subgraphs such that  $\text{tw}(G_i) \geq p$  for every  $i \in \llbracket 1, h \rrbracket$ .*

These results have been used to obtain ceilings. In fact, [CC13] also contains the following result.

**Lemma 5.6** ([CC13, from the proof of Theorem 5.4]). *If  $r$  is an integer and  $\mathcal{H}$  is a class of graphs such that every graph of treewidth at least  $r$  contains an  $\mathcal{H}$ -subgraph, then there is a ceiling  $f_4$  for  $(\text{tw}, \mathcal{H}, \nu)$  such that  $f_4(k) = kr^2 \text{polylog}(kr)$ .*

Let us now see the role of ceilings with respect to the Erdős–Pósa property. A function  $f: \mathbb{R} \rightarrow \mathbb{R}$  is said to be *superadditive* if  $f(x) + f(y) \leq f(x + y)$  for every pair  $x, y$  of positive reals. The main argument of the next lemma has been first used in [FST11]. Different statements of this results appeared in [CC13, RST16, CRST15b].

**Lemma 5.7.** *Let  $\mathcal{H}$  be a family of connected graphs. If  $f$  is a superadditive ceiling for  $(\text{tw}, \mathcal{H}, \nu)$  then  $\mathcal{H}$  has the  $\nu$ -Erdős–Pósa property with gap  $k \mapsto 5 \cdot f(k) \log(k + 1)$ .*

*Proof.* Let us show the following for every integer  $k$ : for every graph  $G$ , if  $\nu\text{-pack}_{\mathcal{H}}(G) = k$  then  $\nu\text{-cover}_{\mathcal{H}} \leq 5f(k) \log(k + 1)$ . The proof is by induction on  $k$ . The base case  $k = 0$  is trivial. Let  $k > 0$ , and let us assume that the above statement holds for every positive integer  $k' < k$  (induction hypothesis).

Let  $G$  be a graph such that  $\nu\text{-pack}_{\mathcal{H}}(G) = k$ . We will rely on the following claim.

*Claim 5.8.* There is a separation  $(A, B)$  of order  $\text{tw}(G) + 1$  of  $G$  such that

$$k/3 \leq \nu\text{-pack}_{\mathcal{H}}(G[A \setminus B]) \leq 2k/3.$$

*Proof.* It is known that every graph  $G$  has a nice tree decomposition with width  $\text{tw}(G)$  [Klo94a]. We therefore can assume that  $(T, r, (X_t)_{t \in V(T)})$  is a nice tree decomposition of  $G$  of optimal width. We define

$$G_t = G \left[ \bigcup_{s \in \text{desc}_{(T,r)}(t)} X_s \right] \quad \text{and} \quad G_t^- = G_t \setminus X_t.$$

Let  $t$  be a vertex of  $T$  at minimal distance from a leaf subject to the requirement  $\nu\text{-pack}_{\mathcal{H}}(G_t^-) > 2k/3$ . Such a vertex exists, as  $\nu\text{-pack}_{\mathcal{H}}(G_r^-) = \nu\text{-pack}_{\mathcal{H}}(G_r) = k$ . Observe that  $t$  is either a forget node, or a join node. Indeed, for every base node  $u$  we have  $\nu\text{-pack}_{\mathcal{H}}(G_u^-) = 0$ . Moreover, every introduce node  $u$  with child  $v$  satisfies  $\nu\text{-pack}_{\mathcal{H}}(G_u^-) = \nu\text{-pack}_{\mathcal{H}}(G_v^-)$ , since  $G_u^- = G_v^-$ .

*First case:*  $t$  is a forget node with child  $u$ . We set  $A = V(G_u)$  and  $B = V(G) \setminus V(G_u^-)$ .

*Second case:*  $t$  is a join node with children  $u_1, u_2$ . We set  $A = V(G_{u_i})$  and  $B = V(G) \setminus V(G_{u_i}^-)$ , where  $u_i$  is a child of  $t$  such that  $\nu\text{-pack}_{\mathcal{H}}(G_{u_i}^-) \geq k/3$ . Such child exist because  $\nu\text{-pack}_{\mathcal{H}}(G_t^-) = \nu\text{-pack}_{\mathcal{H}}(G_{u_1}^-) + \nu\text{-pack}_{\mathcal{H}}(G_{u_2}^-)$  (as  $t$  is a join node) and  $\nu\text{-pack}_{\mathcal{H}}(G_t^-) > 2k/3$ , by definition of  $t$ .

It is clear that in both cases  $(A, B)$  is a separation of order  $\text{tw}(G)+1$ . The inequality  $\text{v-pack}_{\mathcal{H}}(G[A]) \leq 2k/3$  follows from the definition of  $t$ . In the first case, we have

$$\begin{aligned} \text{v-pack}_{\mathcal{H}}(G[A \setminus B]) &= \text{v-pack}_{\mathcal{H}}(G_u^-) \\ &\geq \text{v-pack}_{\mathcal{H}}(G_t^-) - 1 \quad (\text{as } t \text{ is a forget node}) \\ &> \frac{2k}{3} - 1 \\ &\geq \frac{k}{3}. \end{aligned}$$

In the second case, the choice of  $i$  ensures that  $k/3 \leq \text{v-pack}_{\mathcal{H}}(G[A \setminus B])$ . □

Observe that  $\text{tw}(G) \leq f(k)$ , by definition of  $f$ . According to [Claim 5.8](#), there is a separation  $(A, B)$  of order  $\text{tw}(G) + 1$  in  $G$  such that  $k/3 \leq \text{v-pack}_{\mathcal{H}}(G[A \setminus B]) \leq 2k/3$ . Let  $k_A = \text{v-pack}_{\mathcal{H}}(G[A \setminus B])$  and  $k_B = \text{v-pack}_{\mathcal{H}}(G[B \setminus A])$ . It follows that  $k_A, k_B \leq \lfloor 2k/3 \rfloor$ .

We then have

$$\begin{aligned} \text{v-cover}_{\mathcal{H}}(G) &\leq \text{v-cover}_{\mathcal{H}}(G[A \setminus B]) + \text{v-cover}_{\mathcal{H}}(G[B \setminus A]) + |A \cap B| \\ &\leq \text{v-cover}_{\mathcal{H}}(G[A \setminus B]) + \text{v-cover}_{\mathcal{H}}(G[B \setminus A]) + f(k) + 1 \\ &\leq 5f(k_A) \log(k_A + 1) + 5f(k_B) \log(k_B + 1) + f(k) + 1. \end{aligned}$$

The first inequality follows from the fact that the graphs in  $\mathcal{H}$  are connected. Indeed, a connected subgraph of  $G \setminus (A \cap B)$  belong to exactly one of  $G[A \setminus B]$  and  $G[B \setminus A]$ , so we can consider these two parts of  $G \setminus (A \cap B)$  independently. Notice that in the case where  $k = 1$ , we get  $k_A = k_B = 0$  and we have  $\text{v-cover}_{\mathcal{H}}(G) \leq f(k) \leq 3f(k) \log(k+1)$ . Therefore we now assume  $k \geq 2$ . We can deduce from  $k_A, k_B \leq \lfloor \frac{2}{3}k \rfloor$  that  $k_A + 1 \leq \frac{3}{4}(k+1)$  and  $k_B + 1 \leq \frac{3}{4}(k+1)$ .

$$\begin{aligned} \text{v-cover}_{\mathcal{H}}(G) &\leq 5 \cdot (f(k_A) + f(k_B)) \log\left(\frac{3(k+1)}{4}\right) + f(k) + 1 \\ &\leq 5 \cdot f(k) \log\left(\frac{3(k+1)}{4}\right) + f(k) + 1 \quad (\text{superadditivity of } f) \\ &\leq 5 \cdot f(k) \log(k+1) - 5 \cdot \log(4/3)f(k) + 2f(k) \\ &\leq 5 \cdot f(k) \log(k+1). \end{aligned} \quad \square$$

**Corollary 5.9** (see also [\[CC13\]](#) and [\[CC14\]](#)). *For every connected planar graph  $H$ , the class  $\mathcal{M}(H)$  has the v-Erdős–Pósa property with gap  $O(k \cdot h^{38} \cdot \text{polylog}(kh))$ , where  $h = V(H) + 2E(H)$ .*

For every connected planar graph  $H$ , [Corollary 5.9](#) provides a gap for  $\mathcal{M}(H)$  that is polynomial in both  $k$  and  $h = |V(H)| + 2|E(H)|$  ([\[RST94a, 1.5\]](#)). In [Subsection 5.4.1](#) we will apply [Lemma 5.6](#) to majors of specific planar graphs in order to obtain better gaps.

Notice that the proof of [Lemma 5.7](#) strongly relies on the fact that  $H$  is connected. The non-connected case requires some more ideas that are originating from [\[RS86\]](#) (also used for forests in [\[FJW13a\]](#)). We expose them hereafter. We will need the two next lemmas.

**Lemma 5.10** ([\[RS86\]](#)). *Let  $q, k$  be two positive integers, let  $T$  be a tree and let  $\mathcal{A}_1, \dots, \mathcal{A}_q$  be families of subtrees of  $T$ . Assume that for every  $i \in \llbracket 1, q \rrbracket$ , there are  $kq$  elements of  $\mathcal{A}_i$  that are pairwise vertex-disjoint. Then for every  $i \in \llbracket 1, q \rrbracket$ , there are  $k$  elements  $T_1^i, \dots, T_k^i$  of  $\mathcal{A}_i$  such that*

$$T_1^1, \dots, T_k^1, T_1^2, \dots, T_k^2, \dots, T_1^q, \dots, T_k^q$$

*are all pairwise vertex-disjoint.*

The next lemma is the Erdős–Pósa property of subtrees of a tree. It can be obtained from the fact that subtrees of a tree have the Helly property.

**Lemma 5.11** (see [\[GL69\]](#)). *Let  $T$  be a tree and let  $\mathcal{A}$  be a collection of subtrees of  $T$ . For every positive integer  $k$ , either  $T$  has (at least)  $k$  vertex disjoint subtrees that belong to  $\mathcal{A}$ , or  $T$  has a subset  $X$  of less than  $k$  vertices such that no subtree of  $T \setminus X$  belongs to  $\mathcal{A}$ .*

We are now ready to deal with disconnected patterns.

**Lemma 5.12** ([\[RS86\]](#)). *Let  $w$  be a positive integer and let  $H$  be a graph on  $q$  connected components.  $\mathcal{M}(H)$  has the  $v$ -Erdős–Pósa property on the class of graphs of treewidth at most  $w$  with gap  $k \mapsto (w - 1)(kq - 1)$ .*

*Proof.* Let  $k$  be a positive integer. We want to show that either  $v\text{-pack}_{\mathcal{M}(H)}(G) \geq k$  or  $v\text{-cover}_{\mathcal{M}(H)}(G) \leq (w - 1)(kq - 1)$ . Let  $H_1, \dots, H_q$  be the connected components of  $H$ . Let  $(T, \mathcal{X})$  be a tree-decomposition of  $G$  of width  $w$ . For every subgraph  $F$  of  $G$ , we denote by  $T(F)$  the subgraph of  $T$  induced by the bags containing vertices of  $F$ . Notice that  $T(F)$  is connected if  $F$  is connected.

For every  $i \in \llbracket 1, q \rrbracket$ , we let  $\mathcal{H}_i$  be the class of subgraphs of  $G$  that are isomorphic to a graph in  $\mathcal{M}(H_i)$  and we consider the class  $\mathcal{T}_i = \{T(F), F \in \mathcal{H}_i\}$ .

If for every  $i \in \llbracket 1, q \rrbracket$ ,  $\mathcal{T}_i$  contains  $kq$  vertex-disjoint trees, then according to [Lemma 5.11](#) there are pairwise vertex-disjoint trees  $\{T_j^i\}_{i \in \llbracket 1, q \rrbracket, j \in \llbracket 1, k \rrbracket}$ . Observe that for every two subgraphs  $F, F'$  of  $G$ , if  $T(F)$  and  $T(F')$  are vertex-disjoint, then so are  $F$  and  $F'$ . Therefore  $G$  has pairwise vertex-disjoint subgraphs  $\{F_j^i\}_{i \in \llbracket 1, q \rrbracket, j \in \llbracket 1, k \rrbracket}$  such that  $F_j^i$  is isomorphic to an element of  $\mathcal{H}_i$  for every  $i \in \llbracket 1, q \rrbracket$  and  $j \in \llbracket 1, k \rrbracket$ . This proves that in this case,  $v\text{-pack}_{\mathcal{M}(H)}(G) \geq k$ .

We therefore now assume that the above condition does not hold, namely there is an index  $i \in \llbracket 1, q \rrbracket$  such that  $\mathcal{T}_i$  contains less than  $kq$  vertex-disjoint trees. [Lemma 5.11](#) implies the existence of a subset  $X$  with  $|X| \leq kq - 1$  such that  $T \setminus X$  is free from subtrees isomorphic to a member of  $\mathcal{T}_i$ . Let  $Y$  denote the union of the bags indexed by vertices in  $X$ . Observe that  $|Y| \leq (w - 1)|X| \leq (w - 1)(kq - 1)$ . The choice of  $Y$  ensures that  $G \setminus Y$  has no subgraph isomorphic to a member of  $\mathcal{H}_i$ . Hence  $v\text{-cover}_{\mathcal{M}(H_i)} \leq (w - 1)(kq - 1)$ . We deduce  $v\text{-cover}_{\mathcal{M}(H)} \leq (w - 1)(kq - 1)$ .  $\square$

**Corollary 5.13.** *For every planar graph  $H$  with  $q$  connected components, the class  $\mathcal{M}(H)$  has the Erdős–Pósa property with gap  $O(qk^2 \cdot h^{38} \cdot \text{polylog}(kh))$ , where  $h = V(H) + 2E(H)$ .*

## Edge version and tree partitions

In the edge variant of the Erdős–Pósa properties we use tree-partition width as a possible edge-analogue of treewidth.

Let  $\mathcal{H}$  be a class of graphs. We define  $\tilde{\mathcal{H}}$  as the set of all the *subgraph minimal elements* of  $\mathcal{H}$ , i.e.,

$$\tilde{\mathcal{H}} = \{H, H \in \mathcal{H} \text{ and none of the subgraphs of } H \text{ belongs to } \mathcal{H}\}.$$

We define  $\Delta(\mathcal{H})$  as the maximum number of edges incident to a vertex in a graph of  $\mathcal{H}$  (counting multiple edges). We also set  $\tilde{\Delta}(\mathcal{H}) = \Delta(\tilde{\mathcal{H}})$ .

*Observation 5.14.* For every graph  $H$  of  $h$  edges, it holds that  $\tilde{\Delta}(\mathcal{M}(H)) \leq h$ ,  $\tilde{\Delta}(\mathcal{T}(H)) \leq h$ ,  $\tilde{\Delta}(\mathcal{I}(H)) \leq 2h$ .

**Lemma 5.15.** *Let  $\mathcal{H}$  be a class of connected non-trivial graphs where  $\tilde{\Delta}(\mathcal{H}) \leq d$ . Then for every  $r \in \mathbb{N}$ ,  $\mathcal{H}$  has the e-Erdős–Pósa-property on  $\mathcal{G}_{\text{tpw} \leq r}$  with gap  $g_r(k) = k \cdot r \cdot (dr + 1)$ .*

*Proof.* Let  $r \in \mathbb{N}$ . We will show the following for every  $k \in \mathbb{N}$ : for every graph  $G \in \mathcal{G}_{\text{tpw} \leq r}$ , if  $\text{e-pack}_{\mathcal{H}}(G) = k$  then  $\text{e-cover}_{\mathcal{H}}(G) \leq g_r(k)$ .

We proceed by induction. The base case  $k = 0$  is trivial. We thus assume that  $k > 0$  and that the above statement holds for every positive integer  $k' < k$  (induction hypothesis).

Let  $G \in \mathcal{G}_{\text{tpw} \leq r}$  be a graph such that  $\text{e-pack}_{\mathcal{H}}(G) = k$ . We assume that  $G$  is connected, as otherwise we can treat each connected component separately.

Let  $(T, s, \{X_t\}_{t \in V(T)})$  be an optimal rooted tree partition decomposition of  $G$ . We define  $G_t = G \left[ \bigcup_{u \in \text{desc}_{(T,s)}(t)} X_u \right]$ . For every edge  $\{u, v\}$  of  $T$  we denote by  $E_{\{u,v\}}$  the edges of  $G$  with the one endpoint in  $X_u$  and the other one in  $X_v$ . Let  $t$  be a vertex of  $T$  of minimum distance from a leaf, subject to  $\text{e-pack}_{\mathcal{H}}(G_t) > 0$ .

Let  $M$  be a subgraph-minimal subgraph of  $G_t$  that is isomorphic to some member of  $\mathcal{H}$  and let  $t_1, \dots, t_p$  be the children of  $t$  such that  $V(G_{t_i}) \cap V(M) \neq \emptyset$  for every  $i \in \llbracket 1, p \rrbracket$ . By minimality of  $M$ , it has no vertex with more than  $\tilde{\Delta}(\mathcal{H}) \leq d$  incident edges. As  $|X_t| \leq r$ , we deduce that  $p \leq rd$ .

Let  $C = E(X_t) \cup \bigcup_{i=1}^p E_{\{t, u_i\}}$ . Notice that  $|C| \leq r + dr^2$ . Let us consider then graph  $G' = G \setminus C$ . Let  $M'$  be a subgraph of  $G'$  that is isomorphic to some member of  $\mathcal{H}$ . By minimality of  $t$ ,  $\text{e-pack}_{\mathcal{H}}(G_{t_i}) = 0$ , for every  $i \in \llbracket 1, p \rrbracket$ . Therefore, if  $M'$  contained an edge  $e \in E(G_{t_i})$  (for some  $i \in \llbracket 1, p \rrbracket$ ), it would also contain an edge of  $E(G) \setminus E(G_{t_i})$ . Since every graph of  $\mathcal{H}$  is connected,  $M'$  would also need to contain some edge of  $E_{t, u_i}$  in order to be connected to edges of  $E(G) \setminus E(G_{t_i})$ . However  $E(G') \cap E_{t, u_i} = \emptyset$ . We deduce that for every subgraph  $M'$  of  $G'$  that is isomorphic to some member of  $\mathcal{H}$ , we have  $E(M') \cap E(M) = \emptyset$ . It follows that every e- $\mathcal{H}$ -packing in  $G'$  is edge-disjoint with  $M$ .

Hence  $\text{e-pack}_{\mathcal{H}}(G') < k$ , as otherwise a packing of size  $k$  in  $G'$  would, together with  $M$ , yield a packing of size  $k + 1$  in  $G$  whereas  $\text{e-pack}_{\mathcal{H}}(G) = k$ . By applying the induction hypothesis on  $G'$ , there is a subset  $D \subseteq E(G')$  such that  $\text{e-pack}_{\mathcal{H}}(G' \setminus D) = 0$  and moreover  $|D| \leq g_r(k - 1)$ . It is easy to see that  $C \cup D$  is an e- $\mathcal{H}$ -cover of  $G$ . Furthermore  $|C \cup D| \leq r(dr + 1) + g_r(k - 1) = g_r(k)$ , as required.  $\square$

In this section we presented decomposition-based techniques aiming to prove the edge-Erdős–Pósa property, and towards this purpose we used tree-partition decompositions as a possible edge-counterpart to tree-decompositions, that are used in the vertex case. Let us briefly mention that there are other tree-like decompositions that deserve attention. For instance, tree-cut decompositions can be successfully used to deal with immersion expansions, as we will see in [Subsection 5.4.4](#). Applications of the techniques described in this section will be presented in [Section 5.4](#).

### 5.3.2 The Erdős–Pósa property from girth

In this section, we give a proof of the Erdős–Pósa Theorem that highlights a technique for proving more general Erdős–Pósa type results. The technique can be informally summarized as follow. We prove that either  $G$  contains a *small* cycle or that it can be reduced to a smaller graph with the same packing and covering number. We then apply induction on either the graph where a small cycle has been deleted (in the first case), or on the reduced graph (in the second case). This technique has been successfully applied in [\[FJW13a, CRST15b\]](#), for instance. Let us first recall the following result.

**Lemma 5.16** ([\[Tho83b\]](#), see also [\[Die05, Theorem 7.4.2\]](#)). *There is a constant  $c \in \mathbb{R}$ , such that for every  $q \in \mathbb{N}_{\geq 1}$ , every graph of minimum degree at least 3 and girth at least  $c \log q$  contains  $K_q$  as a minor.*

A direct consequence of this result is the following trichotomy.

**Corollary 5.17.** *For every graph  $G$  and every integer  $q > 1$ , one of the following holds:*

- (i)  $G$  has a cycle on at most  $c \log q$  vertices;
- (ii)  $G$  has a vertex of degree at most 2;
- (iii)  $G$  contains  $K_q$  as a minor,

where  $c$  is the constant of [Lemma 5.16](#).

We now prove the lemma that implies the classic Erdős–Pósa Theorem both for the vertex and its edge version. Recall that  $A_x(G)$  denotes  $V(G)$  or  $E(G)$ , depending if  $x = v$  or  $x = e$ .

**Lemma 5.18.** *For every  $q \in \mathbb{N}^+$  and every  $x \in \{v, e\}$ , the set  $\mathcal{M}(\theta_2)$  has the  $x$ -Erdős–Pósa property for the class of  $K_q$ -minor-free with gap  $O(k \cdot \log q)$ .*

*Proof.* We will prove that for every non-negative integer  $k$  and every  $K_q$ -minor-free graph  $G$ , either  $G$  has  $k$   $x$ -disjoint cycles, or  $G$  has a subset  $X \subseteq A_x(G)$  of size at most  $ck \log q$  such that  $G \setminus X$  is a forest, where  $c$  is the constant of [Lemma 5.16](#). We proceed by induction on the pair  $(k, G)$ , with the well-founded order defined by  $(k', G') \leq (k, G) \iff (k' \leq k \text{ and } |A_x(G')| \leq |A_x(G)|)$ , for all graphs  $G, G'$  and non-negative integers  $k, k'$ .

The base cases corresponding to  $k = 0$  or  $|A_x(G)| = 0$  are trivial. Let us now assume that  $k \geq 1$ ,  $|A_x(G)| \geq 1$ , and that the lemma holds for every pair  $(k', G')$  such that  $(k', G') \leq (k, G)$ .

According to [Corollary 5.17](#), either  $G_i$  has a cycle  $C$  on at most  $c \log q$  vertices, or it has a vertex  $v$  of degree at most two, or it contains  $K_q$  as a minor. The last case is not possible, as we require  $G$  to be  $K_q$ -minor-free.

Whenever the first case applies, we set  $G' = G \setminus A_x(C)$  and we consider the pair  $(k - 1, G')$ . If  $G'$  contains  $k - 1$   $x$ -disjoint cycles, then  $G$  contains  $k$   $x$ -disjoint cycles obtained by adding  $C$  to those of  $G'$  and we are done. Otherwise, the induction hypothesis implies the existence of a subset  $X' \subseteq A_x(G')$  with  $|X'| \leq c(k - 1) \log q$  such that  $G' \setminus X'$  is a forest. Then by definition of  $C$ ,  $X = X' \cup A_x(C)$  has size at most  $c \log q$  and  $G \setminus X$  is a forest, as required.

In the second case, we delete  $v$  if it is isolated and we contract an edge  $e$  incident with it otherwise. Notice that since we cannot apply the first case, this contraction does not decrease the number of cycles in  $G$ . Also, we can assume without loss of generality that  $v$  (respectively  $e$ ) is not part of a minimum  $x$ -cover of cycles in  $G$ , as any vertex adjacent to  $v$  (respectively edge incident with  $e$ ) covers all the cycles covered by  $v$  (respectively  $e$ ). Therefore the obtained graph  $G'$  satisfies  $x\text{-pack}_{\mathcal{H}}(G') = x\text{-pack}_{\mathcal{H}}(G)$  and  $x\text{-cover}_{\mathcal{H}}(G) = x\text{-cover}_{\mathcal{H}}(G')$ . It is not hard to see that  $A_x(G') < A_x(G)$ . Therefore we can apply the induction hypothesis on  $G'$  and obtain the desired result on  $G'$ , that immediately translates to  $G$  by the above remarks.  $\square$

By setting  $q = 3k$  and observing that every graph containing  $K_{3k}$  as a minor also contains  $k$  vertex-disjoint cycles (hence also edge-disjoint), [Lemma 5.18](#) yields the vertex and edge versions of the classic Erdős–Pósa theorem as a corollary.

The technique presented in this section has been used to show the following result.

**Theorem 5.19** ([\[FJW13a\]](#)). *For every forest  $H$ ,  $\mathcal{M}(H)$  has the  $v$ -Erdős–Pósa property with gap  $O_H(k)$ .*

Other applications will be presented in [Section 5.4](#). To extend the idea of [Lemma 5.18](#) in order to prove that some graph class  $\mathcal{H}$  has the  $x$ -Erdős–Pósa property with gap  $f: \mathbb{N} \rightarrow \mathbb{N}$ , one should show that for every positive integer  $k$  and every graph  $G$  with  $x\text{-pack}_{\mathcal{H}}(G) \leq k$ ,

- either there is a graph  $G'$  with  $x\text{-pack}_{\mathcal{H}}(G) = x\text{-pack}_{\mathcal{H}}(G')$  and  $x\text{-cover}_{\mathcal{H}}(G) = x\text{-cover}_{\mathcal{H}}(G')$  and such that  $|G'| + \|G'\| < |G| + \|G\|$  (reduction case);
- or  $G$  has a subgraph isomorphic to a member of  $\mathcal{H}$  on at most  $f(k)/k$  vertices/edges (progress case).

## 5.4 Applications to selected classes of graphs

In this section, we show classes of graphs where the techniques presented in [Section 5.3](#) can be applied to yield Erdős–Pósa type results. The classes we will consider are the following:

- majors of wheels, yurts, and graphs of pathwidth at most two in [Subsection 5.4.1](#);
- majors of  $\theta_r$  in [Subsection 5.4.2](#);
- immersion models of connected planar subcubic graphs in [Subsection 5.4.4](#);

### 5.4.1 Wheels, yurts, and graphs of pathwidth at most two

The gap provided by [Corollary 5.9](#) relies on general exclusion theorems for majors of planar graphs. In this section we provide an application of the results of [Section 5.3.1](#) to decrease the contribution of  $H$ , for specific planar patterns where a better exclusion theorem is known. The classes we will consider are the majors of the following graphs: wheels, double wheels, graphs of pathwidth at most 2, and yurt graphs; and we will use the exclusion theorems introduced in [Section 4.3](#).

**Theorem 5.20.** *The class  $\mathcal{M}(H)$  has the  $\nu$ -Erdős–Pósa-property with gap:*

- (i)  $k \mapsto kh^2 \text{polylog}(kh)$  if  $H$  is a wheel of order  $h$ ;
- (ii)  $k \mapsto kh^4 \text{polylog}(kh)$  if  $H$  is a graph of pathwidth at most two with  $|V(H)| + |E(H)| = h$  or a double wheel of order  $h$ ;
- (iii)  $k \mapsto kh^8 \text{polylog}(kh)$  if  $H$  is a yurt graph of order  $h$ .

*Proof.* Let us prove (i). We denote by  $W_h$  the wheel of order  $h$ , for every positive integer  $h$ . According to [Theorem 4.27](#), for every positive integer  $h$ , every graph of treewidth at least  $36h - 2$  contains a major of  $W_h$ . [Lemma 5.6](#) then implies that there is a ceiling  $f_h$  for  $(\text{tw}, \mathcal{M}(W_h), \nu)$ , with  $f_h(k) = k(36h - 2)^2 \text{polylog}(kh)$ . Thanks to [Lemma 5.7](#), this ceiling yields a gap  $g_h(k) = 5k(36h - 2)^2 \text{polylog}(kh) \cdot \log(k + 1)$  and we are done. The proofs of (ii) and (iii) follow the very same path, using exclusion theorems [Theorem 4.28](#), [Theorem 4.29](#), and [Theorem 4.31](#).  $\square$

### 5.4.2 Pumpkins

A way to extend the classic Erdős–Pósa Theorem is to consider generalizations of the class of cycles. A class that attracted some attention in this direction is  $\mathcal{M}(\theta_r)$  for  $r \in \mathbb{N}$ , which is the class of cycles when  $r = 2$ . For instance, Fomin et al. gave in [\[FST11\]](#) the following extension.

**Theorem 5.21** ([\[FST11\]](#)). *There is a function  $f_r(k) = O(k^2 r^2)$  such that for every  $r \in \mathbb{N}$ ,  $\mathcal{M}(\theta_r)$  has the  $\nu$ -Erdős–Pósa property with gap  $f_r$ .*

In an unpublished manuscript [\[FJS13\]](#), Fiorini et al. proved that the classic Erdős–Pósa Theorem can be extended to  $\mathcal{M}(\theta_r)$  (instead of cycles) without increasing the order of magnitude of the gap,  $O(k \log k)$ .

**Theorem 5.22** ([\[FJS13\]](#), see also [\[CRST15b\]](#)). *There is a function  $f_r(k) = O_r(k \log k)$  such that for every  $r \in \mathbb{N}$ ,  $\mathcal{M}(\theta_r)$  has the  $\nu$ -Erdős–Pósa property with gap  $f_r$ .*

We present in this section the following result.

**Theorem 5.23.** *There is a function  $f_r$  with  $f_r(k) = O(k^2 r^2 \text{polylog } kr)$  and  $f_r(k) = O(k^4 r^2 \text{polylog } kr)$ , such that for every  $r \in \mathbb{N}$ ,  $\mathcal{M}(\theta_r)$  has the e-Erdős–Pósa property with gap  $f_r$ .*

**Theorem 5.23** is an edge-analogue of **Theorem 5.22**. The bound on the gap is worse in terms of  $k$  but it indicated the contribution of  $r$ , which is polynomial.

We will in **Chapter 6** prove the following result, that completes the symmetry between the vertex and edge settings.

**Theorem 5.24.** *There is a function  $f_r(k) = O_r(k \log k)$  such that for every  $x \in \{v, e\}$  and every  $r \in \mathbb{N}$ ,  $\mathcal{M}(\theta_r)$  has the  $x$ -Erdős–Pósa property with gap  $f_r$ .*

Let us here prove **Theorem 5.23**. Recall that we proved in **Section 4.2** that every 2-connected graph that has a vertex of degree at least  $2kr$  has a subgraph that is an edge-disjoint union of  $k$   $\mathcal{M}(\theta_r)$ -subgraphs. Let us show the next result.

**Lemma 5.25.** *Let  $r \in \mathbb{N}$ . If  $\mathcal{M}(\theta_r)$  has the  $v$ -Erdős–Pósa property with a gap  $f$  that is superadditive, then it has the e-Erdős–Pósa property with gap  $k \mapsto 2kr \cdot f(k)$ .*

We first need some intermediate lemmas in order to be able to use **Theorem 5.3**.

**Lemma 5.26.** *Let  $d \in \mathbb{N}$ . The class  $\mathcal{M}(\theta_r)$  has the e-Erdős–Pósa property for the class of graphs of maximum degree at most  $d$  with gap  $k \mapsto d \cdot f(k)$ , where  $f$  is the (vertex-) gap provided by **Theorem 5.22**.*

*Proof.* Let  $G$  be a graph satisfying  $\Delta(G) \leq d$  and let  $k$  be a positive integer. If  $G$  contains  $k$  vertex-disjoint  $\mathcal{M}(\theta_r)$ -subgraphs, these subgraphs are in particular edge-disjoint hence  $\text{e-pack}_{\mathcal{M}(\theta_r)}(G) \geq k$ . On the other hand, if  $\text{v-pack}_{\mathcal{M}(\theta_r)}(G) < k$ , according to **Theorem 5.22** there is a subset  $X \subseteq V(G)$  such that  $|X| \leq f(k)$  and  $G \setminus X$  does not contain any  $\mathcal{M}(\theta_r)$ -subgraph. Let  $Y$  be the set of edges incident to the vertices in  $X$ . As  $\Delta(G) \leq d$  we have  $|Y| \leq d \cdot f(k)$ . Notice that any  $\mathcal{M}(\theta_r)$ -subgraph of  $G \setminus Y$  does not contain a vertex from  $X$ , hence the existence of such a subgraph would contradict the definition of  $X$ . This proves that  $\text{e-cover}_{\mathcal{M}(\theta_r)} \leq d \cdot f(k)$  and we are done.  $\square$

**Lemma 5.27.** *If a class of 2-connected graphs  $\mathcal{H}$  has the e-Erdős–Pósa property for 2-connected graphs with a superadditive gap, then it has the e-Erdős–Pósa property (for all graphs) with the same gap.*

*Proof.* Let  $f$  be the gap mentioned in the statement of the lemma. Let  $G$  be a graph and let  $G_1, \dots, G_t$  be its 2-connected components, for some positive integer  $t$ . Let  $k$  be an integer and let  $p_i = \text{e-pack}_{\mathcal{H}}(G_i)$ , for every  $i \in \llbracket 1, t \rrbracket$ . If  $p_1 + \dots + p_t \geq k$ , then  $\text{e-pack}_{\mathcal{H}}(G) \geq k$ , as two subgraphs in two distinct 2-connected components of  $G$  are always edge-disjoint. On the other hand, let us assume that  $p_1 + \dots + p_t < k$ . We have  $\text{e-cover}_{\mathcal{H}}(G_i) \leq f(p_i)$  for every  $i \in \llbracket 1, t \rrbracket$ . Since no  $\mathcal{H}$ -subgraph of  $G$  contains edges

from two or more distinct 2-connected components, we can cover  $\mathcal{H}$ -subgraphs in each 2-connected component in order to cover them in the graph:

$$\begin{aligned}
\text{e-cover}_{\mathcal{H}}(G) &\leq \sum_{i=1}^t \text{e-cover}_{\mathcal{H}}(G_i) \\
&\leq \sum_{i=1}^t f(p_i) \\
&\leq f\left(\sum_{i=1}^t p_i\right) && (f \text{ is superadditive}) \\
&\leq f(k-1).
\end{aligned}$$

□

We can now prove [Lemma 5.25](#).

*Proof of [Lemma 5.25](#).* By applying [Theorem 5.3](#) to [Lemma 4.16](#) and [Lemma 5.26](#), we obtain the result for 2-connected graphs. [Lemma 5.27](#) then allows us to extend it to graphs that are not 2-connected. □

We also need the following result before we prove [Theorem 5.23](#).

**Theorem 5.28** ([\[BvLTT97b, Theorem 14\]](#)). *For every  $r \in \mathbb{N}$ , every graph of treewidth at least  $2r - 1$  contains a  $K_{2,r}$  major.*

Let us now prove [Theorem 5.23](#). We first restate it.

**Theorem 5.23.** *There is a function  $f_r$  with  $f_r(k) = O(k^2 r^2 \text{polylog } kr)$  and  $f_r(k) = O(k^4 r^2 \text{polylog } kr)$ , such that for every  $r \in \mathbb{N}$ ,  $\mathcal{M}(\theta_r)$  has the e-Erdős–Pósa property with gap  $f_r$ .*

*Proof.* [Lemma 5.6](#) applied to  $\mathcal{M}(\theta_r)$  using [Theorem 5.28](#) yields a gap  $f_r(k) = O(kr^2 \text{polylog } kr)$  for the vertex-Erdős–Pósa property of  $\mathcal{M}(\theta_r)$ . Then, [Lemma 5.25](#) yields a gap  $2kr \cdot f_r(k) = O(k^2 r^3 \text{polylog } kr)$  for the edge version, as required. A gap of order of magnitude  $O(k^4 r^2 \text{polylog } kr)$  can be obtained the same way using [Theorem 5.5](#) instead of [Lemma 5.6](#). □

### 5.4.3 Double pumpkins

For every  $r, r' \in \mathbb{N}$ , we denote by  $\theta_{r,r'}$  the graph obtained by identifying one vertex of  $\theta_r$  with one vertex of  $\theta_{r'}$ . In this section, we use the tools provided by [Section 5.3](#) to prove that  $\mathcal{M}(\theta_{r,r'})$  have the edge-Erdős–Pósa-property for simple graphs.

**Theorem 5.29.** *For every  $r, r' \in \mathbb{N}$ ,  $\mathcal{M}(\theta_{r,r'})$  has the edge-Erdős–Pósa property for simple graphs.*

We must note that at the time of writing, the graphs  $\{\theta_{r,r'}\}_{r,r' \in \mathbb{N}}$  are the only graphs on at least three vertices, the majors of which are known to have the edge-Erdős–Pósa property. This fact must be compared to the results on the vertex variant where the graphs  $H$  for which  $\mathcal{M}(H)$  has the vertex-Erdős–Pósa property have been completely characterized: they are exactly the planar graphs [RS86, Theorem 8.2] (see Section 5.6). This raises the following question.

*Question 5.30.* What are the graphs, the majors of which have the edge-Erdős–Pósa property?

**Lemma 5.47** provides a partial answer to **Question 5.30**: all these graphs are planar. Prior to the proof of **Theorem 5.29**, we need to introduce a result of Ding et al. [DO96], the statement of which requires additional definitions.

**Walls, fans, paths, and stars.** Wall have been defined in **Subsection 2.3.2**. As a reminder, the 7-wall is depicted in **Figure 5.1**. The  $n$ -fan is the graph obtained by adding a dominating vertex to a path on  $n$  vertices. A collection of paths is said to be *independent* if two paths of the collection never share interior vertices. The  $n$ -star is the graph obtained by replacing every edge of  $K_{1,n}$  with  $n$  independent paths of two edges. The  $n$ -path is the graph obtained by replacing every edge of an  $n$ -edge path with  $n$  independent paths of two edges. Examples of these graphs are depicted in **Figure 5.1**. The *wall number* (respectively *fan number*, *star number*, and *path number*) of a graph  $G$  is defined as the largest integer  $k$  such that  $G$  contains a subdivision of a  $k$ -wall (respectively of a  $k$ -fan, of a  $k$ -star, of a  $k$ -path), or infinity is no such integer exists. Let  $\gamma(G)$  denote the maximum of the wall number, fan number, star number, and path number of a graph  $G$ .

Ding et al. gave in [DO96] the following characterization of classes of graphs of bounded tree-partition width in terms of excluded topological minors.

**Theorem 5.31** ([DO96]). *There is a function  $f_5: \mathbb{N} \rightarrow \mathbb{N}$  such that every simple graph  $G$  satisfies  $\text{tpw}(G) \leq f_5(\gamma(G))$ .*

In other words, for every integer  $k$ , every simple graph of large enough tree-partition width contains a subdivision of one of the following graphs: the  $k$ -wall, the  $k$ -fan, the  $k$ -path, or the  $k$ -star.

Notice that for every  $r, r' \in \mathbb{N}$ ,  $r' \leq r$ , the graph  $\theta_{r,r'}$  is a minor of the following graphs: the  $r$ -path, the  $r$ -star, the  $(r + r' + 1)$ -fan, and the  $r$ -wall (for  $r \geq 6$ ). Hence, every simple graph of large enough tree-partition width contains a  $\theta_{r,r'}$ -major. This can easily be generalized to edge-disjoint packings, as follows.

**Lemma 5.32.** *For every  $r, r', k \in \mathbb{N}$ , every graph  $G$  satisfying  $\gamma(G) \geq k(r + r' + 7)$  contains an  $e\text{-}\mathcal{M}(\theta_{r,r'})$ -packing of size  $k$ .*

Using **Theorem 5.31**, we get the following corollary.

**Corollary 5.33.** *For every  $r, r', k \in \mathbb{N}$ , every simple graph  $G$  satisfying  $\text{tpw}(G) \geq f_5(k(r + r' + 7))$  contains an  $e\text{-}\mathcal{M}(\theta_{r,r'})$ -packing of size  $k$ .*




Figure 5.1: Unavoidable patterns of graphs of large tree-partition width.

In other words, [Corollary 5.33](#) provides a ceiling for the triple  $(\text{tpw}, \mathcal{M}(\theta_{r,r'}), e)$  in simple graphs. The proof of [Theorem 5.29](#) now follows by a straightforward application of [Theorem 5.3](#) to [Corollary 5.33](#) and [Lemma 5.15](#).

#### 5.4.4 Planar subcubic graphs

In this section, we apply the tools of [Section 5.3](#) to immersion expansions of planar subcubic graphs. The result we prove is the following.

**Theorem 5.34.** *For every connected planar subcubic graph  $H$  on  $h > 0$  edges and every  $x \in \{v, e\}$ , the class  $\mathcal{I}(H)$  has the  $x$ -Erdős–Pósa property with a gap that is polynomial in both  $h$  and the packing number.*

We will show in [Section 5.5](#) that neither the planarity requirement, nor the subcubicity can be dropped.

The main tools of our proof are the graph invariants of tree-cut width and tree-partition width. Our proof uses the exclusion result [Theorem 4.66](#) which implies that (as the disjoint union of planar subcubic graphs is planar subcubic), for every fixed planar subcubic graph  $H$  and every positive integer  $k$ , every graph of large enough tree-cut width contains  $k$  vertex-disjoint immersion expansions of  $H$ . This allows us to focus on graphs of bounded tree-cut width. By applying suitable reductions, we finally reduce the problem to graphs of bounded tree partition width ([Lemma 5.35](#)). The result then follows by the application of [Theorem 5.3](#).

## From tree-cut decompositions to tree-partitions

The purpose of this section is to reduce the proof of [Theorem 5.34](#) (when  $x = e$ ) for host graphs with bounded tree-cut width to the case where host graphs have bounded tree-partition width. In particular, we prove the following lemma.

**Lemma 5.35.** *For every connected graph  $G$ , and every connected graph  $H$  with at least one edge, there is a graph  $G'$  and a graph  $H'$  such that*

- $\text{tpw}(G') \leq (\text{tcw}(G) + 1)^2/2$ ,
- $\text{e-pack}_{\mathcal{I}(H')}(G') \leq \text{e-pack}_{\mathcal{I}(H)}(G)$ , and
- $\text{e-cover}_{\mathcal{I}(H)}(G) \leq \text{e-cover}_{\mathcal{I}(H')}(G')$ .

Observe that, with the notation of [Lemma 5.35](#), if we prove that

$$\text{e-cover}_{\mathcal{I}(H')}(G') \leq f(\text{e-pack}_{\mathcal{I}(H')}(G'))$$

for some non-decreasing function  $f: \mathbb{N} \rightarrow \mathbb{N}$ , then it immediately implies

$$\text{e-cover}_{\mathcal{I}(H)}(G) \leq f(\text{e-pack}_{\mathcal{I}(H)}(G)).$$

For every vertex  $v$  of a graph  $G$ , we denote by  $\text{mdeg}_G(v)$  (or  $\text{mdeg}(v)$  when  $G$  is given by the context) the number of edges (counting multiplicities) incident with  $v$ . For every graph  $G$ , we define  $G^+$  as the graph obtained if, for every vertex  $v$ , we add two new vertices  $v'$  and  $v''$  and the edges  $\{v', v''\}$  (of multiplicity 2),  $\{v, v'\}$  and  $\{v, v''\}$  (both of multiplicity 1). Observe that for every  $G$ , every vertex of  $G^+$  is incident to at least 3 edges (counting multiplicities). We also define  $G^*$  as the graph obtained by adding, for every vertex  $v$ , the new vertices  $v'_1, \dots, v'_{\text{mdeg}(v)}$  and  $v''_1, \dots, v''_{\text{mdeg}(v)}$  and the edges  $\{v'_i, v''_i\}$  (of multiplicity 2),  $\{v, v'_i\}$ , and  $\{v, v''_i\}$  (both of multiplicity 1), for every  $i \in \llbracket 1, \text{deg}(v) \rrbracket$ . If  $v$  is a vertex of  $G$ , then we denote by  $Z_{v,i}$  the subgraph  $G^*[\{v, v'_i, v''_i\}]$  for every  $i \in \llbracket 1, \text{mdeg}_G(v) \rrbracket$ .

*Observation 5.36.* Let  $H$  and  $G$  be two graphs, and let  $(\phi, \psi)$  be an  $H$ -immersion model in  $G$ . Then for every vertex  $x$  of  $G$ , we have  $\text{mdeg}_H(x) \leq \text{mdeg}_G(\phi(x))$ .

Our first aim is to prove the following three lemmata.

**Lemma 5.37.** *Let  $G$  be a graph, let  $H$  be a connected graph with at least one edge and let  $G'$  be a subdivision of  $G^*$ . Then we have*

- $\text{e-pack}_{\mathcal{I}(H^+)}(G^*) = \text{e-pack}_{\mathcal{I}(H^+)}(G')$  and
- $\text{e-cover}_{\mathcal{I}(H^+)}(G^*) = \text{e-cover}_{\mathcal{I}(H^+)}(G')$ .

*Proof.* We denote by  $S$  the set of subdivision vertices added during the construction of  $G'$  from  $G^+$ . As  $G'$  is a subdivision of  $G^*$ , we have  $\text{e-pack}_{\mathcal{I}(H^+)}(G') \geq \text{e-pack}_{\mathcal{I}(H^+)}(G^*)$  and  $\text{e-cover}_{\mathcal{I}(H^+)}(G') \geq \text{e-cover}_{\mathcal{I}(H^+)}(G^*)$ .

As a consequence of [Observation 5.36](#) and the fact that every vertex of  $H^+$  is incident with at least 3 edges (counting multiplicities), if  $M$  is an  $H^+$ -immersion expansion in  $G'$  then no branch vertex of  $M$  belongs to  $S$ . Indeed, every vertex of  $S$  has multidegree 2 in  $G'$ . Therefore, by dissolving in  $M$  the vertices of  $S$  that belong to  $V(M)$ , we obtain an  $H^+$ -immersion expansion in  $G^*$ . It follows that  $\text{e-pack}_{\mathcal{I}(H^+)}(G^*) \geq \text{e-pack}_{\mathcal{I}(H^+)}(G')$ , hence  $\text{e-pack}_{\mathcal{I}(H^+)}(G^*) = \text{e-pack}_{\mathcal{I}(H^+)}(G')$ .

On the other hand, let  $X$  be an  $H^+$ -cover of  $G^*$  and let  $X'$  be a set of edges constructed by taking, for every  $e \in X$ , an edge of the path of  $G'$  connecting the endpoints of  $e$  that has been created by subdividing  $e$ . Assume that  $X'$  is not an  $H^+$ -cover of  $G'$ . According to the remark above, this implies that  $X$  is not an  $H^+$ -cover of  $G^*$ , a contradiction. Hence  $X'$  is an  $H^+$ -cover of  $G'$  and thus  $\text{e-cover}_{\mathcal{I}(H^+)}(G^*) = \text{e-cover}_{\mathcal{I}(H^+)}(G')$ .  $\square$

**Lemma 5.38.** *For every two graphs  $H$  and  $G$  such that  $H$  is connected and has at least one edge, we have  $\text{e-pack}_{\mathcal{I}(H^+)}(G^*) \leq \text{e-pack}_{\mathcal{I}(H)}(G)$ .*

*Proof.* In  $G^*$  (respectively  $H^+$ ), we say that a vertex is *original* if it belongs to  $V(G)$  (respectively  $V(H)$ ). Let  $(\phi, \psi)$  be an  $H^+$ -immersion model in  $G^*$ .

We first show that if  $u$  is an original vertex of  $H^+$ , then  $\phi(u)$  is an original vertex of  $G^*$ . By contradiction, let us assume that  $\phi(u)$  is not original, for some original vertex  $u$  of  $H^+$ . Then  $\phi(u) = v'_i$  or  $\phi(u) = v''_i$ , for some  $v \in V(G)$  and  $i \in \llbracket 1, \text{mdeg}_G(v) \rrbracket$ .

Observe that since  $H$  is connected and has at least one edge, every vertex of  $H^+$  has degree at least three: let  $x, y$ , and  $z$  be the endpoints of three multiedges incident with  $u$ . Then  $\psi(\{u, x\})$ ,  $\psi(\{u, y\})$ , and  $\psi(\{u, z\})$  are edge-disjoint paths connecting  $\phi(u)$  to three distinct vertices. This is not possible because there is an edge cut of size two,  $\{\{v, v'_i\}, \{v, v''_i\}\}$ , separating the two vertices  $v'_i$  and  $v''_i$  (among which is  $\phi(u)$ ) from the rest of the graph. Consequently, if  $u \in V(H^+)$  is original, then  $\phi(u)$  is original.

Let us now consider an edge  $\{u, v\} \in E(H)$ . By the above remark,  $\phi(u)$  and  $\phi(v)$  are original vertices of  $G^*$ . It is easy to see that  $\psi(\{u, v\})$  contains only original vertices of  $G^*$ . Indeed, if this path contained a non-original vertex  $w'$  or  $w''$  for some original vertex  $w$  of  $V(G^*)$ , it would use  $w$  twice in order to reach  $u$  and  $v$ , what is not allowed. Therefore, from the definition of  $H^+$ , the pair  $(\phi|_{V(H)}, \psi|_{E(H)})$  is an  $H$ -immersion model of  $G$ .

We proved that every  $H^+$ -immersion-expansion of  $G^*$  contains an  $H$ -immersion-expansion that belongs to the subgraph  $G$  of  $G^*$ . Consequently every  $H^+$ -packing of  $G^*$  contains an  $H$ -packing of the same size that belongs to  $G$ , and the desired inequality follows.  $\square$

**Lemma 5.39.** *For every two graphs  $H$  and  $G$  such that  $H$  is connected and has at least one edge, we have  $\text{e-cover}_{\mathcal{I}(H)}(G) \leq \text{e-cover}_{\mathcal{I}(H^+)}(G^*)$ .*

*Proof.* Similarly to the proof of [Lemma 5.38](#), we say that an edge of  $G^*$  is *original* if it belongs to  $E(G)$ . Let  $X \subseteq E(G^*)$  be a minimum cover of  $H^+$ -immersion expansions in  $G^*$ .

*First case:* all the edges in  $X$  are original. In this case,  $X$  is an  $H$ -cover of  $G$  as well. Indeed, if  $G \setminus X$  contains an  $H$ -immersion expansion  $M$ , then  $G^* \setminus X$  contains  $M^*$  that, in turn, contains  $H^+$ . Hence in this case,  $\text{e-cover}_{\mathcal{I}(H)}(G) \leq \text{e-cover}_{\mathcal{I}(H^+)}(G^*)$ .

*Second case:* there is an edge  $e \in X$  that is not original. Let  $v$  be the original vertex of  $G^*$  such that either  $e \in Z_{v,l}$  for some  $l \in \llbracket 1, \text{mdeg}_G(v) \rrbracket$ . Let us first show the following claim.

*Claim:* For every  $i \in \llbracket 1, \text{mdeg}_G(v) \rrbracket$ , there is an edge of  $Z_{v,i}$  that belongs to  $X$ .

*Proof of claim:* Looking for a contradiction, let us assume that we have  $E(Z_{v,i}) \cap X = \emptyset$ , for some  $i \in \llbracket 1, \text{mdeg}_G(v) \rrbracket$ . Clearly  $i \neq l$ . By minimality of  $X$ , the graph  $G \setminus (X \setminus \{e\})$  contains an  $H^+$ -immersion expansion  $M$  that uses  $e$ . Observe that  $M' = M \setminus E(Z_{v,l}) \cup E(Z_{v,i})$  contains an  $H^+$ -immersion expansion (since  $Z_{v,l}$  and  $Z_{v,i}$  are isomorphic). Hence,  $M'$  is a subgraph of  $G \setminus (X \setminus \{e\})$  that contains an  $H^+$ -immersion expansion. This is not possible as  $X$  is a cover, so we reach the contradiction we were looking for and the claim holds.  $\diamond$

We build a set  $Y$  as follows. For every edge  $f \in X$ , if  $f$  is original then we add to  $Y$ . Otherwise, if  $v_f$  is the (original) vertex of  $G^*$  such that  $e \in E(Z_{v_f,i})$  for some  $i \in \llbracket 1, \text{mdeg}_G(v_f) \rrbracket$ , then we add to  $Y$  all edges that are incident to  $v_f$  in  $G$ .

The above claim ensures that when a non-original edge  $f$  of  $X$  is encountered, then  $X$  contains an edge in each of  $Z_{v_f,1}, \dots, Z_{v_f, \text{mdeg}_G(v_f)}$ . Therefore, the same set of edges, of size  $\text{mdeg}_G(v_f)$ , will be added to  $Y$  when encountering an other edge from  $Z_{v_f,1}, \dots, Z_{v_f, \text{mdeg}_G(v_f)}$ . Consequently,  $|X| = |Y|$ .

Let us now show that  $Y$  is an  $H^+$ -cover of  $G^*$ . Suppose that there exists an  $H^+$ -immersion expansion  $M$  in  $G^* \setminus Y$ . Observe that since  $H$  is connected and has at least one edge,  $M$  does not belong to  $\bigcup_{i \in \{1, \dots, \text{mdeg}_G(u)\}} Z_{u,i}$  for every original vertex  $u$  of  $G^*$ . Let

$$Z = \bigcup_{u \in V(G)} \bigcup_{i \in \{1, \dots, \text{mdeg}_G(u)\}} E(Z_{u,i})$$

Then  $M$  is a subgraph of  $G \setminus (Y \cup Z)$ . As  $X \subseteq Y \cup Z$ , this contradicts the fact that  $X$  is a cover. Therefore,  $Y$  is an  $H^+$ -cover. Moreover all the edges in  $Y$  are original. As this situation is treated by the first case above, we are done.  $\square$

If  $(T, \{X_t\}_{t \in V(T)})$  is a tree-cut decomposition of a graph  $G$  and  $t \in V(T)$ , we say than an edge of  $G$  crosses the bag  $X_t$  if its endpoints belongs to bags  $X_{t_1}$  and  $X_{t_2}$ , for some  $t_1, t_2 \in V(T)$  such that  $t$  belongs to the interior of the (unique) path of  $T$  connecting  $t_1$  to  $t_2$ . We are now ready to prove [Lemma 5.35](#), that we recall hereafter.

**Lemma 5.35.** *For every connected graph  $G$ , and every connected graph  $H$  with at least one edge, there is a graph  $G'$  and a graph  $H'$  such that*

- $\text{tpw}(G') \leq (\text{tcw}(G) + 1)^2/2$ ,
- $\text{e-pack}_{\mathcal{I}(H')}(G') \leq \text{e-pack}_{\mathcal{I}(H)}(G)$ , and
- $\text{e-cover}_{\mathcal{I}(H)}(G) \leq \text{e-cover}_{\mathcal{I}(H')}(G')$ .

*Proof.* Let  $k = \text{tcw}(G)$ . We examine the nontrivial case where  $G$  is not a tree, i.e.,  $\text{tcw}(G) \geq 2$ . Let us consider the graph  $G^*$ . We claim that  $\text{tcw}(G^*) = \text{tcw}(G)$ . Indeed, starting from an optimal tree-cut decomposition of  $G$ , we can, for every vertex  $v$  of  $G$  and for every  $i \in \llbracket 1, \text{mdeg}_G(v) \rrbracket$ , create a bag that is a children of the one of  $v$  and contains  $\{v'_i, v''_i\}$ . According to the definition of  $G^*$ , this creates a tree-cut decomposition  $\mathcal{D} = ((T, s), \{X_t\}_{t \in V(T)})$  of  $G^*$ . Observe that for every vertex  $x$  that we introduced to the tree of the decomposition during this process,  $\text{adh}_{\mathcal{D}}(x) = 2$  and the corresponding bag has size two. This proves that  $\text{tcw}(G^*) \leq \max(\text{tcw}(G), 2) = \text{tcw}(G)$ . As  $G$  is a subgraph of  $G^*$ , we obtain  $\text{tcw}(G) \leq \text{tcw}(G^*)$  and the proof of the claim is complete.

According to [Proposition 2.8](#), we can assume that  $G^*$  has a nice rooted tree-cut decomposition of width  $\leq k$ . For notational simplicity we again denote it by  $\mathcal{D} = ((T, s), \{X_t\}_{t \in V(T)})$  and, obviously, we can also assume that all leaves of  $T$  correspond to non-empty bags.

Our next step is to transform the rooted tree-cut decomposition  $\mathcal{D}$  into a rooted tree-partition  $\mathcal{D}' = ((T, s), \{X'_t\}_{t \in V(T)})$  of a subdivision  $G'$  of  $G^*$ . Notice that the only differences between two decompositions are that, in a tree-cut decomposition, empty bags are allowed as well as edges connecting vertices of bags corresponding to non-adjacent vertices of  $T$ .

We proceed as follows: if  $X$  is a bag crossed by edges, we subdivide every edge crossing  $X$  and add the obtained subdivision vertex to  $X$ . By repeating this process we decrease at each step the number of bags crossed by edges, that eventually reaches zero. Let  $G'$  be the obtained graph and observe that  $G'$  is a subdivision of  $G$ . As  $G$  is connected, the obtained rooted tree-cut decomposition  $\mathcal{D}' = ((T, s), \{X'_t\}_{t \in V(T)})$  is a rooted tree partition of  $G'$ .

Recall that we say that a vertex  $t \in V(T)$  is thin if  $\text{adh}(t) \leq 2$ , bold otherwise. Notice that the adhesion of any bag of  $T$  in  $\mathcal{D}$  is the same as in  $\mathcal{D}'$ . However, the bags of  $\mathcal{D}'$  may grow during the construction of  $G'$ . Let  $t$  be a vertex of  $T$  and let  $\{t_1, \dots, t_m\}$  be the set of children of  $t$ . We claim that  $|X'_t| \leq (k+1)^2/2$ .

Let  $E_t$  be the set of edges crossing  $X_t$  in  $G$ . Let  $H_t$  be the torso of  $\mathcal{D}$  at  $t$ , and let  $H'_t = H_t \setminus X_t$ . Observe that  $|E_t|$  is the same as the number of edges in  $H'_t$ . Let  $z_p$  be the vertex of  $H'_t$  corresponding to the parent of  $t$ , and similarly for each  $i \in \{1, \dots, m\}$  let  $z_i$  be the vertex of  $H'_t$  corresponding to the child  $t_i$  of  $t$ . Notice that if  $t_i$  is a thin child of  $t$ , then  $z_i$  can be adjacent to only  $z_p$  as  $\mathcal{D}$  is a nice rooted tree-cut decomposition. Thus the sum of the number of incident edges with  $z_i$  in  $H'_t$  for all thin children  $t_i$  of  $t$  is at most  $\text{adh}_{\mathcal{D}}(t) \leq k$ . On the other hand, if  $t_i$  is a bold child of  $t$ , then  $z_i$  has at least 3 neighbors in  $H_t$ , and thus it is contained in the 3-center of  $(H_t, X_t)$ . Thus, the number of all bold children of  $t$  is bounded by  $k - |X_t|$ . Since each vertex in  $H'_t$  is incident with at most  $k$  edges, the total number of edges in  $H'_t$  is at most  $(k - |X_t| + 1)k/2 + k$ . As  $|E(H'_t)| = |E_t| = |X'_t \setminus X_t|$ , it implies that  $|X'_t| \leq |X_t| + k \cdot (k - |X_t| + 2)/2 \leq \max\{2k, k(k+2)/2\} \leq (k+1)^2/2$ . We conclude that  $G'$  has a rooted tree-partition of width at most  $(\text{tcw}(G) + 1)^2/2$ .

Recall that  $G'$  is a subdivision of  $G^*$ . By the virtue of [Lemma 5.39](#), [Lemma 5.38](#), and [Lemma 5.37](#) (relating the packing and covering numbers of  $\mathcal{I}(H)$  and  $\mathcal{I}(H^+)$  in  $G$  and  $G'$ ) we obtain that  $\text{e-pack}_{\mathcal{I}(H^+)}(G') \leq \text{e-pack}_{\mathcal{I}(H)}(G)$  and  $\text{e-cover}_{\mathcal{I}(H)}(G) \leq$

$\text{e-cover}_{\mathcal{I}(H^+)}(G')$ . Hence  $G'$  satisfies the desired properties.  $\square$

Therefore we get the following lemma.

**Lemma 5.40.** *Let  $H$  be a graph on  $h$  edges, let  $r$  be an integer and let  $G$  be a graph such that  $\text{tcw}(G) \leq r$ . Then  $\text{e-cover}_{\mathcal{I}(H)}(G) \leq \text{e-pack}_{\mathcal{I}(H)}(G) \cdot h(r+1)^4$ .*

*Proof.* Let  $G'$  and  $H'$  be the graphs given by [Lemma 5.35](#). As  $\text{tcw}(G) \leq r$  we have  $\text{tpw}(G') \leq (r+1)^2/2$ . Applying [Lemma 5.15](#), we get:

$$\text{e-cover}_{\mathcal{I}(H')}(G') \leq \text{e-pack}_{\mathcal{I}(H')}(G') \cdot h(r+1)^4.$$

Then [Lemma 5.35](#) provides the desired inequality (cf. the remark following it).  $\square$

Recall that our current goal is to prove [Theorem 5.34](#), that we restate for convenience.

**Theorem 5.34.** *For every connected planar subcubic graph  $H$  on  $h > 0$  edges and every  $x \in \{v, e\}$ , the class  $\mathcal{I}(H)$  has the  $x$ -Erdős–Pósa property with a gap that is polynomial in both  $h$  and the packing number.*

The edge version of this result follows from the application of the Master Theorem [Theorem 5.3](#), using the ceiling provided by [Theorem 4.66](#) together with [Lemma 5.40](#).

### The vertex case

To prove the vertex version of [Theorem 5.34](#) is a much easier task. For this, we follow the same methodology by using the graph parameter of treewidth instead of tree-cut width, and topological minors instead of immersions. We use the following vertex-counterpart of [Theorem 4.66](#).

**Lemma 5.41.** *For every  $h \in \mathbb{N}$  there is a function  $f: \mathbb{N} \rightarrow \mathbb{N}$  with  $f(k) = (h \cdot k)^{O(1)}$  such that, for every planar subcubic graph  $H$  with  $|V(H)| + |E(H)| = h$ ,  $f$  is a ceiling for  $(\text{tw}, \mathcal{I}(H), v)$ .*

*Proof.* A equivalent statement of [Theorem 4.26](#) is that every graph of treewidth at least  $f_1(k)$  contains a subdivision of a wall of height and width  $\Omega(k)$  as a subgraph, for every  $k \in \mathbb{N}$ . According to [Lemma 4.70](#), every planar subcubic graph  $H$  is an immersion of the wall of width and height  $h$ , where  $h = |V(H)| + |E(H)|$ . Therefore the wall of width and height  $kh$  contains  $k$ -edge-disjoint  $\mathcal{I}(H)$ -subgraphs. We deduce that every graph of treewidth at least  $f_1(c \cdot hk)$  contains  $k$  vertex-disjoint  $\mathcal{I}(H)$ -subgraphs, for some constant  $c$  not depending on  $H$ . We note that  $f_1$  is polynomial and this concludes the proof.  $\square$

The vertex version of [Theorem 5.34](#) follows by the application of [Theorem 5.3](#) to [Lemma 5.41](#) and [Lemma 5.7](#).

## 5.5 Negative results

Let us now here state several negative results on the Erdős–Pósa property of classes related to containment relations.

In the proofs below, we use the notion of *Euler genus* of a graph  $G$ . The *Euler genus* of a non-orientable surface  $\Sigma$  is equal to the non-orientable genus  $\tilde{g}(\Sigma)$  (or the crosscap number). The *Euler genus* of an orientable surface  $\Sigma$  is  $2g(\Sigma)$ , where  $g(\Sigma)$  is the orientable genus of  $\Sigma$ . We refer to the book of Mohar and Thomassen [MT01] for more details on graph embeddings. The *Euler genus* of a graph  $G$  is the minimum integer  $\gamma$  such that  $G$  can be embedded on a surface of the Euler genus  $\gamma$ .

**Lemma 5.42.** *Let  $H$  be a non-planar graph. Then  $\mathcal{T}(H)$  does not have the  $\nu$ -Erdős–Pósa property.*

*Proof.* Informally, we will construct, for every positive integer  $k$ , a graph  $G_k$  by “thickening” the vertices and edges of  $H$ . From the non-planarity of  $H$  and the way this graph is constructed, we will deduce that  $\nu\text{-pack}_{\mathcal{T}(H)}(G_k) = 1$ . On the other hand, the connectivity provided by the thickening of  $H$  will ensure that the removal of any  $k - 1$  vertices will leave a subdivision of  $H$  unaltered.

For every integers  $k > 0$  and  $d$ , we denote by  $\Gamma_{d,k}$  the graph obtained from a grid of width  $dk$  and height  $d + k - 1$  by adding  $k$  vertices  $a_1, \dots, a_k$  (that we call *apices*) and connecting  $a_1$  to the  $d$  first vertices on the first row of the grid (starting from the left),  $a_2$  to the  $d$  next vertices, and so on. For every  $i \in \llbracket 0, d - 1 \rrbracket$ , the set of vertices at indices  $\{ik + j, j \in \llbracket 0, k - 1 \rrbracket\}$  on the last row of  $\Gamma_{d,k}$  is called the  *$i$ -th port* of  $\Gamma_{d,k}$ . We will refer to the vertex at index  $ik + j$  of the last row as the  *$j$ -th vertex of the  $i$ -th port*. See Figure 5.2 for a drawing of  $\Gamma_{4,3}$ . On this drawing, the ports are  $U_0, \dots, U_3$ .

Let  $k$  be a positive integer. For every vertex  $v$  of  $H$ , we arbitrarily choose an ordering of its neighbors and we denote by  $\sigma_v(u)$  the rank of  $u$  in this ordering (that takes values from 0 to  $\deg(v) - 1$ ), for every neighbor  $u$  of  $v$ . We also let  $F_v$  be a copy of the graph  $\Gamma_{\deg(v),k}$ .

The graph  $G_k$  can be constructed from the disjoint union of the graphs of  $\{F_v, v \in V(H)\}$  by adding, for every pair  $u, v$  of adjacent vertices of  $H$ , the edge connecting the  $i$ -th vertex of the  $\sigma_v(u)$ -th port of  $F_v$  to the  $i$ -th vertex of the  $\sigma_u(v)$ -th port of  $F_u$ , for every  $i \in \llbracket 0, k - 1 \rrbracket$ . Informally, we connect the vertices of the  $\sigma_v(u)$ -th port of  $F_v$  to the vertices of the  $\sigma_u(v)$ -th port of  $F_u$  using “parallel” edges. Figure 5.3 depicts the graph  $G_k$  when  $G = K_5$  and  $k = 3$ .

The Euler genera of  $G_k$  and  $H$  are equal, however, as  $H$  is not planar, the Euler genus of the disjoint union of two copies of  $H$  is larger than that of  $H$  (see [BHK62]) and we get that  $\nu\text{-pack}_{\mathcal{T}(H)}(G_k) < 2$ . On the other hand, our construction ensures that  $\nu\text{-pack}_{\mathcal{T}(H)}(G_k) \geq 1$ .

Let us now show that for every subset  $X \subseteq V(G_k)$  with  $|X| < k$  we have  $\nu\text{-pack}_{\mathcal{T}(H)}(G \setminus X) \geq 1$ . This would complete the proof, since  $\{G_k, k \in \mathbb{N}_{\geq 1}\}$  would be an infinite family of graphs that have no  $\nu\text{-}\mathcal{T}(H)$ -packings of size 2 but where a minimum  $\nu\text{-}\mathcal{T}(H)$ -cover can be arbitrarily large.

Let  $u$  and  $v$  be two adjacent vertices of  $H$ , and let  $d = \deg(v)$ . For every  $i \in \llbracket 0, k - 1 \rrbracket$ , let  $C_i$  denote the vertices that are




Figure 5.2: The gadget  $\Gamma_{4,3}$  used in Lemma 5.42.

- either in the same column of  $F_u$  as the  $i$ -th vertex of the  $\sigma_u(v)$ -th port of  $F_u$ ;
- or in the same column of  $F_v$  as the  $i$ -th vertex of the  $\sigma_v(u)$ -th port of  $F_v$ .

The family  $\{C_i, i \in \llbracket 1, k \rrbracket\}$  contains  $k$  vertex disjoint elements, therefore at least one of them does not contain any vertex from  $X$  (as  $|X| < k$ ). Therefore, for every edge  $\{u, v\}$  of  $H$  there is an edge  $f(\{u, v\})$  between a vertex  $x$  of the  $\sigma_u(v)$ -th port of  $F_u$  and a vertex  $y$  of the  $\sigma_v(u)$ -th port of  $F_v$  such that no vertex of the same column as  $x$  in  $F_u$  (respectively  $y$  in  $F_v$ ) belong to  $X$ . Using the same argument we can show that for every vertex  $v \in V(H)$  there is an apex  $a$  such that the columns of  $F_v$  adjacent to  $a$  are free of vertices of  $X$ . Also we know that at least  $d$  rows do not contain vertices from  $X$ , as the grid of  $F_v$  has height  $d + k - 1$ . Therefore  $F_v$  contains as a subgraph a grid  $S_v$  such that:

1. an apex  $a$  is adjacent to  $d$  vertices of the first row of  $S_v$ ;
2. for every vertex  $u$  adjacent to  $v$ , the edge  $f(\{u, v\})$  shares one vertex the last row of  $S_v$ ;




Figure 5.3: The “thickened”  $K_5$  for  $k = 3$ . Edges with dashed ends are connected to the aligned edges at the opposite side of the figure. This graph contains a subdivision of  $K_5$  but not two vertex-disjoint ones, and the removal of any two vertices leaves one subdivision of  $K_5$  unaltered.

3. no vertex of the last row of  $S_v$  belong to two edges  $f(\{u, v\})$  and  $f(\{u', v\})$  for some distinct neighbors  $u, u'$  of  $v$ ;
4.  $S_v$  has height and width at least  $d$ ;
5.  $S_v$  does not contain any vertex of  $X$ .

We deduce that  $F_v \setminus X$  contains  $d$  paths  $P_0, \dots, P_{d-1}$  that have only the apex  $a$  as common vertex and such that  $P_i$  connects  $a$  to an endpoint of  $f(\{v, u_i\})$ , where  $u_i$  is

the neighbor of  $v$  of rank  $i$ , for every  $i \in \llbracket 0, d - 1 \rrbracket$ . It is now easy to see that the graph

$$G_k \left[ \bigcup_{v \in V(H)} \bigcup_{i=0}^{\deg_H(v)-1} V(P_i^v) \right]$$

contains a subdivision of  $H$  that does not contain any vertex of  $X$ . This concludes the proof.  $\square$

The proof of [Lemma 5.42](#) can be adapted to the setting of the edge-Erdős–Pósa property under the additional requirement that the pattern is subcubic.

**Lemma 5.43.** *Let  $H$  be a subcubic non-planar graph. Then  $\mathcal{T}(H)$  does not have the e-Erdős–Pósa property.*

*Proof.* Let  $k$  be a positive integer. We use the same construction of  $G_k$  as in the proof of [Lemma 5.42](#) with the following modifications: each vertex  $v$  of degree  $d \geq 4$  of  $G_k$  is replaced by a subcubic tree, the leaves of which are the neighbors of  $v$ . Let us call  $G'_k$  the graph we obtain. It is not hard to see that the genera of  $G'_k$  and  $G_k$  are equal. Moreover, as  $G'_k$  is subcubic, every e- $\mathcal{T}(H)$ -packing is also an v- $\mathcal{T}(H)$ -packing. We then obtain as previously that  $\text{e-pack}_{\mathcal{T}(H)}(G'_k) = 1$ . The arguments to show that  $\text{e-cover}_{\mathcal{T}(H)}(G'_k) \geq k$  are identical to the ones used in the proof of [Lemma 5.42](#).  $\square$

In fact, [Lemma 5.42](#) and [Lemma 5.43](#) can be used to prove that more general classes do not have the Erdős–Pósa property, as follows. As we will see in [Corollary 5.45](#) and [Corollary 5.46](#), the conditions of [Lemma 5.44](#) already encompass several well-studied classes.

**Lemma 5.44.** *Let  $x \in \{v, e\}$ , let  $H$  be a non-planar graph and let  $\mathcal{H}$  be a class of graphs such that:*

- (i)  $\mathcal{T}(H) \subseteq \mathcal{H}$ ; and
- (ii)  $H$  is graph of minimum Euler genus in  $\mathcal{H}$ ;
- (iii) if  $x = e$ , then  $H$  is subcubic.

*Then  $\mathcal{H}$  does not have the  $x$ -Erdős–Pósa property.*

*Proof.* Let  $k$  be a positive integer. We again consider the constructions of  $G_k$  and  $G'_k$  used in the proofs of [Lemma 5.42](#) and [Lemma 5.43](#). Let  $J_k$  be  $G_k$  if  $x = v$  and  $J_k = G'_k$  if  $x = e$ . Let show that  $\text{v-pack}_{\mathcal{H}}(J_k) = 1$ . For this, let us assume that there is an  $x$ - $\mathcal{T}(H)$ -packing  $F_1, \dots, F_p$ , for some  $p \in \mathbb{N}_{\geq 2}$  in  $J_k$ . It is crucial to note that in both the cases  $x = v$  and  $x = e$ , the subgraphs  $F_1, \dots, F_p$  are vertex-disjoint. In fact, when  $x = v$ , this follows from the definition of a v- $\mathcal{T}(H)$ -packing, and if  $x = e$  it is because  $G'_k$  is

subcubic. Then we have:

$$\begin{aligned}
\gamma(J_k) &\geq \gamma(F_1 \cup \dots \cup F_p) \\
&= \sum_{i=1}^p \gamma(F_i) && \text{(see [BHK62])} \\
&\geq p \cdot \gamma(H) && \text{(by minimality of } H) \\
\gamma(J_k) &> \gamma(H) && \text{(contradiction).}
\end{aligned}$$

We reached a contradiction, hence  $\nu\text{-pack}_{\mathcal{H}}(J_k) = 1$ . On the other hand,

$$\nu\text{-cover}_{\mathcal{H}}(J_k) \geq \nu\text{-cover}_{\mathcal{T}(H)}(J_k) \geq k.$$

The last inequality can be found in the proof of [Lemma 5.42](#) or [Lemma 5.43](#) (depending if  $x = \nu$  or  $x = e$ ). This concludes the proof.  $\square$

**Corollary 5.45** (proved in [\[RS86\]](#) for  $\mathcal{M}(H)$ ). *For every non-planar graph  $H$ , none of  $\mathcal{I}(H)$  and  $\mathcal{M}(H)$  have the  $\nu$ -Erdős–Pósa property.*

**Corollary 5.46.** *For every subcubic non-planar graph  $H$ , none of  $\mathcal{I}(H)$  and  $\mathcal{M}(H)$  have the  $e$ -Erdős–Pósa property.*

[Corollary 5.46](#) can be strengthened by dropping the degree condition on  $H$  when considering minor models of  $H$ , as follows.

**Lemma 5.47.** *For every non-planar graph  $H$ ,  $\mathcal{M}(H)$  does not have the  $e$ -Erdős–Pósa property.*

*Proof.* Let  $k$  be a positive integer. Again we use the graph  $G'_k$  constructed as in [Lemma 5.43](#). We modify it by replacing every apex  $a$  by a subcubic tree, the leaves of which are the neighbors of  $a$ . Let  $G''_k$  denote the graph that we obtain. Observe that  $G''_k$  is subcubic. Therefore, using the same argument as in the proof of [Lemma 5.43](#) we can show that  $e\text{-pack}_{\mathcal{M}(H)}(G) = 1$ . In the sequel we use the terminology of the proof of [Lemma 5.42](#). Let  $F''_v$  denote the graph obtained from  $F_v$  by replacing every vertex  $u$  of degree at least 4 by a subcubic tree, the leaves of which are the neighbors of  $u$ , for every  $v \in V(H)$ . The proof that  $e\text{-cover}_{\mathcal{M}(H)}(G) \geq k$  goes as in the proof of [Lemma 5.42](#), except that we obtain, for every  $v \in V(H)$ , that  $F''_v \setminus X$  contains a tree, the leaves of which are endpoints of  $f(\{v, u_i\})$  for  $i \in \llbracket 0, d-1 \rrbracket$  (instead of paths connecting an apex to endpoints of  $f(\{v, u_i\})$ ). Fortunately this is enough to guarantee that  $G''_k \setminus X$  contains  $H$  as a minor, and we are done.  $\square$

Thomassen in [\[Tho88\]](#) provided an example of a tree such that  $H \notin \mathcal{EP}_{\leq t.m.}^{\nu}$  (the same graph does not belong to  $\mathcal{EP}_{\leq t.m.}^e$  neither). Inspired by this construction we give another such graph that, additionally, is 2-connected.

**Lemma 5.48.** *There is a 2-connected (respectively 3-connected) planar graph that belongs to none of  $\mathcal{EP}_{\leq t.m.}^{\nu}$ ,  $\mathcal{EP}_{\leq t.m.}^e$ ,  $\mathcal{EP}_{\leq \text{imm.}}^{\nu}$ , and  $\mathcal{EP}_{\leq \text{imm.}}^e$ .*




Figure 5.4: A 2-connected graph  $H$  for which  $\mathcal{I}(H)$  does not have the (vertex or edge) Erdős–Pósa property.

*Informal proof.* This graph  $H$  is depicted in Figure 5.4. To see that  $H \notin \mathcal{EP}_{\leq t.m.}^v$  and  $H \notin \mathcal{EP}_{\leq t.m.}^e$ , consider as host graph  $G$  the graph in Figure 5.5. This graph consists of a main body that is a wall of height 3 and three triples of graphs attached at its upper, leftmost, and lower paths. Each of these triples consists of three copies of some of the 3-connected components of  $H$ . Notice that  $G$  does not contain more than one  $H$ -immersion expansion. However, in order to cover all  $H$ -immersion expansions of  $G$  one needs to remove at least 3 edges/vertices. By increasing the height of the wall of  $G$ , we may increase the minimum size of an  $\mathcal{I}(H)$ -vertex/edge cover while no  $\mathcal{I}(H)$ -vertex/edge packing of size greater than 1 will appear. It is easy to modify  $H$  so to make it 3-connected: just add a new vertex and make it adjacent with the tree vertices of degree 4. The resulting graph  $H'$  remains planar. The same arguments, applied to an easy modification of the host graph, can prove that  $H'$  is not a graph in  $\mathcal{H}^v$  or  $\mathcal{H}^e$ .  $\square$




Figure 5.5: The host graph  $G$ .

## 5.6 Summary of results

### 5.6.1 Results in terms of containment relations

For every partial order  $\preceq$  on graphs, and for every graph  $H$ , let

$$\mathcal{G}_{\preceq}(H) = \{G, H \preceq G\}.$$

For instance,  $\mathcal{G}_{\leq m}(H)$  is the class of all graphs containing  $H$  as a minor. For every  $x \in \{v, e\}$ , we define

$$\mathcal{EP}_{\succeq}^x = \{H, \mathcal{G}_{\succeq}(H) \text{ has the } x\text{-Erdős-Pósa property}\}$$

A general question on Erdős-Pósa properties is to characterize  $\mathcal{EP}_{\succeq}^x$  for several containment relations. We start with the following easy observation.

**Lemma 5.49.** *If  $\preceq$  is the subgraph or the induced subgraph relation,  $x \in \{v, e\}$ , and  $H$  is a non-trivial graph, then  $\mathcal{G}_{\preceq}(H)$  has the  $x$ -Erdős-Pósa property, with gap  $f : k \mapsto k \cdot |A_x(G)|$ . In other words,  $\mathcal{EP}_{\succeq}^x$  is the set of all graphs.*

*Proof.* Let  $H$  and  $G$  be two graphs and let  $k = x\text{-pack}_{\mathcal{G}_{\preceq}(H)}(G)$ . Let  $M_1, \dots, M_k$  be a  $v\text{-}\mathcal{G}_{\preceq}(H)$ -packing (respectively  $e\text{-}\mathcal{G}_{\preceq}(H)$ -packing) of size  $k$  with the minimal number of vertices (respectively edges). Observe that in this case,  $|M_i| = |H|$  (respectively  $\|M_i\| = \|H\|$ ) for every  $i \in \llbracket 1, k \rrbracket$ . Let  $X = \bigcup_{i=1}^k V(M_i)$  (respectively  $X = \bigcup_{i=1}^k E(M_i)$ ). As the packing we consider is of size  $k$ , the graph  $G \setminus X$  does not have any subgraph isomorphic to a member of  $\mathcal{G}_{\preceq}(H)$ . Hence  $X$  is an  $v\text{-}\mathcal{G}_{\preceq}(H)$ -cover (respectively  $e\text{-}\mathcal{G}_{\preceq}(H)$ -cover), and besides we have  $|X| = k \cdot |H|$  (respectively  $|X| = k \cdot \|H\|$ ).  $\square$

Notice that in case  $x = v$ , it is not necessary to demand that  $H$  is non-trivial in the statement of [Lemma 5.49](#).

Let us summarize known results that are related to the most common containment relations.

**Subgraphs and induced subgraphs:**  $\mathcal{EP}_{\succeq}^x$  is the class of all graphs, both for  $\preceq$  being the subgraph and induced subgraph relation, for every  $x \in \{v, e\}$  ([Lemma 5.49](#)).

**Minors:**  $\mathcal{EP}_{\leq m}^v$  is the class of planar graphs [[RS86](#), Theorem 8.2]. About the edge version, [Theorem 5.24](#), [Theorem 5.23](#), and [Theorem 5.29](#) imply that  $\mathcal{EP}_{\leq m}^e$  includes the class  $\{\theta_r\}_{r \in \mathbb{N}_{\geq 1}} \cup \{\theta_{r,r'}\}_{r,r' \in \mathbb{N}}$  and we show in [Lemma 5.47](#) that  $\mathcal{EP}_{\leq m}^e$  is a subclass of planar graphs.

**Topological Minors:**  $\mathcal{EP}_{\leq t.m.}^v$  has been characterized in the unpublished manuscript [[LPW14](#)]. There are trees, 2-connected and 3-connected graphs that belongs to none of  $\mathcal{EP}_{\leq t.m.}^v$  and  $\mathcal{EP}_{\leq t.m.}^e$  ([\[Tho88\]](#) and [Lemma 5.48](#)). The class  $\mathcal{EP}_{\leq t.m.}^v$  does not contain any non-planar graph ([Lemma 5.42](#)) and  $\mathcal{EP}_{\leq t.m.}^e$  does not contain any non-planar subcubic graph ([Lemma 5.43](#)).

**Immersion:** As proved in [Subsection 5.4.4](#),  $\mathcal{EP}_{\leq imm.}^v$  contains all planar subcubic graphs and  $\mathcal{EP}_{\leq imm.}^e$  contains all non-trivial, connected, planar subcubic graphs. Moreover,  $\mathcal{EP}_{\leq imm.}^v$  does not contain any non-planar graph ([Corollary 5.45](#)) and  $\mathcal{EP}_{\leq imm.}^e$  does not contain any subcubic non-planar graph ([Corollary 5.46](#)). On the other hand there is a 3-connected planar graph that belongs to none of  $\mathcal{EP}_{\leq imm.}^v$  and  $\mathcal{EP}_{\leq imm.}^e$ . [Lemma 5.48](#).

## 5.6.2 Results in terms of graph classes

In the following sections we list positive and negative results on the Erdős–Pósa property, and open problems.

Let us define the notation used in all the tables of [Section 5.6.2](#) and [Section 5.6.2](#). The fourth column of the tables gives the type of the packings/covers the current line is about. The character  $v$  (respectively  $e$ ) refers to vertex-disjoint (respectively edge-disjoint) packings and vertex (respectively edge) covers. We write  $v/e$  when the mentioned result holds for both the vertex and the edge version. The symbol  $v_{1/2}$  (resp.  $e_{1/2}$ ) indicates that the packing is allowed to use at most twice each vertex (resp. each edge) and that the cover contains vertices (resp. edges). Finally,  $w$  stands for vertex covers and packings where every vertex  $v$  of the host graph can be used at most  $w(v)$  times by every packing, where  $w$  is a function mapping reals to the vertices of the host graph. The more specific definitions are given in the corresponding sections.

### Positive results

We provide a series of tables presenting known results on the Erdős–Pósa property of some graph classes, sorted depending on the pattern. Results related to other structures (matroids, hypergraphs, geometry) and to fractional versions are not mentioned here.

A dash in the “gap” column means that the authors did not explicitly provided a gap function, even though one might be computable from the proof. The fourth column refers to the type of packing/cover, as defined above.

**Acyclic patterns.** Let  $G$  be a graph. For every  $S, T \subseteq V(G)$ , an  $(S, T)$ -*path* of  $G$  is a path with the one endpoint in  $S$  and the other one in  $T$ . An  $S$ -*path* is a path with both endpoints (which are distinct) in  $S$ . If  $S$  is a collection of subsets of  $V(G)$ , an  $S$ -*path* is a path that has endpoints in two different elements of  $S$ . A generalization of these settings have been introduced in [\[MW15\]](#), where the pairs of vertices that can be connected by a path are specified by an auxiliary graph. If  $S \subseteq V(G)$  and  $H$  (demand graph) is a graph with vertex set  $S$ , a path of  $G$  is said to be  $H$ -*valid* if its endpoints are adjacent vertices of  $H$ .

Ref.	Guest class $\mathcal{H}$	Host class $\mathcal{G}$	T.	Gap
[Kön31]	$K_2$	bipartite	v	$k$
[LY78]	directed cuts	any	e	$k$
[Men27]	$(S, T)$ -paths	any	v/e	$k$
[Grü38]	directed $(S, T)$ -paths	any (digraph)	v/e	$k$
[Gal64]	$S$ -paths	any	v	$2k$
[Mad78b]	$\mathcal{S}$ -paths	any	v	see [Sch01]
[Mad78a]	$\mathcal{S}$ -paths	any	e	see [SS04]
[CGG <sup>+</sup> 06]	non-zero directed $\mathcal{S}$ -paths	edge-group-labeled digraphs	v	$2k - 2$
[MW15]	$H$ -valid paths, $H$ with no matching of size $t$	any	v	$2^{2^{O(k+t)}}$
[FJW13a], Th. 5.19	$\mathcal{M}(H)$ , $H$ forest	any	v	$O_{\mathcal{H}}(k)$

**Triangles.** A graph is *flat* if every edge belongs to at most two triangles.

Ref.	Guest class $\mathcal{H}$	Host class $\mathcal{G}$	T.	Gap
[Tuz90]	triangles	planar graphs	e	$2k$
		$G$ with $\ G\  \geq 7 G ^2/16$	e	$2k$
		tripartite graphs	e	$7k/3$
[Kri95]	triangles	$\mathcal{T}(K_{3,3})$ -free graphs	e	$2k$
[HK88]	triangles	tripartite graphs	e	$1.956k$
[Hax99]	triangles	any	e	$(3 - \frac{3}{23})k$
[ALBT11]	triangles	odd-wheel-free graphs	e	$2k$
		4-colorable graphs		
[HKT11]	triangles	$K_4$ -free planar graphs	e	$3k/2$
		$K_4$ -free flat graphs		

**Cycles.** The statement of the results in [DZ02, DXZ03] requires additional definition. An odd ring is a graph obtained from an odd cycle by replacing every edge  $\{u, v\}$  by either a triangle, or three triangles  $uab$  and  $ucd$  together with the edges  $\{b, c\}$  and  $\{a, d\}$ . We denote by  $\mathcal{G}_1$  the class of graphs with no induced subdivision of the following:  $K_{2,3}$ , a wheel, or an odd ring. We denote by  $\mathcal{G}_2$  the class of graphs with no induced subdivision of the following:  $K_{3,3}$ , a wheel, or an odd ring.  $\vec{\mathcal{C}}_{\geq t}$  is the class of directed cycles of length at least  $t$ .

The results on directed cycles also need few more definitions. A digraph is *strongly planar* if it has a planar drawing such that for every vertex  $v$ , the edges with head  $v$

form an interval in the cyclic ordering of edges incident with  $v$  (definition from [GT10]). An *odd double circuit* is a digraph obtained from an undirected circuit of odd length more than 2 by replacing each edge by a pair of directed edges, one in each direction.  $F_7$  is the digraph obtained from the directed cycle on vertices  $v_1, \dots, v_7, v_1$ , by adding the edges creating the directed cycle  $v_1, v_3, v_5, v_7, v_2, v_4, v_6, v_1$ . We denote by  $\mathcal{F}$  the class of digraphs with no butterfly minor isomorphic to an odd double circuit, or  $F_7$  (for the definition of butterfly minors of digraphs see [GT10, JRST01, AKKW16]).

Results related to cycles with length constraints, with prescribed vertices, or to extensions of cycles are presented in the forthcoming tables.

Ref.	Guest class $\mathcal{H}$	Host class $\mathcal{G}$	T.	Gap
[EP65]	cycles	any	v	$O(k \log k)$
[Sim67]	cycles	any	v	$< (4 + o(1)) k \log k$
[Vos68]	cycles	any	v	$\leq (2 + o(1)) k \log k$
[Die05]	cycles	any	e	$(2 + o(1))k \log k$
[DZ02]	cycles	$\mathcal{G}_1$ , weighted	w	$k$
[DXZ03]	cycles	$\mathcal{G}_2$	v	$k$
[KLL02b]	cycles	planar graphs	v	$5k$
		outerplanar graphs	v	$2k$
[MYZ13]	cycles	planar graphs	v	$3k$
			e	$4k - 1$
[RRST96a]	directed cycles	any	v	–
[RS96]	directed cycles	planar	v	$O(k \log(k) \log \log k)$
[GT10]	directed cycles	strongly planar	v	$k$
		$\mathcal{F}$		
[LY78]	directed cycles	planar	e	$k$
[Sey96]	directed cycles	eulerian with a linkless embedding in 3-space	e	$k$
[HJW16]	cycles non homologous to zero	embedded graphs	$v_{1/2}$	–

**Cycles with length constraints.** The class of cycles of length at least  $t$  is referred to as  $\mathcal{C}_{\geq t}$ . For every positive integer  $k$  with, we say that a graph is  $k$ -near bipartite if every set  $X$  of vertices contains a stable of size at least  $|X|/2 - k$ .

Ref.	Guest class $\mathcal{H}$	Host class $\mathcal{G}$	T.	Gap
[Ree99]	odd cycles	planar graphs	v	superexponential
[FHRV05]	odd cycles	planar graphs	v e	$10k$ $2k$
[Tho01b]	odd cycles	$2^{3^{9k}}$ -connected graphs	v	$2k - 2$
[RR01]	odd cycles	$576k$ -connected graphs	v	$2k - 2$
[KR09]	odd cycles	$24k$ -connected graphs	v	$2k - 2$
[Ree99]	odd cycles	$k$ -near bipartite graphs	v	–
[KN07]	odd cycles	embeddable in an orientable surface of Euler genus $t$	v/e	–
[BR00]	odd cycles	any	e	–
[KV04]	odd cycles	planar graphs	e	$2k$
[KK12]	odd cycles	4-edge-connected graphs	e	$2^{2^{O(k \log k)}}$
[Ree99]	odd cycles	any	$v_{1/2}$	–
[Tho88]	cycles of length $0 \pmod t$	any	v	–
[KW05]	non-zero cycles	$(15k/2)$ -connected group-labeled graphs	v	$2k - 2$
[Wol11]	non-zero cycles	group-labeled graphs, c.f. [Wol11]	v	$c^{k^{c'}}$ for some $c, c'$
	cycles of non-zero length $\pmod{2t+1}$	any		
[HJW16]	doubly non-zero cycles, c.f. [HJW16]	doubly group-labeled graphs	$v_{1/2}$	–
	odd cycles non homologous to zero	embedded graphs		
[BBR07b]	$\mathcal{C}_{\geq t}$	any	v	$(13 + o_t(1))tk^2$
[FH14]	$\mathcal{C}_{\geq t}$	any	v	$(6t + 4 + o_t(1))k \log k$
[MNŠW16]	$\mathcal{C}_{\geq t}$	any	v	$6kt + (10 + o(1))k \log k$
[HM13]	$\vec{\mathcal{C}}_{\geq 3}$	any	v	–

**Extensions of cycles.** A *dumb-bell* is a graph obtained by connecting two cycles by a (non-trivial) path.

Ref.	Guest class $\mathcal{H}$	Host class $\mathcal{G}$	T.	Gap
[Sim67]	dumb-bells	any	v	$< (4000 + o(1))k \log k$
[FLM <sup>+</sup> 13]	$\mathcal{M}(\theta_t)$	any	v	$O(t^2 k^2)$
[FJS13]	$\mathcal{M}(\theta_t)$	any	v	$O_t(k \log k)$
[RST16]	$\mathcal{M}(\theta_t)$	any	e	$O(k^2 t^2 \text{polylog } kt)$ $O(k^4 t^2 \text{polylog } kt)$
[CRST15b], Th. 5.24	$\mathcal{M}(\theta_t)$	any	v/e	$O_t(k \log k)$

**Majors.** For every digraph  $D$ , we denote by  $\vec{\mathcal{M}}(D)$  (respectively  $\vec{\mathcal{T}}(G)$ ,  $\vec{\mathcal{I}}(G)$ ) the class of all digraphs that contain  $D$  as a directed minor (respectively directed topological minor, directed immersion). Refer to [CS11, CFS12, FS13] for a definition of these notions.

We also denote by  $\vec{\mathcal{M}}_{\bowtie}(D)$  (respectively  $\vec{\mathcal{T}}_{\bowtie}(G)$ ) the class of all digraphs that contain  $D$  as a butterfly-minor (respectively as a butterfly topological minor).  $\vec{\mathcal{P}}$  (respectively  $\vec{\mathcal{W}}$ ) is the class of all graphs that are butterfly minors of a cylindrical directed grid (respectively butterfly topological minors of a cylindrical directed wall). See for instance [AKKW16] for a definition of the cylindrical directed grid and wall and [JRST01, AKKW16] for a definition of butterfly (topological) minors.

For every  $s \in \mathbb{N}$ , a digraph is said to be  $s$ -semicomplete if for every vertex  $v$  there are at most  $s$  vertices that are not connected to  $v$  by an arc (in either direction). A semicomplete digraph is a 0-semicomplete digraph.

Ref.	Guest class $\mathcal{H}$	Host class $\mathcal{G}$	T.	Gap
[RS86], Lemma 5.12	$\mathcal{M}(H)$ , $H$ planar	any $\{G, \text{tw}(G) \leq t\}$	v v	– $(t-1)(k \text{cc}(H) - 1)$
[DKW10]	$\mathcal{M}(K_t)$	$O(kt)$ -connected graphs	v	–
[FST11]	$\mathcal{M}(H)$ , $H$ planar connected	$K_t$ -minor free	v	$O_{H,t}(k)$
[RT13b]	$\mathcal{M}(H)$ , $\text{pw}(H) \leq 2$ and $H$ connected	any	v	$2^{O( H ^2)} \cdot k^2 \log k$
[CC13]+ [CC14], Cor. 5.9	$\mathcal{M}(H)$ , $H$ planar connected	any	v	$O( H ^{O(1)} \cdot k \text{polylog } k)$
[CRST15b], Lem. 5.15, Th. 5.29	$\mathcal{M}(H)$ , $H$ con- nected	$\{G, \text{tpw}(G) \leq t\}$	v/e	$O_{H,t}(k)$
	$\mathcal{M}(\theta_{t,t'})$	simple graphs	e	–
[AKKW16]	$\vec{\mathcal{M}}_{\bowtie}(H)$ , $H \in \vec{\mathcal{P}}$	any	v	–
[Ray16]	$\vec{\mathcal{M}}(H)$ , for any di- graph $H$	$t$ -semicomplete, for fixed $t$	v	–

**Subdivisions.** For every  $t \in \mathbb{N}$ ,  $\mathcal{T}_{(0 \bmod t)}(H)$  denotes the class of subdivisions of  $H$  where every edge is subdivided  $0 \bmod t$  times.  $\mathcal{L}$  is a graph class defined in the

(unpublished) manuscript [LPW14]. See the previous section for the definition of  $\vec{\mathcal{T}}(G)$  and  $\vec{\mathcal{W}}$ .

Ref.	Guest class $\mathcal{H}$	Host class $\mathcal{G}$	T.	Gap
[Tho88]	$\mathcal{T}_{(0 \bmod t)}(H)$ , $H$ planar subcubic	any	v	–
[LPW14]	$\mathcal{T}(H)$ , $H \in \mathcal{L}$	any	v	–
[AKKW16]	$\vec{\mathcal{T}}_{\boxtimes}(H)$ , $H \in \vec{\mathcal{W}}$	any	v	–
[Ray16]	$\vec{\mathcal{T}}(H)$ , for any digraph $H$	$t$ -semicomplete, for fixed $t$	v	–

**Immersion expansions.** A graph  $H$  is a *half-integral immersion* of a graph  $G$  if  $H$  is an immersion of the graph obtained by  $G$  after duplicating the multiplicity of all its edges. We denote by  $\mathcal{I}_{1/2}(H)$  the class of all graphs containing  $H$  as a half-integral immersion. See above the definition of  $\vec{\mathcal{I}}(G)$ .

Ref.	Guest class $\mathcal{H}$	Host class $\mathcal{G}$	T.	Gap
[Liu15]	$\mathcal{I}(H)$	4-edge-connected	e	–
[GKRT16], Th. 5.34 Lemma 5.40	$\mathcal{I}(H)$ , $H$ planar subcubic connected non-trivial	any	e	$(\ H\  \cdot k)^{O(1)}$
	$\mathcal{I}(H)$ , $H$ connected non-trivial	$\{G, \text{tpw}(G) \leq t\}$ $\{G, \text{tcw}(G) \leq t\}$	e	$\ H\  \cdot t^2 \cdot k$
[Liu15]	$\mathcal{I}_{1/2}(H)$	any	$e_{1/2}$	–
[Ray16]	$\vec{\mathcal{I}}(H)$ , $H$ strongly connected, $H \neq K_1$	semicomplete	v	$2^{O(k^2 \ H\ ^2)}$

**Patterns with prescribed vertices.** Let us first present the two settings of Erdős–Pósa problems with prescribed vertices that we want to deal with here. The first type is when the guest class consists of fixed subgraphs of the host graph. For instance, one can consider a family  $\mathcal{F}$  of (non necessarily disjoint) subtrees of a tree  $T$ , and compare the maximum number of disjoint elements in  $\mathcal{F}$  with the minimum number of vertices/edges of  $T$  meeting all elements of  $\mathcal{F}$ . We will refer to these guest classes by words indicating that we are dealing with substructures (like “subtrees”). We stress that in this setting, the host class is allowed to contain one subgraph  $F$  of the host graph, but not one other subgraph  $F'$  even if  $F$  and  $F'$  are isomorphic. For every positive integer  $t$ , a  $t$ -path is a disjoint union of  $t$  paths, and a  $t$ -subpath of a  $t$ -path  $G$  is a subgraph that has a connected intersection with every connected component of  $G$ . The concept of  $t$ -forests and  $t$ -subforests is defined similarly.

In order to introduce the second type of problem, we need the following definition. Let  $x \in \{v, e\}$ . If  $\mathcal{H}$  is a class of graphs,  $G$  is a graph and  $S \subseteq A_x(G)$ , then a  $S$ - $\mathcal{H}$ -subgraph of  $G$  is a subgraph of  $G$  isomorphic to some member of  $\mathcal{H}$  and that contain one edge/vertex of  $S$ . We are now interested in comparing, for every graph

$G$  and every  $S \subseteq A_x(G)$ , the maximum number of  $S$ - $\mathcal{H}$ -subgraph of  $G$  with the minimum number of elements of  $A_x(G)$  that meet all  $S$ - $\mathcal{H}$ -subgraphs of  $G$ . We refer to these problems by prefixing the guest class with an “ $S$ ” (like in “ $S$ -cycles”). The authors of [HJW16] consider  $(S_1, S_2)$ -cycles for  $S_1, S_2 \subseteq V(G)$ : such cycles must meet both of  $S$  and  $S'$ . A generalization of this type of problem has been introduced in [KM15]: instead of one set  $S$ , one considers three subsets  $S_1, S_2, S_3$  of  $V(G)$  and a  $(S_1, S_2, S_3)$ -subgraph is required to intersect at least two sets of  $S_1, S_2$  and  $S_3$ . Note that some results on patterns with prescribed vertices have been stated in the table on acyclic patterns.

Ref.	Guest class $\mathcal{H}$	Host class $\mathcal{G}$	T.	Gap
[HS58]	subpaths	paths	v	$k$
[GL69]	$t$ -subpaths	$t$ -paths	v	$O(k^{t!})$
	subgraphs $H$ with $\text{cc}(H) \leq t$	paths	v	–
	$t$ -subforests	$t$ -forests	v	–
[GL69]	subtrees of a tree	trees	v	$k$
[Kai97]	$t$ -subpaths	$t$ -paths	v	$(t^2 - t + 1)k$
[Alo98]	$t$ -subpaths	$t$ -paths	v	$2t^2k$
[Alo02]	subgraphs $H$ with $\text{cc}(H) \leq t$	trees	v	$2t^2k$
	subgraphs $H$ with $\text{cc}(H) \leq t$	$\{G, \text{tw}(G) \leq w\}$	v	$2(w + 1)t^2k$
[KiKM11]	$S$ -cycles	any	v	$O(k^2 \log k)$
[PW12]	$S$ -cycles	any	v/e	$O(k \log k)$
[BJS14]	$S$ -cycles $\cap \mathcal{C}_{\geq t}$	any	v	$O(tk \log k)$
[Joo14]	odd $S$ -cycles	$50k$ -connected graphs	v	$O(k)$
[KK13]	odd $S$ -cycles	any	$v_{1/2}$	–
[KKKK13]	odd directed $S$ -cycles	any	$v_{1/2}$	–
[HJW16]	$(S_1, S_2)$ -cycles	any	v	–
[KM15]	$(S_1, S_2, S_3)$ - $\mathcal{M}(H)$ , $H$ planar	any	v	–

#### Classes with bounded parameters.

Ref.	Guest class $\mathcal{H}$	Host class $\mathcal{G}$	T.	Gap
[Tho88]	any family of connected graphs	$\{G, \text{tw}(G) \leq t\}$	v	$k(t + 1)$
[FJW13a]	$\{H, \text{pw}(H) \geq t\}$	any	v	$O_t(k)$
[CRST15b]	any finite family of connected graphs	$\{G, \text{tpw}(G) \leq t\}$	v/e	$O_t(k)$

## Negative results

The next table present lower bounds on the gap for several graph classes, as well as graph classes that do not have the Erdős–Pósa property. It indicates to which extend the results of the table of [Section 5.6.2](#) are best possible. The notation used here are the same as in the previous section, where they are defined.

### Cycles and paths.

Ref.	Guest class $\mathcal{H}$	Host class $\mathcal{G}$	T.	Gap
[Tuz90]	triangles	all graphs	e	$\geq 2k$
[EP65]	cycles	all graphs	v	$\Omega(k \log k)$
[Sim67]	cycles	all graphs	v	$> (\frac{1}{2} + o(1)) k \log k$
[Vos68]	cycles	all graphs	v	$\geq (\frac{1}{8} + o(1)) k \log k$
[KLL02b]	cycles	planar graphs	v	$\geq 2k$
[MYZ13]	cycles	planar graphs	e	$\geq 4k - c,$ $c \in \mathbb{N}$
[DL88]	odd cycles	all graphs	v	none
[Tho88]	cycles of length $p \bmod t,$ $p \in \llbracket 1, t - 1 \rrbracket$	all graphs	v	none
[Ree99]	odd cycles	all graphs	e	none
[Tho01b]	odd cycles	planar graphs	v	$\geq 2k - 2$
[KV04]	odd cycles	planar graphs	e	$\geq 2k$
[PW12]	$S$ -cycles	any	v	$\Omega(k \log k)$
[KK13]	odd $S$ -cycles	all graphs	v	none
[KKKK13]	odd directed $S$ -cycles	all graphs	v	none
[FH14]	$\mathcal{C}_{\geq t}$	all graphs	v	$\Omega(k \log k),$ $t$ fixed $\Omega(t), k$ fixed
[MNŠW16]	$\mathcal{C}_{\geq t}$	all graphs	v	$\geq (k - 1)t$ $\geq \frac{(k-1) \log k}{8}$
[Sim67]	dumb-bells	all graphs	v	$> (1 + o(1))k \log k$
[MW15]	$H$ -valid paths, $H$ with no matching of size $t$	all graphs	v	$f(k, t)$ (i.e. unavoidable dependency in $t$ )

**Patterns related to containment relations.**

Ref.	Guest class $\mathcal{H}$	Host class $\mathcal{G}$	T.	Gap
from [EP65]	$\mathcal{M}(H)$ , $H$ has a cycle	all graphs	v	$\Omega(k \log k)$
[RS86]	$\mathcal{M}(H)$ , $H$ non-planar	all graphs	v	none
Lemma 5.47	$\mathcal{M}(H)$ , $H$ non-planar	all graphs	e	none
Lemma 5.42	$\mathcal{T}(H)$ , $H$ non-planar	all graphs	v	none
[Tho88]	$\mathcal{T}_{(p \bmod t)}(H)$ , $H$ planar sub-cubic, $p \in \llbracket 1, t-1 \rrbracket$	all graphs	v	none
[Tho88]	$\mathcal{T}(H)$ , for infinitely many trees $H$ with $\Delta(H) = 4$	planar graphs	e	none
Lemma 5.43	$\mathcal{T}(H)$ , $H$ non-planar sub-cubic	all graphs	e	none
copying [Tho88]	$\mathcal{I}(H)$ , for infinitely many trees $H$ with $\Delta(H) = 4$	planar graphs	e	none
Cor. 5.45	$\mathcal{I}(H)$ , $H$ non-planar	all graphs	v	none
Cor. 5.46	$\mathcal{I}(H)$ , $H$ non-planar sub-cubic	all graphs	e	none
[GKRT16]	$\mathcal{I}(H)$ , for some 3-connected $H$ with $\Delta(H) = 4$	planar graphs	e	none
[Liu15]	$\mathcal{I}(H)$ , for every $H$	3-edge-connected graphs	e	none
[AKKW16]	$\vec{\mathcal{M}}(G)$ , $G \notin \vec{\mathcal{P}}$ $\vec{\mathcal{T}}(G)$ , $G \notin \vec{\mathcal{W}}$	all graphs	v	none

# CHAPTER 6

---

## FROM THE ERDŐS–PÓSA PROPERTY TO APPROXIMATION

In this chapter we show how the combinatorial connection between invariants of packing and covering provided by the Erdős–Pósa property can be used to design approximation algorithms. We focus on packing and covering graphs from  $\mathcal{M}(\theta_r)$  for any  $r \in \mathbb{N}$ , in both the vertex and edge setting. Drawing upon combinatorial results presented in [Chapter 4](#), we give an algorithmic proof that  $\mathcal{M}(\theta_r)$  has the vertex- and edge-Erdős–Pósa property with gap  $O(k \log k)$ , which is optimal. Using the algorithmic machinery of our proofs we introduce a unified approach for the design of an  $O(\log \text{OPT})$ -approximation algorithm for  $v\text{-pack}_{\theta_r}$ ,  $v\text{-cover}_{\theta_r}$ ,  $e\text{-pack}_{\theta_r}$ , and  $e\text{-cover}_{\theta_r}$  that runs in  $O(n \cdot \log(n) \cdot m)$  steps.

This chapter contains material that previously appeared in the following article: *An  $O(\log \text{OPT})$ -approximation for covering/packing minor models of  $\theta_r$* , co-authored with Dimitris Chatzidimitriou, Ignasi Sau, and Dimitrios M. Thilikos, WAOA 2015, Patras, Greece, 2015 [[CRST15b](#), [CRST15a](#)].

### 6.1 Introduction

From the algorithmic point of view, the computation of  $x\text{-pack}_{\mathcal{H}}$  (for  $x \in \{v, e\}$ ) corresponds to the general family of graph packing problems, while the computation of  $x\text{-cover}_{\mathcal{H}}$  belongs to the general family of graph modification problems where the modification operation is the removal of vertices/edges (depending on whether  $x = v$  or  $x = e$ ). Interestingly, particular instantiations of  $\mathcal{H} = \mathcal{M}(\theta_r)$  generate known, well studied, NP-hard problems. For instance, asking whether  $v\text{-cover}_{\mathcal{M}(\theta_r)} \leq k$  generates VERTEX COVER for  $r = 1$ , FEEDBACK VERTEX SET for  $r = 2$ , and DIAMOND HITTING SET for  $r = 3$  [[FJP10](#), [FLMS12](#)]. Moreover, asking whether  $x\text{-pack}_{\mathcal{M}(\theta_r)}(G) \geq k$  corresponds to VERTEX CYCLE PACKING [[BTY11](#), [KLL02a](#)] and EDGE CYCLE PACKING [[ACR03](#), [KNS<sup>+</sup>07](#)] when  $x = v$  and  $x = e$ , respectively. Finally, asking whether

$|E(G)| - \text{e-cover}_{\mathcal{M}(\theta_r)}(G) \leq k$  corresponds to the MAXIMUM CACTUS SUBGRAPH<sup>1</sup>. All parameters keep being NP-complete to compute because the aforementioned base cases can be reduced to the general one by replacing each edge by one of multiplicity  $r - 1$ .

From the approximation point of view, it was proven in [FLMS12] that, when  $H$  is a planar graph, there is a randomized polynomial  $O(1)$ -approximation algorithm for  $\text{v-cover}_{\mathcal{M}(H)}$ . For the cases of  $\text{v-cover}_{\mathcal{M}(\theta_r)}$  and  $\text{v-pack}_{\mathcal{M}(\theta_r)}$ ,  $O(\log n)$ -approximations are known for every  $r \geq 1$  because of [JPS<sup>+</sup>11, JPS<sup>+</sup>14] (see also [SV05]). Moreover,  $\text{v-cover}_{\mathcal{M}(\theta_3)}$  admits a deterministic 9-approximation [FJP10]. About the edge variant, it is known, from [KNY05], that there is a polynomial  $O(\sqrt{\log n})$ -approximation algorithm for  $\text{e-pack}_{\mathcal{M}(\theta_2)}(G)$ . Notice also that it is trivial to compute  $\text{e-cover}_{\mathcal{M}(\theta_1)}(G)$  in polynomial time. However, to our knowledge, nothing is known about the computation of  $\text{e-cover}_{\mathcal{M}(\theta_r)}(G)$  for  $r \geq 3$ .

In this section we introduce a unified approach for the study of the combinatorial interconnections and the approximability of the parameters  $\text{v-cover}_{\mathcal{M}(\theta_r)}$ ,  $\text{e-cover}_{\mathcal{M}(\theta_r)}$ ,  $\text{v-pack}_{\mathcal{M}(\theta_r)}$ , and  $\text{e-pack}_{\mathcal{M}(\theta_r)}$ . Our main combinatorial result is the following theorem, already stated in the previous chapter.

**Theorem 5.24.** *There is a function  $f_r(k) = O_r(k \log k)$  such that for every  $x \in \{v, e\}$  and every  $r \in \mathbb{N}$ ,  $\mathcal{M}(\theta_r)$  has the  $x$ -Erdős–Pósa property with gap  $f_r$ .*

Our proof is unified and treats simultaneously the covering and the packing parameters for both the vertex and the edge cases. This verifies the optimal combinatorial bound for the case where  $x = v$  [FJS13] and optimally improves (in terms of  $k$ ) the bound given in Subsection 5.4.2 (which appeared in [RST16]) for the case where  $x = e$ . In this section, when giving the running time of an algorithm with input some graph  $G$ , we agree that  $n = |V(G)|$  and  $m = |E(G)|$ . Based on the proof of Theorem 5.24, we prove the following algorithmic result.

**Theorem 6.1.** *For every  $r \in \mathbb{N}_{\geq 2}$  and every  $x \in \{v, e\}$ , there exists an  $O(n \cdot \log(n) \cdot m)$ -step algorithm that, given a graph  $G$ , outputs an  $O(\log \text{OPT})$ -approximation for  $x$ -cover $_{\mathcal{M}(\theta_r)}$  and  $x$ -pack $_{\mathcal{M}(\theta_r)}$ .*

Theorem 6.1 improves the results in [JPS<sup>+</sup>11, JPS<sup>+</sup>14] for the cases of  $\text{v-cover}_{\mathcal{M}(\theta_r)}$  and  $\text{v-pack}_{\mathcal{M}(\theta_r)}$  and is, to our knowledge, the first approximation algorithm for  $\text{e-cover}_{\mathcal{M}(\theta_r)}$  and  $\text{e-pack}_{\mathcal{M}(\theta_r)}$  for  $r \geq 3$ .

**Overview of the proof.** Our proofs are based on the notion of *partitioned protrusion* that, roughly, is a tree-structured subgraph of  $G$  with small boundary to the rest of  $G$  (see Subsection 6.2.2 for the precise definition). Partitioned protrusions, that we met in Section 4.1 under the name of *edge-protrusions*, can be seen as the edge-analogue of the notion of protrusions introduced in [BFL<sup>+</sup>09a] (see also [BFL<sup>+</sup>09b]).

<sup>1</sup>The MAXIMUM CACTUS SUBGRAPH problem asks, given a graph  $G$  and an integer  $k$ , whether  $G$  contains a subgraph with  $k$  edges where no two cycles share an edge. It can be reduced to the VERTEX CYCLE PACKING problem on cubic graphs which, in turn, can be proved to be NP-complete using a simple variant of the NP-completeness proof of EXACT COVER BY 2-SETS [Gol15].

Our approach makes strong use of the main result of [Section 4.1](#), that is equivalently stated as [Theorem 4.4](#) in this section. According to this result, there exists a polynomial algorithm that, given a graph  $G$  and an integer  $k$  as an input, outputs one of the following:

1. a collection of  $k$  edge/vertex disjoint  $\theta_r$ -majors of  $G$ ,
2. a  $\theta_r$ -major  $J$  with  $O(\log k)$  edges, or
3. a large partitioned protrusion of  $G$ .

Our approximation algorithm does the following for each  $k \leq |V(G)|$ . If the first case of the above combinatorial result applies on  $G$ , we can safely output a packing of  $k$   $\theta_r$ -majors in  $G$ . In the second case, we make some progress as we may remove the vertices/edges of  $J$  from  $G$  and then set  $k := k - 1$ . In order to deal with the third case, we prove that in a graph  $G$  with a sufficiently large partitioned protrusion, we can either find some  $\theta_r$ -major with  $O(\log k)$  edges (which is the same as in the second case), or we can replace it by a smaller graph where both  $\text{x-cover}_{\mathcal{M}(\theta_r)}$  and  $\text{x-pack}_{\mathcal{M}(\theta_r)}$  remain invariant ([Lemma 6.2](#)). The proof that such a reduction is possible is given in [Section 6.3](#) and is based on a suitable dynamic programming encoding of partial packings and coverings that is designed to work on partitioned protrusions.

Notice that the “essential step” in the above procedure is the second case that reduces the packing number of the current graph by 1 to the price of reducing the covering number by  $O(\log k)$ . This is the main argument (previously used in [\[FJW13b\]](#)) that supports the claimed  $O(\log \text{OPT})$ -approximation algorithm ([Theorem 6.1](#)) and the corresponding Erdős–Pósa relations in [Theorem 5.24](#).

**Organization of the chapter.** In [Section 6.2](#) we provide all concepts and notation that we use in our proofs. [Section 6.3](#) contains the proof of [Lemma 6.2](#), which is the main technical part of the chapter. The presentation and analysis of our approximation algorithm is done in [Section 6.4](#), where [Theorem 5.24](#) and [Theorem 6.1](#) are proven.

## 6.2 Definitions specific to this section

### 6.2.1 Basic definitions

Recall that we use  $n(G)$  and  $m(G)$  to denote  $|G|$  and  $\|G\|$ , respectively, for every graph  $G$ . If  $\mathcal{H}$  is a finite collection of graphs, we set  $n(\mathcal{H}) = \sum_{H \in \mathcal{H}} n(H)$ ,  $m(\mathcal{H}) = \sum_{H \in \mathcal{H}} m(H)$ .

Given a graph  $H$  and a graph  $J$  that are both subgraphs of the same graph  $G$ , we define the subgraph  $H \cap_G J$  of  $G$  as the graph  $(V(H) \cap V(J), E(H) \cap E(J))$ .

Given a graph  $G$  and a set  $S \subseteq V(G)$ , such that all vertices in  $S$  have degree 2 in  $G$ , we define  $\text{diss}(G, S)$  as the graph obtained from  $G$  after we dissolve in it all vertices in  $S$ .

**Topological minors.** Whereas the results presented in this chapter deal with majors, the tools that we provide in [Section 6.3](#) are expressed in the setting of subdivisions. We show in [Section 6.4](#) how this more general setting can be applied to majors. If  $G$  is a graph and  $\mathcal{H}$  is a finite collection of connected graphs, recall that an  $\mathcal{H}$ -subdivision in  $G$  is a subgraph  $M$  of  $G$  that is a subdivision of a graph, denoted by  $\hat{M}$  in this chapter, that is isomorphic to a member of  $\mathcal{H}$ . Clearly, the vertices of  $\hat{M}$  are vertices of  $G$  and its edges correspond to paths in  $G$  between their endpoints such that internal vertices of a path do not appear in any other path. We refer to the vertices of  $\hat{M}$  in  $G$  as the *branch vertices* of the  $\mathcal{H}$ -subdivision  $M$ , whereas internal vertices of the paths between branch vertices are called *subdivision vertices* of  $M$ .

**Approximation algorithms.** A way to approach problems that are believed to be computationally hard is via approximation algorithms. The main idea is to trade accuracy for speed. As approximation algorithms appear very locally in this thesis, we only give the few required definitions.

In an *optimization problem*, one is typically asked the minimum (or maximum, depending on the problem) size of an object satisfying predefined properties.

For instance, in the Independent Set optimization problem, one is asked, for a graph  $G$ , the maximum size of a subset of  $V(G)$  that consist of pairwise non-independent vertices.

Informally, an *approximation algorithm* for an optimization problem is an algorithm that computes an approximate solution for this problem. In order to give a guarantee on the performance of the algorithm, we use the following definition: for a constant  $c \in \mathbb{R}$ ,  $c > 1$ , we say that an algorithm is a *c-approximation* for a given problem if on every input it returns a solution that is within a factor  $c$  of the optimal solution. If, for some function  $f: \mathbb{N} \rightarrow \mathbb{R}$ , the algorithm returns on every input of size  $n$  a solution which is within a factor  $f(\text{OPT})$  (resp.  $f(n)$ ) of the optimal solution  $\text{OPT}$ , then we call it an  $f(\text{OPT})$ -approximation (resp.  $f(n)$ -approximation).

## 6.2.2 Boundaried graphs

Informally, a boundaried graph is used to represent a graph that has been obtained by “dissecting” a larger graph along some of its edges, where the boundary vertices correspond to edges that have been cut. In this section we formally define boundaried graphs and related notions. We also give a notion of equivalence that is a cornerstone of our algorithms.

**Boundaried graphs** A *boundaried graph*  $\mathbf{G} = (G, B, \lambda)$  is a triple consisting of a graph  $G$ , a set  $B$  of vertices of degree one (called *boundary*), and a bijection  $\lambda$  from  $B$  to a subset of  $\mathbb{N}_{\geq 1}$ . The edges with at least one endpoint in  $B$  are called *boundary edges*. We define  $E^s(G)$  as the subset of  $E(G)$  of boundary edges. We stress that instead of  $\mathbb{N}_{\geq 1}$  we could choose any other set of symbols to label the vertices of  $B$ . We denote the set of labels of  $\mathbf{G}$  by  $\Lambda(\mathbf{G}) = \lambda(B)$ . Given a collection  $\mathcal{H}$  of graphs, we say that a  $\mathbf{G}$  is  $\mathcal{H}$ -free if  $G \setminus B$  is  $\mathcal{H}$ -free.

Two boundaried graphs  $\mathbf{G}_1$  and  $\mathbf{G}_2$  are *compatible* if  $\Lambda(\mathbf{G}_1) = \Lambda(\mathbf{G}_2)$ . Let now  $\mathbf{G}_1 = (G_1, B_1, \lambda_1)$  and  $\mathbf{G}_2 = (G_2, B_2, \lambda_2)$  be two compatible boundaried graphs. We define the graph  $\mathbf{G}_1 \oplus \mathbf{G}_2$  as the graph obtained by first taking the disjoint union of  $G_1$  and  $G_2$ , then, for every  $i \in \Lambda(\mathbf{G}_1)$ , identifying  $\lambda_1^{-1}(i)$  with  $\lambda_2^{-1}(i)$ , and finally dissolving all resulting identified vertices. Suppose that  $e$  is an edge of  $G = \mathbf{G}_1 \oplus \mathbf{G}_2$  that was created after dissolving the vertex resulting from the identification of a vertex  $v_1$  in  $B_1$  and a vertex  $v_2$  in  $B_2$  and that  $e_i$  is the boundary edge of  $G_i$  that has  $v_i$  as endpoint, for  $i = 1, 2$ . Then we say that  $e$  is the *heir* of  $e_i$  in  $G$ , for  $i = 1, 2$ , and we denote this by  $\text{heir}_G(e_i)$ . For  $i \in \{1, 2\}$ , if  $S \subseteq E(G_i)$ , then

$$\text{heir}_G(S) = (E(G_i) \cap S) \cup \{\text{heir}_G(e) \mid e \in E^s(G_i) \cap S\}.$$

For reasons of notational consistency, if  $V \subseteq V(G_i)$ , we denote  $\text{heir}_G(V) = V$ .

Figure 6.1 shows the result of the operation  $\oplus$  on two graphs. Boundaries are drawn in gray and their labels are written next to them. The graphs  $\mathbf{G}_1$  and  $\mathbf{G}_2$  on this picture are compatible as  $\Lambda(\mathbf{G}_1) = \Lambda(\mathbf{G}_2) = \{0, 1, 2, 3\}$ .




Figure 6.1: Gluing graphs together:  $G = \mathbf{G}_1 \oplus \mathbf{G}_2$ .

For every  $t \in \mathbb{N}_{\geq 1}$ , we denote by  $\mathcal{B}_t$  all boundaried graphs whose boundary is labeled by numbers in  $\llbracket 1, t \rrbracket$ . Given a boundaried graph  $\mathbf{G} = (G, B, \lambda)$  and a subset  $S$  of  $V(G)$  such that all vertices in  $S$  have degree 2 in  $G$ , we define  $\text{diss}(\mathbf{G}, S)$  as the graph  $\hat{\mathbf{G}} = (\hat{G}, B, \lambda)$  where  $\hat{G} = \text{diss}(G, S)$ .

Let  $W$  be a graph and  $S$  be a non-empty subset of  $V(W)$ . An  $S$ -splitting of  $W$  is a pair  $(\mathbf{G}_S, \mathbf{G}_{S^c})$  consisting of two boundaried graphs  $\mathbf{G}_S = (G_S, B_S, \lambda_S)$  and  $\mathbf{G}_{S^c} = (G_{S^c}, B_{S^c}, \lambda_{S^c})$  that can be obtained as follows: First, let  $W^+$  be the graph obtained by subdividing in  $W$  every edge with one endpoint in  $S$  and the other in  $V(W) \setminus S$  and let  $B$  be the set of created vertices. Let  $\lambda$  be any bijection from  $B$  to a subset of  $\mathbb{N}_{\geq 1}$ . Then  $G_S = W^+[S \cup B]$ ,  $G_{S^c} = W^+ \setminus S$ ,  $B_S = B_{S^c} = B$ , and  $\lambda_S = \lambda_{S^c} = \lambda$ . Notice that there are infinite such pairs, depending on the numbers that are assigned to the boundaries of  $\mathbf{G}_S$  and  $\mathbf{G}_{S^c}$ . Moreover, keep in mind that all the boundary edges of  $G_S$  are non-loop edges with exactly one endpoint in  $B$  and the same holds for the boundary edges of  $G_{S^c}$ . An example of a splitting is given in Figure 6.2, where boundaries are depicted by gray vertices.

We say that  $\mathbf{G}' = (G', B', \lambda')$  is a *boundaried subgraph* of  $\mathbf{G} = (G, B, \lambda)$  if  $G'$  is a subgraph of  $G$ ,  $B' \subseteq B$  and  $\lambda' = \lambda|_{B'}$ . On the other hand,  $\mathbf{G}$  is a subgraph of a (non-boundaried) graph  $H$  if  $\mathbf{G} = \mathbf{H}_S$  for some  $S$ -splitting  $(\mathbf{H}_S, \mathbf{H}_{S^c})$ , where  $S \subseteq V(H)$ .




Figure 6.2: Cutting a graph:  $(\mathbf{G}_S, \mathbf{G}_{S^c})$  is an  $S$ -splitting of  $W$ , where  $S$  consists of all the white vertices.

If  $H$  is a graph,  $G$  is a subgraph of  $H$ , and  $\mathbf{F} = (F, B, \lambda)$  is a boundaried subgraph of  $H$ , we define  $G \cap_H \mathbf{F}$  as follows. Let  $S = V(G) \cap (V(F) \setminus B)$  and let  $G^+$  be the graph obtained by subdividing once every edge of  $G$  that has one endpoint in  $S$  and the other in  $V(G) \setminus S$ . We call  $B'$  the set of created vertices and let  $G' = G^+[S \cup B']$ . Then  $G'$  is a subgraph of  $F$  where  $B' \subseteq B$ . For every  $v \in B'$ , we set  $\lambda'(v) = \lambda(v)$ , which is allowed according to the previous remark. Then  $G \cap_H \mathbf{F} = (G', B', \lambda')$ . Observe that  $G \cap_H \mathbf{F}$  is one side of an  $S$ -splitting of  $G$ .

Given two boundaried graphs  $\mathbf{G}' = (G', B', \lambda')$  and  $\mathbf{G} = (G, B, \lambda)$ , we say that they are *isomorphic* if there is an isomorphism from  $G'$  to  $G$  that respects the labelings of  $B$  and  $B'$ , i.e., maps every vertex  $x \in B'$  to  $\lambda^{-1}(\lambda'(x)) \in B$ . Given a boundaried graph  $\mathbf{G} = (G, B, \lambda)$ , we denote  $n(\mathbf{G}) = n(G) - |B|$  and  $m(\mathbf{G}) = m(G)$ .

Given a boundaried graph  $\mathbf{G} = (G, B, \lambda)$  and an  $x \in \{v, e\}$ , we set  $A_x(\mathbf{G}) = V(G) \setminus B$  or  $A_x(\mathbf{G}) = E(G)$ , depending on whether  $x = v$  or  $x = e$ .

**Partial structures.** Let  $\mathcal{F}$  be a finite family of connected graphs. A boundaried subgraph  $\mathbf{J}$  of a boundaried graph  $\mathbf{G}$  is a *partial  $\mathcal{F}$ -subdivision* if there is a boundaried graph  $\mathbf{H}$  which is compatible with  $\mathbf{G}$  and a boundaried subgraph  $\mathbf{J}'$  of  $\mathbf{H}$  which is compatible with  $\mathbf{J}$  such that  $\mathbf{J} \oplus \mathbf{J}'$  is an  $\mathcal{F}$ -subdivision of  $\mathbf{G} \oplus \mathbf{H}$ . Intuitively, this means that  $\mathbf{J}$  can be extended into an  $\mathcal{F}$ -subdivision in some larger graph. In this case, the  $\mathcal{F}$ -subdivision  $\mathbf{J} \oplus \mathbf{J}'$  is said to be an *extension* of  $\mathbf{J}$ .

Similarly, for every  $p \in \mathbb{N}_{\geq 1}$ , a collection of boundaried subgraphs  $\mathcal{J} = \{\mathbf{J}_1, \dots, \mathbf{J}_p\}$  of a graph  $\mathbf{G}$  is a *partial  $x$ - $\mathcal{F}$ -packing* if there is a boundaried graph  $\mathbf{H}$  which is compatible with  $\mathbf{G}$  and a collection of boundaried subgraphs  $\{\mathbf{J}'_1, \dots, \mathbf{J}'_p\}$  of  $\mathbf{H}$  such that  $\{\mathbf{J}_1 \oplus \mathbf{J}'_1, \dots, \mathbf{J}_p \oplus \mathbf{J}'_p\}$  is an  $x$ - $\mathcal{F}$ -packing of  $\mathbf{G} \oplus \mathbf{H}$ . The obtained packing is said to be an *extension* of the partial packing  $\mathcal{J}$ . A partial packing is  $\mathcal{T}(\mathcal{F})$ -free if none of its members is an  $F$ -subdivision for some  $F \in \mathcal{F}$ . Observe that since every graph in  $\mathcal{F}$  is connected, every partial subdivision of an  $\mathcal{T}(\mathcal{F})$ -free partial packing in  $\mathbf{G}$  must contain at least one boundary vertex of  $\mathbf{G}$ .

**Partitions and protrusions.** In order to decompose graphs along edge cuts, we introduce the following edge-counterpart of the notion of (*vertex*) *protrusion* introduced in [BFL<sup>+</sup>09a, BFL<sup>+</sup>09b].

Given a rooted tree-partition  $\mathcal{D} = (T, s, \mathcal{X})$  of  $G$  and a vertex  $i \in V(T)$ , we define

$$T_i = T[\text{desc}_{T,s}(i)], \quad V_i = \bigcup_{h \in V(T_i)} X_h, \quad \text{and} \quad G_i = G[V_i].$$

Let  $W$  be a graph and  $t \in \mathbb{N}_{\geq 1}$ . A pair  $\mathbf{P} = (\mathbf{G}, \mathcal{D})$  is a  $t$ -partitioned protrusion of  $W$  if there exists an  $S \subseteq V(W)$  such that

- $\mathbf{G} = (G, B, \lambda)$  is a boundaried graph where  $\mathbf{G} \in \mathcal{B}_t$  and  $\mathbf{G} = \mathbf{G}_S$  for some  $S$ -splitting  $(\mathbf{G}_S, \mathbf{G}_{S^c})$  of  $W$  and
- $\mathcal{D} = (T, s, \{X_u\}_{u \in V(T)})$  is a rooted tree-partition of  $G \setminus B$  of width at most  $t$ , where  $X_s$  is the set of neighbors in  $G$  of the vertices in  $B$ .

Given a family  $\mathcal{F}$  of graphs, we say that a  $t$ -partitioned protrusion  $(\mathbf{G}, \mathcal{D})$  of a graph  $W$  is  $\mathcal{F}$ -free if  $\mathbf{G}$  is  $\mathcal{F}$ -free. For every vertex  $u \in V(T)$ , we also define the  $t$ -partitioned protrusion  $\mathbf{P}_u$  of  $W$  as a pair  $\mathbf{P}_u = (\mathbf{G}_u, \mathcal{D}_u)$ , where  $\mathcal{D}_u = (T_u, u, \{X_v\}_{v \in V_u})$  and  $\mathbf{G}_u = \mathbf{G}_{V_u}$  for some  $V_u$ -splitting  $(\mathbf{G}_{V_u}, \mathbf{G}_{V_u^c})$  of  $W$ . We choose the labeling function of  $\mathbf{G}_s$  so that it is the same as the one of  $\mathbf{G}$ , i.e.,  $\mathbf{G}_s = \mathbf{G}$ . Notice that the labelings of all other  $\mathbf{G}_u$ 's are arbitrary. For every  $u \in V(T)$  we define

$$\mathcal{G}_u = \{\mathbf{G}_l\}_{l \in \text{children}_{(T,s)}(u)}.$$

### 6.2.3 Encodings, signatures, and folios

In this section we introduce tools that we will use to sort boundaried graphs depending on the subdivisions that are realizable inside.

**Encodings.** Let  $\mathcal{H}$  be a family of graphs, let  $t \in \mathbb{N}_{\geq 1}$ , and let  $x \in \{v, e\}$ . If  $\mathbf{G} = (G, B, \lambda) \in \mathcal{B}_t$  is a boundaried graph and  $S \subseteq A_x(\mathbf{G})$ , we define  $\text{pp}_{\mathcal{H}}^x(\mathbf{G}, S)$  (pp as the initials of *partial packings*) as the collection of all sets  $\{(\mathbf{J}_1, L_1), \dots, (\mathbf{J}_\sigma, L_\sigma)\}$  such that

- $\{\mathbf{J}_1, \dots, \mathbf{J}_\sigma\}$  is a partial  $x$ - $\mathcal{T}(\mathcal{H})$ -packing of  $G \setminus S$  of size  $\sigma$  and
- $L_i = V(\hat{M}_i) \cap V(G)$ , where  $M_i$  is an extension of  $J_i$ , for every  $i \in \llbracket 1, \sigma \rrbracket$ .

In other words,  $L_i$  contains branch vertices of the partial  $\mathcal{H}$ -subdivision  $\mathbf{J}_i$  for every  $i \in \llbracket 1, \sigma \rrbracket$  (see [Figure 6.3](#) and [Figure 6.4](#)). The set  $\text{pp}_{\mathcal{H}}^x(\mathbf{G}, S)$  encodes all different restrictions in  $\mathbf{G}$  of partial  $x$ - $\mathcal{H}$ -packings that avoid the set  $S$ . Given a boundaried graph  $\mathbf{G} = (G, B, \lambda)$  and a set  $L \subseteq V(G)$  such that every vertex of  $V(G) \setminus L$  has degree 2 in  $G$ , we define  $\kappa(\mathbf{G}, L)$  as the boundaried graph obtained from  $\mathbf{G}$  by dissolving every vertex of  $V(G) \setminus L$ , i.e.,  $\kappa(\mathbf{G}, L) = (\text{diss}(\mathbf{G}, V(G) \setminus L), B, \lambda)$ . In the definition of  $\kappa$  we assume that the boundary vertices of  $\kappa(\mathbf{G}, L)$  remain the same as in  $\mathbf{G}$  while the other vertices are treated as new vertices (see [Figure 6.5](#)).

This allows us to introduce the following notation aimed at representing, intuitively, the essential part of each partial packing:

$$\text{cpp}_{\mathcal{H}}^x(\mathbf{G}, S) = \{\hat{\mathcal{J}} = \{\hat{\mathbf{J}}_1, \dots, \hat{\mathbf{J}}_\sigma\} = \{\kappa(\mathbf{J}_1, L_1), \dots, \kappa(\mathbf{J}_\sigma, L_\sigma)\} \mid \{(\mathbf{J}_1, L_1), \dots, (\mathbf{J}_\sigma, L_\sigma)\} \in \text{pp}_{\mathcal{H}}(\mathbf{G}, S)\}$$

(here, cpp is mnemonic for *compressed partial packings*).




Figure 6.3: An  $e\mathcal{T}(H)$ -packing in  $\mathbf{G}$ , where branch vertices are circled.




Figure 6.4: A partial subdivision from the packing of [Figure 6.3](#), where  $L$  is the set of subdivision vertices.

**Isomorphisms.** If  $\mathbf{G} = (G, B, \lambda)$  and  $\mathbf{G}' = (G', B', \lambda')$  are two compatible boundaried graphs in  $\mathcal{B}_t$ ,  $S \in V(G)$ , and  $S' \in V(G')$ , we say that a member  $\hat{J}$  of  $\text{cpp}_{\mathcal{H}}^{\times}(\mathbf{G}, S)$  and a member  $\hat{J}'$  of  $\text{cpp}_{\mathcal{H}}^{\times}(\mathbf{G}', S')$  are *isomorphic* if there is a bijection between them such that paired elements are isomorphic. We also say that  $\text{cpp}_{\mathcal{H}}^{\times}(\mathbf{G}, S)$  and  $\text{cpp}_{\mathcal{H}}^{\times}(\mathbf{G}', S')$  are *isomorphic* if there is a bijection between them such that paired elements are isomorphic.

We now come to the point where we can define, for every boundaried graph, a signature encoding all the possible partial packings that can be realized in this graph.

**Signatures and folios.** For every  $y \in \mathbb{N}$ , we set

$$\text{sig}_{\mathcal{H}}^{\times}(\mathbf{G}, y) = \{\text{cpp}_{\mathcal{H}}^{\times}(\mathbf{G}, S), S \subseteq A_{\times}(G), |S| = y\}$$

and, given two compatible boundaried graphs  $\mathbf{G}, \mathbf{G}' \in \mathcal{B}_t$  and a  $y \in \mathbb{N}$ , we say that  $\text{sig}_{\mathcal{H}}^{\times}(\mathbf{G}, y)$  and  $\text{sig}_{\mathcal{H}}^{\times}(\mathbf{G}', y)$  are *isomorphic* if there is a bijection between them such that paired elements are isomorphic.

Finally, for  $\rho \in \mathbb{N}$ , we set

$$\text{folio}_{\mathcal{H}, \rho}(\mathbf{G}) = (\text{sig}_{\mathcal{H}}^{\vee}(\mathbf{G}, 0), \dots, \text{sig}_{\mathcal{H}}^{\vee}(\mathbf{G}, \rho), \text{sig}_{\mathcal{H}}^{\text{e}}(\mathbf{G}, 0), \dots, \text{sig}_{\mathcal{H}}^{\text{e}}(\mathbf{G}, \rho)).$$

Given two  $\mathbf{G}, \mathbf{G}' \in \mathcal{B}_t$ , a  $\rho \in \mathbb{N}$ , and a finite collection of connected graphs  $\mathcal{H}$ , we say that  $\mathbf{G} \simeq_{\mathcal{H}, \rho} \mathbf{G}'$  if  $\mathbf{G}$  and  $\mathbf{G}'$  are compatible, neither  $\mathbf{G}$  nor  $\mathbf{G}'$  contains an  $\mathcal{H}$ -subdivision, and the elements of  $\text{folio}_{\mathcal{H}, \rho}(\mathbf{G})$  and  $\text{folio}_{\mathcal{H}, \rho}(\mathbf{G}')$  are coordinate-wise isomorphic.




Figure 6.5: The *compression* of the partial packing of Figure 6.4:  $\hat{\mathbf{J}} = \kappa(\mathbf{J}, L)$ .

### 6.3 The reduction procedure

Let  $\mathcal{H}$  be a finite set of connected graphs. In this section we show that one can, in linear time, either find a *small*  $\mathcal{H}$ -subdivision in a  $t$ -partitionned protrusion, or *reduce* it so that the parameters of packing and covering (wrt.  $\mathcal{H}$ -subdivisions) remain unchanged. More formally, the purpose of this section is to prove the following lemma.

**Lemma 6.2.** *There exists a function  $f_6: \mathbb{N}^2 \rightarrow \mathbb{N}$  and an algorithm that, given a positive integer  $t$ , a finite collection  $\mathcal{H}$  of connected graphs where  $h = m(\mathcal{H})$ , and a  $t$ -partitionned protrusion  $\mathbf{P} = (\mathbf{G}, (T, s, \mathcal{X}))$  of a graph  $W$  with  $n(\mathbf{G}) > f_6(h, t)$ , outputs either*

- an  $\mathcal{H}$ -subdivision of  $W$  with at most  $f_6(h, t)$  edges, or
- a graph  $W'$  such that

$$\begin{aligned} \text{x-pack}_{\mathcal{T}(\mathcal{H})}(W') &= \text{x-pack}_{\mathcal{T}(\mathcal{H})}(W), \\ \text{x-cover}_{\mathcal{T}(\mathcal{H})}(W') &= \text{x-cover}_{\mathcal{T}(\mathcal{H})}(W), \text{ and} \\ n(W') &< n(W). \end{aligned}$$

Furthermore, this algorithm runs in time  $O_{t,h}(n(T))$ .

Before giving the proof of Lemma 6.2, we need to prove several intermediate results. In the sequel, unless stated otherwise, we assume that  $x \in \{v, e\}$ ,  $t \in \mathbb{N}_{\geq 1}$  and that  $\mathcal{H}$  is a finite collection of connected graphs. We set  $h = m(\mathcal{H})$ .

**Lemma 6.3.** *There are two functions  $f_7: \mathbb{N}^2 \rightarrow \mathbb{N}$  and  $f_8: \mathbb{N}^2 \rightarrow \mathbb{N}$  such that, for every graph  $W$  and every  $t$ -partitionned protrusion  $(\mathbf{G}, (T, s, \mathcal{X}))$  of  $W$ , if  $\mathcal{P}$  is an  $\mathcal{H}$ -free partial  $x$ - $\mathcal{H}$ -packing in  $\mathbf{G}$  then:*

- (a) *The partial subdivisions of graphs of  $\mathcal{H}$  that are contained in  $\mathcal{P}$  have in total at most  $f_7(h, t)$  branch vertices.*
- (b)  *$\mathcal{P}$  intersects at most  $f_8(h, t)$  graphs of  $\mathcal{G}_s$ .*

*Proof.* Proof of (a). First, note that  $\mathcal{P}$  has cardinality at most  $t$ . Indeed, since every element of  $\mathcal{P}$  is a partial subdivision (because the packing is  $\mathcal{T}(\mathcal{H})$ -free) of a connected graph, it contains a boundary edge of  $\mathbf{G}$  (which by definition has degree one). Also, two distinct partial subdivisions in  $\mathcal{P}$  are (at least) edge-disjoint. Finally, each of

these partial subdivisions contains at most  $\max_{H \in \mathcal{H}} n(H) \leq h$  branch vertices. Consequently, the number of branch vertices of graphs of  $\mathcal{H}$  induced by the elements of  $\mathcal{P}$  in  $\mathbf{G}$  is at most  $t \cdot h$ . Hence the function  $f_7(h, t) := t \cdot h$  upper-bounds the amount of branch vertices each  $\mathcal{T}(\mathcal{H})$ -free partial packing can contain.

Proof of (b). Let  $\zeta$  be the maximum multiplicity of an edge in a graph of  $\mathcal{H}$ . Because of (a),  $\mathcal{P}$  has at most  $f_7(h, t)$  branch vertices of graphs of  $\mathcal{H}$ , so at most  $f_7(h, t)$  graphs of  $\mathcal{G}_s$  may contain such vertices. Besides,  $\mathcal{P}$  might also contain paths free of branch vertices linking pairs of branch vertices. Since there are at most  $(f_7(h, t))^2$  such pairs and no pair will need to be connected with more than  $\zeta \leq h$  distinct paths, it follows that at most  $(f_7(h, t))^2 \cdot h$  graphs of  $\mathcal{G}_s$  contain vertices from these paths. Therefore, the elements of  $\mathcal{P}$  intersects all together at most  $f_7(h, t) + (f_7(h, t))^2 \cdot h =: f_8(h, t)$  graphs of  $\mathcal{G}_s$ .  $\square$

**Lemma 6.4.** *The size of the image of the function  $\text{cpp}_{\mathcal{H}}^{\times}$ , when its domain is restricted to*

$$\{(\mathbf{G}, S), \mathbf{G} \in \mathcal{B}_t \text{ is } \mathcal{T}(\mathcal{H})\text{-free and } S \subseteq A_{\times}(\mathbf{G})\},$$

*is upper-bounded by a function of  $h$  and  $t$ .*

*Proof.* Let  $\mathbf{G} \in \mathcal{B}_t$  be  $\mathcal{T}(\mathcal{H})$ -free and let  $S \subseteq A_{\times}(\mathbf{G})$ . By Lemma 6.3(a), every  $\mathcal{T}(\mathcal{H})$ -free partial  $\times$ - $\mathcal{H}$ -packing in  $\mathbf{G}$  contains at most  $f_7(h, t)$  branch vertices. A partial packing may in addition use at most  $t$  boundary vertices. Let  $\mathcal{C}_{h,t}$  be the class of all boundaried graphs of  $\mathcal{B}_t$  on at most  $f_7(h, t) + t$  vertices. Clearly the size of this class is a function depending on  $h$  and  $t$  only. Recall that the elements of the set  $\text{cpp}_{\mathcal{H}}^{\times}(\mathbf{G}, S)$  are obtained from partial  $\times$ - $\mathcal{H}$ -packings by dissolving internal vertices of the paths linking branch vertices, hence every element of  $\text{cpp}_{\mathcal{H}}^{\times}(\mathbf{G}, S)$  is a boundaried graph of  $\mathcal{B}_t$  having at most  $f_7(h, t) + t$  vertices. Therefore, for any  $\mathcal{T}(\mathcal{H})$ -free boundaried graph  $\mathbf{G} \in \mathcal{B}_t$  and subset  $S \subseteq A_{\times}(\mathbf{G})$ , we have  $\text{cpp}_{\mathcal{H}}^{\times}(\mathbf{G}, S) \subseteq \mathcal{C}_{h,t}$ . As a consequence, the image of the function  $\text{cpp}_{\mathcal{H}}^{\times}$  when restricted to  $\mathcal{T}(\mathcal{H})$ -free boundaried graphs  $\mathbf{G} \in \mathcal{B}_t$  (and subsets  $S \subseteq A_{\times}(\mathbf{G})$ ) is a subset of the power set of  $\mathcal{C}_{h,t}$ , so its size is upper-bounded by a function that depends only on  $h$  and  $t$ .  $\square$

**Corollary 6.5.** *There is a function  $f_9: \mathbb{N}^2 \rightarrow \mathbb{N}$  such that the relation  $\simeq_{\mathcal{H},t}$  partitions  $\mathcal{T}(\mathcal{H})$ -free boundaried graphs of  $\mathcal{B}_t$  into at most  $f_9(h, t)$  equivalence classes.*

The following follows directly from the definition of  $\text{cpp}_{\mathcal{H}}^{\times}$ .

*Remark 6.6.* Let  $\mathbf{F}, \mathbf{G} \in \mathcal{B}_t$  be two compatible boundaried graphs and let  $k \in \mathbb{N}$ . The following are equivalent:

- $\mathbf{F} \oplus \mathbf{G}$  has an  $\times$ - $\mathcal{T}(H)$ -packing of size  $k$ ;
- there is a  $\hat{\mathcal{J}} \in \text{cpp}_{\mathcal{H}}^{\times}(\mathbf{G}, \emptyset)$  such that  $\mathbf{F} \oplus \mathbf{U}\hat{\mathcal{J}}$  has an  $\times$ - $\mathcal{T}(H)$ -packing of size  $k$ .

The choice of the definition of the relation  $\simeq$  is justified by the following lemma. Informally speaking, it states that we can replace a  $t$ -partitioned protrusion of a graph with any other  $\simeq_{\mathcal{H},t}$ -equivalent  $t$ -partitioned protrusion without changing the covering and packing number of the graph. The reduction algorithm that we give after this lemma relies on this powerful property.

**Lemma 6.7** (protrusion replacement). *Let  $\mathbf{F}, \mathbf{G}, \mathbf{G}' \in \mathcal{B}_t$  be three compatible boundaried graphs such that  $\mathbf{G} \simeq_{\mathcal{H},t} \mathbf{G}'$ . For every  $k \in \mathbb{N}$ , we have:*

- (i) *there is an  $x\text{-}\mathcal{T}(H)$ -packing of size  $k$  in  $\mathbf{F} \oplus \mathbf{G}$  iff there is one in  $\mathbf{F} \oplus \mathbf{G}'$ ; and*
- (ii) *there is an  $x\text{-}\mathcal{T}(H)$ -cover of size  $k$  in  $\mathbf{F} \oplus \mathbf{G}$  iff there is one in  $\mathbf{F} \oplus \mathbf{G}'$ .*

*Proof.* Proof of (i), “ $\Rightarrow$ ”. Let  $\mathcal{M}$  be an  $x\text{-}\mathcal{T}(\mathcal{H})$ -packing of size at least  $k$  in  $\mathbf{F} \oplus \mathbf{G}$ , whose set of branch vertices is  $L$ . We define

$$\begin{aligned} \mathbf{J}_{\mathbf{F}} &= (\mathbf{U}\mathcal{M}) \cap_{\mathbf{F} \oplus \mathbf{G}} \mathbf{F}, \\ \mathbf{J}_{\mathbf{G}} &= (\mathbf{U}\mathcal{M}) \cap_{\mathbf{F} \oplus \mathbf{G}} \mathbf{G}, \text{ and} \\ \hat{\mathbf{J}}_{\mathbf{G}} &= \bigcup_{M \in \mathcal{M}} \kappa(M \cap_{\mathbf{F} \oplus \mathbf{G}} \mathbf{G}, L \cap V(G)). \end{aligned}$$

Note that  $\hat{\mathbf{J}}_{\mathbf{G}} \in \text{cpp}_{\mathcal{H}}^x(\mathbf{G}, \emptyset)$  and that  $\mathbf{F} \oplus \hat{\mathbf{J}}_{\mathbf{G}}$  has an  $x\text{-}\mathcal{T}(\mathcal{H})$ -packing of size at least  $k$  (cf. **Remark 6.6**). By definition of  $\simeq$ , there is a bijection  $\psi$  between  $\text{cpp}_{\mathcal{H}}^x(\mathbf{G}, \emptyset)$  and  $\text{cpp}_{\mathcal{H}}^x(\mathbf{G}', \emptyset)$ . Let  $\hat{\mathbf{J}}'_{\mathbf{G}}$  be the image of  $\hat{\mathbf{J}}_{\mathbf{G}}$  by  $\psi$ . Since  $\hat{\mathbf{J}}'_{\mathbf{G}}$  and  $\hat{\mathbf{J}}_{\mathbf{G}}$  are isomorphic,  $\mathbf{F} \oplus \hat{\mathbf{J}}'_{\mathbf{G}}$  also has an  $x\text{-}\mathcal{T}(\mathcal{H})$ -packing of size at least  $k$ . By **Remark 6.6**, this implies that such a packing exists in  $\mathbf{F} \oplus \mathbf{G}'$  as well. The direction “ $\Leftarrow$ ” is symmetric as  $\mathbf{G}$  and  $\mathbf{G}'$  play the same role.

Proof of (ii), “ $\Rightarrow$ ”. Let  $C \subseteq A_x(\mathbf{F} \oplus \mathbf{G})$  be a minimum  $x\text{-}\mathcal{H}$ -covering of  $\mathbf{F} \oplus \mathbf{G}$  of size at most  $k$ . Let  $S = C \cap A_x(\mathbf{G})$ . Since we assume that  $\mathbf{G}$  is  $\mathcal{T}(\mathcal{H})$ -free and that  $C$  is minimum, we can also assume that  $|S| \leq t$  (otherwise we could get a smaller covering by taking the  $t$  boundary vertices/edges of  $\mathbf{G}$ ). By our assumption that  $\mathbf{G} \simeq_{\mathcal{H},t} \mathbf{G}'$ , there is an isomorphism between  $\text{sig}_{\mathcal{H}}^x(\mathbf{G}, |S|)$  and  $\text{sig}_{\mathcal{H}}^x(\mathbf{G}', |S|)$ . Let  $S' \subseteq A_x(\mathbf{G}')$  be a set such that  $\text{cpp}_{\mathcal{H}}^x(\mathbf{G}, S)$  is sent to  $\text{cpp}_{\mathcal{H}}^x(\mathbf{G}', S')$  by this isomorphism. Then observe that every partial packing  $\mathcal{J}'$  of  $\mathbf{G}' \setminus S'$ , such that  $(\mathbf{F} \setminus C) \oplus (\mathbf{U}\mathcal{J}')$  has an  $\mathcal{H}$ -subdivision, can be translated into a partial packing  $\mathcal{J}$  of  $\mathbf{G} \setminus S$  such that  $(\mathbf{F} \setminus C) \oplus (\mathbf{U}\mathcal{J})$  also has such a subdivision, in the same way as in the proof of (i) above. As  $C$  is a cover, this would lead to contradiction. Therefore  $\text{pp}_{\mathcal{H}}^x(\mathbf{G}, S)$  does not contain such a partial packing. As a consequence,  $C \cap A_x(\mathbf{F}) \cup S'$  is a covering of  $\mathbf{F} \oplus \mathbf{G}'$  of size at most  $k$ . As in the previous case, the proof of direction “ $\Leftarrow$ ” comes from the symmetry in the statement.  $\square$

**Lemma 6.7** can be rewritten as follows.

**Corollary 6.8.** *Under the assumptions of **Lemma 6.7**, we have  $x\text{-pack}(\mathbf{F} \oplus \mathbf{G}) = x\text{-pack}(\mathbf{F} \oplus \mathbf{G}')$  and  $x\text{-cover}(\mathbf{F} \oplus \mathbf{G}) = x\text{-cover}(\mathbf{F} \oplus \mathbf{G}')$ .*

Recall that  $f_9(h, t)$  denotes the number of equivalence classes of  $\simeq_{\mathcal{H},t}$  among boundaried graphs of  $\mathcal{B}_t$ . For every  $h, t \in \mathbb{N}$ , let  $f_{10}(h, t) = f_9(h, t) \cdot f_8(h, t)$  and let  $f_6(h, t) = 2ht^3 \cdot (f_{10}(h, t))^{f_9(h, t)+1}$ . Let us give some intuition about these definitions. The first remark is an application of the pigeonhole principle.

*Remark 6.9.* In a collection of more than  $f_{10}(h, t)$   $\mathcal{T}(\mathcal{H})$ -free boundaried graphs of  $\mathcal{B}_t$ , there is one that is equivalent (w.r.t.  $\simeq_{\mathcal{H},t}$ ) to  $f_8(h, t)$  other graphs of the collection.

**Lemma 6.10.** *If  $(T, s, \mathcal{X})$  is a rooted tree-partition of a graph  $G$  with the following properties:*

- $(T, \mathcal{X})$  has width at most  $t$ ;
- $T$  has height at most  $f_9(h, t)$ ; and
- $T$  has degree at most  $f_{10}(h, t) + 1$ ,

*then  $G$  has at most  $f_6(h, t)$  vertices, and every  $\mathcal{H}$ -subdivision of  $G$  has at most  $f_6(h, t)$  edges.*

*Proof.* Indeed, the above assumptions imply that  $T$  has at most  $(f_{10}(h, t))^{f_9(h, t)+1}$  vertices. Every bag of  $(T, \mathcal{X})$  contains at most  $t$  vertices of  $G$ , therefore  $G$  has at most  $(f_{10}(h, t))^{f_9(h, t)+1} \cdot t \leq f_6(h, t)$  vertices. Also, every bag induces a subgraph with at most  $t(t-1)/2$  multiedges (i.e. without counting multiplicities), and for every edge  $f$  of  $T$  we have  $|E_f| \leq t$ , hence every bag contributes for at most  $t^2 + t$  multiedges. Therefore  $G$  has at most  $(t+t^2)(f_{10}(h, t))^{f_9(h, t)+1}$  multiedges. Now, observe that a multiedge of  $G$  is used at most  $h$  times by an  $\mathcal{H}$ -subdivision, since every path connecting two branch vertices of a subdivision uses a given multiedge at most once. We deduce that an  $\mathcal{H}$ -subdivision of  $G$  contains at most  $h \cdot (t+t^2)(f_{10}(h, t))^{f_9(h, t)+1} \leq f_6(h, t)$  edges.  $\square$

The two next lemmas are the main tools used in the proof of [Lemma 6.2](#). Under different conditions, they provide either a small subdivision, or a reduced graph. [Lemma 6.11](#) considers the case where a vertex of  $T$  has high degree, whereas [Lemma 6.12](#) deals with the situation where  $T$  has a long path.

**Lemma 6.11.** *There is an algorithm that, given a  $t$ -partitioned protrusion  $\mathbf{P} = (\mathbf{G}, (T, s, \mathcal{X}))$ , and a vertex  $u \in V(T)$  with more than  $f_{10}(h, t)$  children such that for every  $v \in \text{children}_{(T, s)}(u)$ , we have  $m(\mathbf{G}_v) \leq f_6(h, t)$ , outputs either*

- an  $\mathcal{H}$ -subdivision with at most  $f_6(h, t)$  edges, or
- a graph  $W'$  such that

$$\begin{aligned} \text{x-pack}_{\mathcal{T}(\mathcal{H})}(W') &= \text{x-pack}_{\mathcal{T}(\mathcal{H})}(W), \\ \text{x-cover}_{\mathcal{T}(\mathcal{H})}(W') &= \text{x-cover}_{\mathcal{T}(\mathcal{H})}(W), \text{ and} \\ n(W') &< n(W). \end{aligned}$$

*Moreover, this algorithm runs in  $O_{h, t}(1)$  steps.*

*Proof.* If  $\mathbf{G}_v$  contains an  $\mathcal{H}$ -subdivision for some child  $v$  of  $u$ , then this subdivision has at most  $f_6(h, t)$  edges and we are done. Therefore we now consider the case where  $\mathbf{G}_v$  is  $\mathcal{H}$ -free for every child  $v$  of  $u$ . This allows us to consider the folios of these boundaryed graphs.

As  $u$  has more than  $f_{10}(h, t)$  children, it contains a collection of  $d = f_8(h, t) + 1$  children  $v_1, \dots, v_d$ , such that  $\mathbf{G}_{v_1} \simeq_{\mathcal{H}, t} \mathbf{G}_{v_i}$  for every  $i \in \llbracket 2, d \rrbracket$  (by [Remark 6.9](#)). Since every  $\text{x-}\mathcal{T}(\mathcal{H})$ -packing of  $W$  will intersect at most  $f_8(h, t)$  bags of children of  $u$

(by [Lemma 6.3\(b\)](#)), we can safely delete one of the  $f_8(h, t) + 1$  equivalent subgraphs mentioned above.

We use the following procedure in order to identify such a subgraph to delete or a small  $\mathcal{H}$ -subdivision;

1. let  $A$  be an array of  $f_9(h, t)$  counters initialized to 0, each corresponding to a distinct equivalence class of  $\simeq_{\mathcal{H}, t}$ ;
2. pick a vertex  $v \in \text{children}_{(T, s)}(u)$  that has not been considered yet;
3. if  $G_v$  contains an  $\mathcal{H}$ -subdivision  $M$ , then return  $M$  and exit;
4. otherwise, increment the counter of  $A$  corresponding to the equivalence class of  $G_v$  by one;
5. if this counter reaches  $d + 1$ , return  $v$ , otherwise go back to [line 2](#).

Notice that the subdivision returned in [line 3](#) has size at most  $f_6(h, t)$  as mentioned above, and that the vertex returned in [line 5](#) has the desired property. The relation  $\simeq_{\mathcal{H}, t}$  has at most  $f_9(h, t)$  equivalence classes ([Corollary 6.5](#)), thus the main loop will be run at most  $f_9(h, t) \cdot f_8(h, t) + 1$  times (by the pigeonhole principle). Eventually, [lines 3](#) and [4](#) can be performed in  $O_{h, t}(1)$ -time given that  $G_v$  has size bounded by a function of  $h$  and  $t$ .

In the end, we return  $W' = W \setminus V(G_v)$  if the algorithm above outputs  $v$  and  $M$  otherwise.  $\square$

**Lemma 6.12.** *There is an algorithm that, given a  $t$ -partitioned protrusion  $\mathbf{P} = (\mathbf{G}, (T, s, \mathcal{X}))$  of a graph  $W$  and a vertex  $u \in V(T)$  of height  $f_9(h, t)$  in  $(T, s)$  such that  $T_u$  has maximum degree at most  $f_{10}(h, t) + 1$ , outputs either*

- an  $\mathcal{H}$ -subdivision with at most  $f_6(h, t)$  edges, or
- a graph  $W'$  such that

$$\begin{aligned} \text{x-pack}_{\mathcal{T}(\mathcal{H})}(W') &= \text{x-pack}_{\mathcal{T}(\mathcal{H})}(W), \\ \text{x-cover}_{\mathcal{T}(\mathcal{H})}(W') &= \text{x-cover}_{\mathcal{T}(\mathcal{H})}(W), \text{ and} \\ n(W') &< n(W). \end{aligned}$$

Moreover, this algorithm runs in  $O_{h, t}(1)$ -time.

*Proof.* As in the proof of [Lemma 6.10](#), we use the fact that every  $\mathcal{H}$ -subdivision of  $G_u$  uses every multiedge at most  $h$  times. A consequence is that the bounded subgraph of  $G_u$  obtained by setting the multiplicity of every multiedge  $e$  to  $\min(\text{mult}(e), h)$  contains an  $\mathcal{H}$ -subdivision iff  $G_u$  does. As the number of vertices and edges of this subgraph is bounded by a function of  $h$  and  $t$ , we can therefore check in  $O_{h, t}(1)$ -time if  $G_u$  contains an  $\mathcal{H}$ -subdivision. If one is found, it has at most  $f_6(h, t)$  edges ([Lemma 6.10](#)) and we are done.

Let us now consider the case where  $\mathbf{G}_u$  is  $\mathcal{T}(\mathcal{H})$ -free. By definition of vertex  $u$ , there is a path on  $f_9(h, t) + 1$  vertices from a leaf of  $T_u$  to  $u$ . Let us arbitrarily choose, for every vertex  $v$  of this path, a  $V_v$ -splitting  $(\mathbf{G}_{G_v}, \mathbf{G}_{G_v^c})$  of  $\mathbf{G}$ . By definition of  $f_9(h, t)$  (the number of equivalence classes in  $\simeq_{\mathcal{H}, t}$  in  $\mathcal{B}_t$ ), there are two distinct vertices  $v, w$  on this path such that  $\mathbf{G}_v \simeq_{\mathcal{H}, t} \mathbf{G}_w$ . As mentioned above, the number edges of  $\mathbf{G}_u$  is bounded by a function of  $h$  and  $t$ , hence finding these two vertices can be done in  $O_{h,t}(1)$ -time. Let us assume without loss of generality that  $s$  is closer to  $v$  than  $w$ . Let  $\mathbf{H}$  be the boundaried graph such that  $W = \mathbf{H} \oplus \mathbf{G}_{G_v}$  and let  $W' = \mathbf{H} \oplus \mathbf{G}_{G_w}$ . By [Corollary 6.8](#), we have  $\text{x-pack}_{\mathcal{T}(\mathcal{H})}(W') = \text{x-pack}_{\mathcal{T}(\mathcal{H})}(W)$  and  $\text{x-cover}_{\mathcal{T}(\mathcal{H})}(W') = \text{x-cover}_{\mathcal{T}(\mathcal{H})}(W)$ . Furthermore, the graph  $W'$  is clearly smaller than  $W$ .  $\square$

We are now ready to prove [Lemma 6.2](#). Let us first recall its statement.

**Lemma 6.2.** *There exists a function  $f_{11}: \mathbb{N}^2 \rightarrow \mathbb{N}$  and an algorithm that, given a positive integer  $t$ , a finite collection  $\mathcal{H}$  of connected graphs where  $h = m(\mathcal{H})$ , and a  $t$ -partitioned protrusion  $\mathbf{P} = (\mathbf{G}, (T, s, \mathcal{X}))$  of a graph  $W$  with  $n(\mathbf{G}) > f_6(h, t)$ , outputs either*

- an  $\mathcal{H}$ -subdivision of  $W$  with at most  $f_6(h, t)$  edges, or
- a graph  $W'$  such that

$$\begin{aligned} \text{x-pack}_{\mathcal{T}(\mathcal{H})}(W') &= \text{x-pack}_{\mathcal{T}(\mathcal{H})}(W), \\ \text{x-cover}_{\mathcal{T}(\mathcal{H})}(W') &= \text{x-cover}_{\mathcal{T}(\mathcal{H})}(W), \text{ and} \\ n(W') &< n(W). \end{aligned}$$

Furthermore, this algorithm runs in time  $O_{t,h}(n(T))$ .

*Proof.* Let us consider the following procedure:

1. by a DFS on  $(T, s)$ , compute the height of each vertex of  $T$  and find (if it exists) a vertex  $v$  of degree more than  $f_{10}(h, t) + 1$  and height at most  $f_9(h, t) - 1$  that has minimum height;
2. if such a vertex  $v$  is found, then apply the algorithm of [Lemma 6.11](#) on  $\mathbf{P}$  and  $v$ , and return the obtained result;
3. otherwise, find a vertex  $u$  of height exactly  $f_9(h, t)$  in  $(T, s)$  and then apply the algorithm of [Lemma 6.12](#) on  $\mathbf{P}$  and  $(T_u, u)$  and return the obtained result.

Observe that since  $n(G) > f_6(h, t)$ , [Lemma 6.10](#) implies that either  $T$  has diameter more than  $f_9(h, t)$ , or it contains a vertex of degree more than  $f_{10}(h, t) + 1$ . Therefore, the vertex  $u$  of [line 3](#) always exists in the case where no vertex of high degree is found in [line 1](#). The correctness of this algorithm follows from [Lemma 6.11](#) and [Lemma 6.12](#). The DFS done on [line 1](#) takes time  $O(n(T))$  and the rest of the algorithm takes time  $O_{h,t}(1)$  according to the aforementioned lemmas.  $\square$

## 6.4 Approximation meets the Erdős–Pósa property

We show in this section an application of the results of [Section 6.3](#). Given a graph  $H$ , we denote by  $\text{ex}(H)$  the set of all graphs that contain  $H$  as a minor, and, subject to this condition, are minimal for the topological minor relation. Whereas [Lemma 6.2](#) is stated in terms of subdivisions, we translate to the setting of majors using the following remark.

*Remark 6.13.* Recall that  $\mathcal{M}(H)$  is defined as the set of subgraph-minimal graphs containing  $H$  as a minor. For every graph  $H$ , the following holds:

$$\mathcal{M}(H) = \mathcal{T}(\text{ex}(H)).$$

Also, the size of  $\text{ex}(H)$  is upper-bounded by some function of  $m(H)$ .

Applied to [Lemma 6.2](#), the above remark yields the next corollary.

**Corollary 6.14.** *There is an algorithm that, given a positive integer  $t$ , a finite collection  $\mathcal{H}$  of connected graphs where  $h = m(\mathcal{H})$ , and a  $t$ -partitioned protrusion  $\mathbf{P} = (\mathbf{G}, (T, s, \mathcal{X}))$  of a graph  $W$  with  $n(\mathbf{G}) > f_6(h, t)$ , outputs either*

- an  $\mathcal{H}$ -major of  $W$  with at most  $f_6(h, t)$  edges, or
- a graph  $W'$  such that

$$\begin{aligned} \text{x-pack}_{\mathcal{M}(\mathcal{H})}(W') &= \text{x-pack}_{\mathcal{M}(\mathcal{H})}(W), \\ \text{x-cover}_{\mathcal{M}(\mathcal{H})}(W') &= \text{x-cover}_{\mathcal{M}(\mathcal{H})}(W), \text{ and} \\ n(W') &< n(W). \end{aligned}$$

Furthermore, this algorithm runs in time  $O_{t,h}(n(T))$ .

Let us restate the main result of [Section 4.1.3](#), that we will use in the subsequent proofs.

**Theorem 4.4.** *There exists an algorithm that, with input three positive integers  $r, w, z$  and an  $n$ -vertex graph  $G$ , outputs one of the following:*

- a  $\theta_r$ -major in  $G$  with at most  $z$  edges,
- a connected  $(2r - 2)$ -edge-protrusion  $Y$  of  $G$  with extension more than  $w$ , or
- an  $H$ -major in  $G$  for some graph  $H$  where  $\delta(H) \geq \frac{1}{r-1} 2^{\frac{z-5r}{4r(2w+1)}}$ ,

in  $O_r(m)$  steps.

## 6.4.1 Reduce or progress

The proof of the next lemma combines [Theorem 4.4](#) and [Corollary 6.14](#).

**Lemma 6.15** (reduce or progress). *There is an algorithm that, with input  $x \in \{v, e\}$ ,  $r \in \mathbb{N}_{\geq 2}$ ,  $k \in \mathbb{N}$  and an  $n$ -vertex graph  $W$ , outputs one of the following:*

- a  $\theta_r$ -major of  $W$  with at most  $O_r(\log k)$  edges;
- a graph  $W'$  where

$$\begin{aligned} x\text{-cover}_{\mathcal{M}(\theta_r)}(W') &= x\text{-cover}_{\mathcal{M}(\theta_r)}(W), \\ x\text{-pack}_{\mathcal{M}(\theta_r)}(W') &= x\text{-pack}_{\mathcal{M}(\theta_r)}(W), \text{ and} \\ n(W') &< n(W); \text{ or} \end{aligned}$$

- an  $H$ -major in  $W$ , for some graph  $H$  with  $\delta(H) \geq k(r+1)$ ,

in  $O_r(m)$  steps.

*Proof.* We set  $t = 2r - 2$ ,  $w = f_6(h, t)$ ,  $z = 2r(w - 1) \log(k(r+1)(r-1)) + 5r$ , and  $h = m(\mathcal{M}(\theta_r))$ . Observe that  $z = O_r(\log k)$  and  $h, t, w = O_r(1)$ . Also observe that our choice for variable  $z$  ensures that  $2^{\frac{z-5r}{2r(w-1)}} / (r-1) = k(r+1)$ .

By applying the algorithm of [Theorem 4.4](#) to  $r, w, z$ , and  $W$ , we obtain in  $O_r(m(W))$ -time either:

**First case:** a  $\theta_r$ -major in  $W$  of at most  $z$  edges,

**Second case:** a  $(2r - 2)$ -edge-protrusion  $Y$  of  $W$  with extension  $> w$ , or

**Third case:** an  $H$ -major  $M$  in  $W$ , for some graph  $H$  with  $\delta(H) \geq k(r+1)$ .

In the first case, we return the obtained  $\theta_r$ -major.

In the second case, by applying the algorithm of [Corollary 6.14](#) on  $Y$ , we get in  $O(n(W))$ -time either a  $\theta_r$ -major of  $W$  on at most  $w = O_r(1)$  vertices, or a graph  $W'$  where, for  $x \in \{v, e\}$ ,  $x\text{-cover}_{\mathcal{H}}(W') = x\text{-cover}_{\mathcal{H}}(W)$ ,  $x\text{-pack}_{\mathcal{H}}(W') = x\text{-pack}_{\mathcal{H}}(W)$  and  $n(W') < n(W)$ .

In the third case, we return the major  $M$ .

In each of the above cases, we get after  $O(m)$  steps either a major of a graph with large minimum degree, or a small  $\theta_r$ -major in  $W$ , or an equivalent graph that has less vertices.  $\square$

It might not be clear yet to what purpose the major of a graph of degree more than  $k(r+1)$  output by the algorithm of [Lemma 6.15](#) can be used. Recall that we presented in [Subsection 4.2.2](#) an algorithm that finds a large packing of  $\theta_r$ -majors in a graph of large minimum degree. Applying this algorithm to the graph output by [Lemma 6.15](#) gives the desired packing.

## 6.4.2 Approximation algorithms

**Theorem 5.24** is a direct combinatorial consequence of the following.

**Theorem 6.16.** *There is a function  $f_{12}: \mathbb{N} \rightarrow \mathbb{N}$  and an algorithm that, with input  $x \in \{v, e\}$ ,  $r \in \mathbb{N}_{\geq 2}$ ,  $k \in \mathbb{N}_{\geq 1}$ , and an  $n$ -vertex graph  $W$ , outputs either a  $x$ - $\mathcal{M}(\theta_r)$ -packing of  $W$  of size  $k$  or an  $x$ - $\mathcal{M}(\theta_r)$ -covering of  $W$  of size at most  $f_{12}(r) \cdot k \cdot \log k$ . Moreover, this algorithm runs in  $O(n \cdot m)$  steps if  $x = e$  and in  $O(n^c + n \cdot m)$  steps if  $x = v$ , where  $c$  is the constant from **Theorem 4.19**.*

*Proof.* Let  $f_{12}: \mathbb{N} \rightarrow \mathbb{N}$  be a function such that each  $\theta_r$ -major output by the algorithm of **Lemma 6.15** has size at most  $f_{12}(r) \cdot \log k$ . We consider the following procedure:

1.  $G := W$ ;  $P := \emptyset$ ;
2. apply the algorithm of **Lemma 6.15** on  $(x, r, k, G)$ :
  - Progress:** if the output is a  $\theta_r$ -major  $M$ , let  $G := G \setminus A_x(M)$  and  $P = P \cup \{M\}$ ;
  - Win:** if the output is a  $H$ -major  $M$  in  $W$  for some graph  $H$  with  $\delta(H) \geq k(r+1)$ , apply the algorithm of **Theorem 4.21** (if  $x = e$ ) or the one of **Theorem 4.23** (if  $x = v$ ) to  $H$  to obtain an  $x$ - $\mathcal{M}(\theta_r)$ -packing of size  $k$  in  $H$ ; using  $M$ , translate this packing into an  $x$ - $\mathcal{M}(\theta_r)$ -packing of size  $k$  in  $W$  and return this new packing;
  - Reduce:** otherwise, the output is a graph  $G'$ : let  $G := G'$ ;
3. if  $|P| = k$  then return  $P$  which is an  $x$ - $\mathcal{M}(\theta_r)$ -packing of size  $k$  in  $W$ ;
4. if  $n(W) = 0$  then return  $P$  which is in this case a  $x$ - $\mathcal{M}(\theta_r)$ -covering of size at most  $f_{12}(r) \log k$  of  $W$ ;
5. Otherwise, go back to **line 2**.

This algorithm clearly returns the desired result. Furthermore, the loop is executed at most  $n(W)$  times and each call to the algorithm of **Lemma 6.15** takes  $O(m(W))$  steps. When the algorithm reaches the “Win” case (which can happen at most once), the calls to the algorithm of **Theorem 4.21** (if  $x = e$ ) or the one of **Theorem 4.23** (if  $x = v$ ), respectively, take  $O(m(H))$  and  $O((n(H))^c)$  steps. Therefore, in total, this algorithm terminates in  $O(n \cdot m)$  steps if  $x = e$  and in  $O(n^c + n \cdot m)$  steps if  $x = v$ .  $\square$

Observe that if the algorithm of **Theorem 6.16** reaches the “Win” case, then the input graph is known to contain an  $x$ - $\mathcal{M}(\theta_r)$ -packing of size at least  $k$ . As a consequence, if we are only interested in the existence of a packing or covering, the call to the algorithm of **Theorem 4.21** or **Theorem 4.23** is not necessary. This gives a faster algorithm for the existential version of **Theorem 6.16**.

**Corollary 6.17.** *There is an algorithm that, with input  $x \in \{v, e\}$ ,  $r \in \mathbb{N}_{\geq 2}$ ,  $k \in \mathbb{N}$ , and a graph  $W$ , outputs 0 only if  $W$  has an  $x$ - $\mathcal{M}(\theta_r)$ -packing of size  $k$  or 1 only if  $W$  has an  $x$ - $\mathcal{M}(\theta_r)$ -covering of size at most  $f_{12}(r) \cdot k \cdot \log k$ . Furthermore this algorithm runs in  $O(n \cdot m)$  steps.*

Notice that there may be graphs where both the outputs 0 and 1 of the algorithm of [Corollary 6.17](#) are valid. We now conclude this section with the proof of [Theorem 6.1](#). We start by restating it.

**Theorem 6.1.** *For every  $r \in \mathbb{N}_{\geq 2}$  and every  $x \in \{v, e\}$ , there exists an  $O(n \cdot \log(n) \cdot m)$ -step algorithm that, given a graph  $G$ , outputs an  $O(\log \text{OPT})$ -approximation for  $x\text{-cover}_{\mathcal{M}(\theta_r)}$  and  $x\text{-pack}_{\mathcal{M}(\theta_r)}$ .*

*Proof.* Let us call  $A$  the algorithm of [Corollary 6.17](#). Let  $k_0 \in \llbracket 1, n(W) \rrbracket$  be an integer such that  $A(x, r, k_0, W) = 1$  and  $A(x, r, k_0 - 1, W) = 0$ , and let us show that the values  $k_0 - 1$  and  $k_0 \log k_0$  are  $O(\log \text{OPT})$ -approximations of  $x\text{-pack}_{\mathcal{M}(\theta_r)}(W)$  and  $x\text{-cover}_{\mathcal{M}(\theta_r)}(W)$ , respectively.

First, notice that for every  $k > x\text{-pack}_{\mathcal{M}(\theta_r)}(W)$ , the value returned by  $A(x, r, k, W)$  is 1. Symmetrically, for every  $k$  such that  $k \log k < x\text{-cover}_{\mathcal{M}(\theta_r)}(W)$ , the value of  $A(x, r, k, W)$  is 0. Therefore, the value  $k_0$  is such that:

$$k_0 - 1 \leq x\text{-pack}_{\mathcal{M}(\theta_r)}(W) \text{ and} \\ x\text{-cover}_{\mathcal{M}(\theta_r)}(W) \leq k_0 \log k_0.$$

As every minimal covering must contain at least one vertex or edge (depending on whether  $x = v$  or  $x = e$ ) of each model of a maximal packing,  $x\text{-pack}_{\mathcal{M}(\theta_r)}(W) \leq x\text{-cover}_{\mathcal{M}(\theta_r)}(W)$ , hence we have:

$$k_0 - 1 \leq x\text{-pack}_{\mathcal{M}(\theta_r)}(W) \leq x\text{-cover}_{\mathcal{M}(\theta_r)}(W) \leq k_0 \log k_0. \quad (6.1)$$

We then deduce:

$$1 \leq \frac{x\text{-pack}_{\mathcal{M}(\theta_r)}(W)}{k_0 - 1} \leq \frac{k_0 \log k_0}{k_0 - 1} \\ = O(\log k_0) \\ = O(\log x\text{-pack}_{\mathcal{M}(\theta_r)}(W)) \quad \text{using (6.1),}$$

and symmetrically for the covering parameter:

$$1 \leq \frac{k_0 \log k_0}{x\text{-cover}_{\mathcal{M}(\theta_r)}(W)} \leq \frac{k_0 \log k_0}{k_0 - 1} \\ = O(\log k_0) \\ = O(\log x\text{-cover}_{\mathcal{M}(\theta_r)}(W)) \quad \text{using (6.1) again.}$$

This proves that the values  $k_0 - 1$  and  $k_0 \log k_0$  are indeed  $O(\log \text{OPT})$ -approximations of  $x\text{-pack}_{\mathcal{M}(\theta_r)}(W)$  and  $x\text{-cover}_{\mathcal{M}(\theta_r)}(W)$ , respectively. The value  $k_0$  can be found by performing a binary search in the interval  $\llbracket 1, n \rrbracket$ , with  $O(\log n)$  calls to Algorithm A. Hence, our approximation algorithm runs in  $O(n \cdot \log(n) \cdot m)$  steps.  $\square$

Notice that all our results are strongly exploiting [Lemma 6.2](#) that holds for every finite collection  $\mathcal{H}$  of connected graphs. Actually, what is missing in order to have an overall generalization of all of our results, is an extension of [Theorem 4.4](#) where  $\mathcal{M}(\theta_r)$  is replaced by any finite collection  $\mathcal{H}$  of connected planar graphs. This is an interesting combinatorial problem even for particular instantiations of  $\mathcal{H}$ .

# CHAPTER 7

---

## CONCLUSION

### 7.1 General contribution

This thesis is centered around results on graph classes excluding a given pattern as a substructure. We focused on theorems providing, for the considered classes, either a decomposition theorem, or a bound on a graph invariant. These results are then applied to problems of well-quasi-ordering and Erdős–Pósa-type problems.

**Decomposition theorems and applications to well-quasi-ordering.** Decomposition theorems sometimes provide enough information about the considered class to obtain well-quasi-ordering results. We proved decomposition theorems for the classes of simple graphs excluding the graphs  and  as induced minors, and  as a contraction. These structural descriptions allowed us to show that the considered classes are well-quasi-ordered by the induced minors and contraction relations, respectively. The study of some infinite antichains for these relations then yielded, in each case, a characterization on the well-quasi-ordered classes defined by one forbidden pattern. Such dichotomies were known for most of the other containment relations that are not well-quasi-orders in general (subgraph, induced subgraph and topological minor). Using structural results of Tutte and Oporowski et al., we obtained a similar characterization of well-quasi-ordered classes of multigraphs with respect to the contraction relation. Considering infinite antichains rather than well-quasi-orders, we characterized all canonical antichains for the induced minor relation and proved that there is no such antichain for the contraction relation.

**Bounds on graph invariants.** A chapter of this thesis is devoted to theorems providing bounds on invariants of the graphs excluding some pattern as a substructure. We considered basic invariants like minimum and maximum degree, structural invariants like treewidth and tree-cut width, and introduced the  $H$ -girth, a family of invariants that generalizes the parameter of girth. The excluded patterns range from parameterized planar graphs (wheels, double wheels, yurts, packings of  $\theta_r$ 's) to larger families of graphs as planar subcubic graphs, cliques, and graphs of pathwidth

at most 2. A summary of the results we proved is given in [Table 7.1](#). For each line, the table can be read as follows: if we consider the class of graphs excluding the pattern of the first column (an example of which is given in the second column) with respect to the relation of the third column, then the parameter given in the fourth column is equal to the expression of the fifth column.

excluded pattern	ex.	relation	par.	value of the parameter
wheel of order $k$		minor	tw	$\Theta(k)$
double wheel of order $k$		minor	tw	$O(k^2 \log^2 k)$
$H$ , $\text{pw}(H) \leq 2$		minor	tw	$O(( H  + \ H\ )^2)$
yurt graph of order $k$		minor	tw	$O(k^4)$
$k \cdot \theta_r$		minor	tw $\delta$	$\Theta(k \log k)$ $\Theta(k)$
edge-disj. union of $k \theta_r$ 's		minor	$\Delta$	$\Theta(k)$
$K_k$		minor	$\theta_r$ -girth	$O(\log k)$
$H$ planar subcubic		immersion	tcw	$O(\ H\ ^{29} \text{polylog } \ H\ )$

Table 7.1: Summary of the exclusion results of [Chapter 4](#). When applicable,  $r$  is considered to be constant. The bounds related to  $K_k$  and to edge-disjoint  $\theta_r$ 's have other requirements that are given in the corresponding theorems. The result on planar subcubic graphs draws upon [Theorem 4.67](#) as obtained by Wollan.

**The Erdős-Pósa property.** A possible application of our results on graph invariants is Erdős-Pósa-type problems. For this, we presented two general schemes. The first one, [Theorem 5.3](#), is a master theorem that unifies known techniques based on tree decompositions and treewidth. We applied it to parameters that are less common for this type of problems, like minimum and maximum degree, and to structural parameters alternative to the treewidth, the tree-partition width and the tree-cut width. This technique, together with exclusion results proved in the previous section, yielded new Erdős-Pósa-type results for various classes of graphs. The second scheme is related to the notion of girth and its aforementioned extensions ( $H$ -girth). We used it to prove that majors of  $\theta_r$  have the edge and vertex Erdős-Pósa property with optimal gap  $k \mapsto c \cdot k \log k$  (for some  $c$  depending on  $r$ ). This provides one of the first results on the edge-Erdős-Pósa property of majors. Last, as an algorithmic application of Erdős-Pósa-type results, we give a  $O(\log \text{OPT})$ -approximation algorithm for the (edge and vertex) packing and covering numbers of  $\mathcal{M}(\theta_r)$ , improving the known  $O(\log n)$ -approximation for the vertex version and providing the first approximation for the edge version (for general  $r$ ). The proof relies on the concept of *edge-protrusion*, a variation of the notion of protrusion that is tailored for edge-related problems.

## 7.2 Perspectives and problems

### 7.2.1 On the $H$ -girth

In this thesis we introduced the  $H$ -girth and considered in particular the  $\theta_r$ -girth (for any positive integer  $r$ ). Recall that for every graph  $H$ , we call  $H$ -girth of a graph the minimum order of an  $H$ -major in this graph. It could be fruitful to study this parameter for different instantiations of  $H$ . As we mention in the corresponding section, the girth is a concept that arises in various areas of Graph Theory and an interesting question is to investigate for which graphs  $H$  the girth can be replaced by  $H$ -girth in these contexts. To give an example related to the topics addressed in this thesis, let us consider the following result of Thomassen.

**Theorem 7.1** ([Tho83b]). *There is a function  $f: \mathbb{N} \rightarrow \mathbb{N}$  such that for every  $r \in \mathbb{N}$ , every graph of minimum degree 3 and girth at least  $f(r)$  contains a  $K_r$ -major.*

The corresponding problem for  $H$ -girth is the following.

*Problem 7.2.* Given a graph  $H$ , does **Theorem 7.1** hold when “girth” is replaced with “ $H$ -girth”?

According to the original theorem,  $K_3$  is a positive instance of **Problem 7.2**. Also, we gave in **Section 4.1** a proof that  $\theta_r$  would be an positive instance too (for every  $r \in \mathbb{N}$ ) if we allowed minor changes of the statement (in the main result of **Section 4.1**, we require the minimum degree to be at least  $3r$ ).

A first step could be to try to extend **Theorem 7.1** to  $K_4$ -girth.

*Question 7.3.* Does **Theorem 7.1** hold when “girth” is replaced with “ $K_4$ -girth”?

It turns out that a positive instances to **Problem 7.2**, even for specific patterns and with slight changes in the statement, may be used to prove Erdős–Pósa-type results. Indeed, we detailed in the section on the Erdős–Pósa property a general approach to prove Erdős–Pósa-type results based on the  $H$ -girth, and we unfolded it to the  $\theta_r$ -girth to obtain our results. In order to obtain results for any other graph  $H$ , the only missing part is a theorem expressing that a graph of large  $H$ -girth either contains a large packing of  $H$ -majors, or can be reduced to a smaller graph with the same parameters of packing and covering. A positive answer to **Problem 7.2** for some graph  $H$  would guarantee that we can prove such a theorem. In the case where the answer to **Problem 7.2** is negative for  $H$ , we can still focus on the weaker goal of identifying positive instances to the following problem.

*Problem 7.4.* Given a graph  $H$ , does **Theorem 7.1** hold when “girth” is replaced with “ $H$ -girth”, if we additionally assume that  $G$  does not contain  $a$ -edge-protrusions with extension more than  $b$ , for some constants  $a, b \in \mathbb{N}$  depending on  $H$ ?

There are other parameters that can be defined in a similar way: given a graph  $G$ , we could consider the minimum order of a subgraph of  $G$  that contains  $H$  as an induced minor (respectively topological minor, immersion, etc.). For each of these variants the above questions can be asked and lead to interesting developments.

## 7.2.2 On the Erdős–Pósa property

The vertex-Erdős–Pósa property of graph majors is completely explained (i.e. we know precisely the graphs whose majors have the vertex-Erdős–Pósa property), mainly thanks to the work of Robertson and Seymour. However, its edge-counterpart is still not well understood.

*Question 7.5.* Characterize graphs whose majors have the edge-Erdős–Pósa property?

We proved in this thesis that the majors of any graph in  $\{\theta_r\}_{r \in \mathbb{N}} \cup \{\theta_{r,r'}\}_{r,r' \in \mathbb{N}}$  have the edge-Erdős–Pósa property (recall that  $\theta_{r,r'}$  can be obtained from  $P_3$  by setting the multiplicities of the edges to  $r$  and  $r'$ , respectively). We also provided a construction that shows that the majors of any non-planar graph fail to have the edge-Erdős–Pósa property in general (i.e. for any host graph). What is the correct boundary between these two classes? Again, a first step is to look at small graphs.

*Question 7.6.* Does  $\mathcal{M}(K_4)$  have the edge-Erdős–Pósa property?

Naturally, similar questions can be asked for other classes defined via a containment relation. Lets us now consider immersions. We do not understand yet the classes of graphs whose immersion-expansions have the (edge or vertex) Erdős–Pósa property.

*Question 7.7.* Characterize graphs whose immersion expansions have the vertex-Erdős–Pósa property (respectively, edge-Erdős–Pósa property)?

In this thesis we proved that the immersion expansions of every planar subcubic graph have the vertex and edge-Erdős–Pósa property. We also gave examples of families of planar graphs with maximum degree at least 4, the immersion expansions of which do not have the vertex and edge Erdős–Pósa property. Furthermore, as for majors, the immersion expansions of a non-planar graph do not have the Erdős–Pósa property in general. Again, we provide weaker questions on this topic.

*Question 7.8.* Does  $\mathcal{I}(W_4)$  have the vertex-Erdős–Pósa property (resp. edge-Erdős–Pósa property), where  $W_4 = \boxtimes$ ?

*Question 7.9.* For every  $r \in \mathbb{N}$ , does  $\mathcal{I}(\theta_r)$  have the vertex-Erdős–Pósa property (resp. edge-Erdős–Pósa property)?

Symmetrically to minors, the status of the Erdős–Pósa property for topological minors is half-solved: the class of graphs whose subdivisions have the vertex-Erdős–Pósa property have been completely characterized in [LPW14]. On the other hand, not much is known about the graphs whose subdivisions have the edge-Erdős–Pósa property. Families of trees with maximum degree 4 whose subdivisions do not have the (vertex and edge) Erdős–Pósa property have been provided in [Tho88], and we gave in [Subsection 5.6.1](#) other examples of such families.

*Question 7.10.* Characterize graphs whose subdivisions have the edge-Erdős–Pósa property?

### 7.2.3 On edge-protrusions

The concept of protrusion has been introduced [BFL<sup>+</sup>09b, BFL<sup>+</sup>09a] in order to easily deal with pieces of a graph that are *simple* (for the considered problem) and have a small *boundary* with the rest of the graph. The boundary is usually a set of vertices and this has certain drawbacks when dealing with edge-deletion problems. In particular, knowing that the protrusion has a small (vertex-)boundary with the rest of the graph does not give any bound on the number of edges one would need to delete in order to separate the protrusion from the rest of the graph. In some of the problems we considered (covering the vertices or edges of the  $\theta_r$ -majors of a graph), we fixed this problem by introducing an edge-counterpart of the notion of protrusion. This notion of edge-protrusion can be easily used with tree-like decompositions that provide small edge-separators, such as tree-cut decomposition and tree-partition. We hope that it will find further combinatorial and algorithmic applications to edge-related problems.

### 7.2.4 On well-quasi-ordering

As we characterized all classes of graphs defined by one forbidden induced minor which are well-quasi-ordered by this relation, a natural line of research is to investigate what happens when we exclude more graphs.

*Question 7.11.* For which graphs  $G$  and  $H$  is the class  $\text{Excl}_{\leq_{i.m.}}(G, H)$  well-quasi-ordered by induced minors?

Of course, the question can be asked for any number of forbidden induced minors. It can even be asked when infinite families of graphs are forbidden. Let us call *fan* a graph containing a vertex whose removal yields a path.

*Question 7.12.* Let  $\mathcal{H}$  be a class of planar graphs that do not contain large fans as induced minors. Is  $(\mathcal{H}, \leq_{i.m.})$  a well-quasi-order?

Several well-quasi-ordering proofs presented in this thesis handle graphs with vertices labelled by the elements of a well-quasi-order. In fact, labels are a standard tool in well-quasi-ordering theory. An other question of interest is to characterize the classes of graphs that are well-quasi-ordered by induced minors when labelled by a well-quasi-order. Note that this question has already been considered for the induced subgraph relation in [AL14].

*Question 7.13.* Characterize the classes of graphs  $\mathcal{H}$  such that, for every well-quasi-order  $(\mathcal{L}, \preceq)$ , the class  $(\text{lab}_{(\mathcal{L}, \preceq)}(\mathcal{H}), \leq_{i.m.})$  is a well-quasi-order.

An other question related to labels has been asked by Pouzet. We say that a class  $\mathcal{G}$  of graphs is *n-well-quasi-ordered* (for some  $n \in \mathbb{N}$ ) if  $(\text{lab}_{([1, n], =)}(\mathcal{G}), \leq_{i.sg.})$  is a well-quasi-order. In other words, the graphs of  $\mathcal{G}$  labeled with integers of  $[1, n]$  are well-quasi-ordered by the induced subgraph relation that respects labels.

*Conjecture 7.14* (Pouzet's conjecture [Pou72]). A class of graphs that is (downwards) closed under induced subgraphs is 2-well-quasi-ordered iff it is *n-well-quasi-ordered*, for every  $n \in \mathbb{N}$ .

An obstruction to well-quasi-ordering is the existence of infinite antichains, which are objects of interest as well. To conclude this section, let us mention a question asked by Ding.

*Question 7.15* ([Din]). Characterize infinite antichains of the class of all fans, for the induced subgraph relation.

Observe that there is a strong connection between the quasi-order of fans ordered by the induced subgraph relation and that of binary words ordered by the factor relation. Every word corresponds to a fan, and every fan to at most two words. Therefore the goal of [Question 7.15](#) is essentially the same as characterizing infinite antichains of  $\{0, 1\}^*$  for the factor relation.

## 7.3 Publications

The results presented in this thesis are mainly originated from nine publications. However, for the sake of consistency, we chose to omit some results obtained during the PhD. Let us briefly present here all of them.

### 7.3.1 On the Erdős–Pósa property

- [[RT13a](#), [RT13b](#)] *Polynomial gap extensions of the Erdős–Pósa Theorem* with Dimitrios M. Thilikos, EuroComb 2013. We prove that the majors of graphs of pathwidth at most two have the vertex-Erdős–Pósa property with gap  $k \mapsto 2^{O(h^2)}k^2 \log k$ . We omitted these results as they are now subsumed by the results presented in [Subsection 5.4.1](#).
- [[RST16](#)] *An edge variant of the Erdős–Pósa property* with Ignasi Sau and Dimitrios M. Thilikos, Discrete Mathematics (2016). These results are presented in [Chapter 5](#), more specifically in [Subsection 5.4.2](#).
- [[CRST15b](#), [CRST15a](#)] *An  $O(\log \text{OPT})$ -approximation for covering/packing minor models of  $\theta_r$* , with Dimitris Chatzidimitriou, Ignasi Sau and Dimitrios M. Thilikos, WAOA 2015. Long version submitted. These results are presented in [Chapter 6](#).
- [[GKRT16](#)] *Packing and covering immersion models of planar subcubic graphs*, with Archontia Giannopoulou, O-jong Kwon, and Dimitrios M. Thilikos, WG 2016. Long version submitted. These results are presented in [Section 4.4](#) and [Subsection 5.4.4](#).
- [[RT16](#)] *Recent techniques and results on the Erdős–Pósa property*, with Dimitrios M. Thilikos. Submitted. These results are presented in [Chapter 5](#).
- [[Ray16](#)] *Hitting minors, subdivisions, and immersions in tournaments*. Submitted. This paper deals with the Erdős–Pósa property in directed graphs. Using exclusion theorems of Chudnovsky, Fradkin, Kim, and Seymour in the class of

tournaments, we show that the majors (respectively subdivisions) of every directed graph have the vertex-Erdős–Pósa property in tournaments, and that the immersion expansions of every strongly connected directed graph have the edge-Erdős–Pósa property in tournaments. We chose to omit this result because it is about directed graphs, whereas all the other results presented in the thesis deal with undirected graphs.

### 7.3.2 On well-quasi-ordering

- [BKRT15a, BKRT15b] *Induced minors and well-quasi-ordering* with Jarosław Błasiok, Marcin Kamiński and Théophile Trunck, EuroComb 2015. Long version submitted. These results are presented in [Chapter 3](#).
- [KRT14] *Multigraphs without big bonds are wqo by contraction* with Marcin Kamiński and Théophile Trunck, submitted. These results are presented in [Section 3.8](#).
- [KRT16] *Well-quasi-ordering  $H$ -contraction-free graphs*, with Marcin Kamiński and Théophile Trunck. Submitted. These results are presented in [Section 3.7](#).

### 7.3.3 On exclusion results

- [RT15] *Low polynomial exclusion of planar graph patterns* with Dimitrios M. Thilikos, to appear in *Journal of Graph Theory* (2016). These results are presented in [Section 4.3](#).
- [CRST15c] *Minors in graphs of large  $\theta_r$ -girth* with Dimitris Chatzidimitriou, Ignasi Sau and Dimitrios M. Thilikos, submitted. These results are presented in [Section 4.1](#).

### 7.3.4 On other topics

- [AKR14] *Scattered packings of cycles* with Aistis Atminas and Marcin Kamiński, to appear in *Theoretical Computer Science* (2016). In this paper, we consider the problem of deciding, given a graph  $G$  and two positive integers  $k$  and  $l$ , whether  $G$  contains  $k$  cycles that are at distance at least  $l$  from each other. We obtain polynomial kernels and hardness results for this problem parameterized by several combinations of  $k$ ,  $l$ , and the maximum degree of the input graph.
- [GPR<sup>+</sup>16a] *Cutwidth: obstructions and algorithmic aspects* with Archontia Giannopoulou, Michał Pilipczuk, Dimitrios M. Thilikos, and Marcin Wrochna, IPEC 2016. Cutwidth is a graph parameter which measures the minimal number of edges of a graph crossing an ordering of its vertices. Our main result is a single-exponential upper bound on the size of immersion-minimal graphs that have cutwidth greater than  $k$  (i.e. obstructions for cutwidth at most  $k$ ).

- [GPR<sup>+</sup>16b] *Linear kernels for edge deletion problems to immersion-closed graph classes* with Archontia Giannopoulou, Michał Pilipczuk, Dimitrios M. Thilikos, and Marcin Wrochna. Submitted. Let  $\mathcal{F}$  be a collection of connected graphs containing at least one planar graph. In [FLMS12], Fomin et al. considered the problem of deciding, with input a graph  $G$  and a positive integer  $k$ , if one can delete  $k$  vertices in  $G$  in order to obtain an  $\mathcal{F}$ -minor free graph. They obtained a randomized constant factor approximation for the optimization version of this problem (i.e. computing the minimum  $k$  such that the above holds) and a randomized FPT algorithm for the parameterized version. We consider the immersion variant of this problem where one is asked if we can delete  $k$  edges in the input graph in order to get an  $\mathcal{F}$ -immersion free graph. In this direction, when  $\mathcal{F}$  contains at least one planar subcubic graph, we obtain a constant factor approximation, a linear kernel, and an FPT algorithm.

---

## BIBLIOGRAPHY

- [ACR03] Alessandro Panconesi Alberto Caprara and Romeo Rizzi. Packing cycles in undirected graphs. *Journal of Algorithms*, 48(1):239 – 256, 2003.
- [ADG04] Julio Aracena, Jacques Demongeot, and Eric Goles. Positive and negative circuits in discrete neural networks. *Neural Networks, IEEE Transactions on*, 15(1):77–83, Jan 2004.
- [AKKW16] Saeed Akhoondian Amiri, Ken-Ichi Kawarabayashi, Stephan Kreutzer, and Paul Wollan. The Erdős–Pósa property for directed graphs. *ArXiv e-prints 0904.0727*, 2016.
- [AKR14] Aistis Atminas, Marcin Kamiński, and Jean-Florent Raymond. Scattered packings of cycles. *ArXiv e-prints*, September 2014.
- [AL14] Aistis Atminas and Vadim V. Lozin. Labelled induced subgraphs and well-quasi-ordering. *Order*, pages 1–16, 2014.
- [ALBT11] S. Aparna Lakshmanan, Cs. Bujtás, and Zs. Tuza. Small edge sets meeting all triangles of a graph. *Graphs and Combinatorics*, 28(3):381–392, 2011.
- [Alo98] Noga Alon. Piercing  $d$ -intervals. *Discrete & Computational Geometry*, 19(3):333–334, 1998.
- [Alo02] Noga Alon. Covering a hypergraph of subgraphs. *Discrete Mathematics*, 257(2–3):249 – 254, 2002.
- [Ara08] Julio Aracena. Maximum number of fixed points in regulatory boolean networks. *Bulletin of Mathematical Biology*, 70(5):1398–1409, 2008.
- [ARS16] Julio Aracena, Adrien Richard, and Lilian Salinas. Number of fixed points and disjoint cycles in monotone Boolean networks. *ArXiv e-prints*, February 2016.

- [BBR07a] E. Birmelé, J. Bondy, and B. Reed. Brambles, prisms and grids. In Adrian Bondy, Jean Fonlupt, Jean-Luc Fouquet, Jean-Claude Fournier, and Jorge L. Ramírez Alfonsín, editors, *Graph Theory in Paris*, Trends in Mathematics, pages 37–44. Birkhäuser Basel, 2007.
- [BBR07b] Etienne Birmelé, J. Adrian Bondy, and Bruce A. Reed. The Erdős-Pósa property for long circuits. *Combinatorica*, 27(2):135–145, 2007.
- [BEM<sup>+</sup>04] S. Bezrukov, R. Elsässer, B. Monien, R. Preis, and J.-P. Tillich. New spectral lower bounds on the bisection width of graphs. *Theoretical Computer Science*, 320(2–3):155 – 174, 2004.
- [BFL<sup>+</sup>09a] Hans L. Bodlaender, Fedor V. Fomin, Daniel Lokshtanov, Eelko Penninkx, Saket Saurabh, and Dimitrios M. Thilikos. (Meta) Kernelization. In *Proceedings of FOCS 2009*, pages 629–638. IEEE Computer Society, 2009.
- [BFL<sup>+</sup>09b] Hans L. Bodlaender, Fedor V. Fomin, Daniel Lokshtanov, Eelko Penninkx, Saket Saurabh, and Dimitrios M. Thilikos. (meta) kernelization. *ArXiv e-prints 0904.0727*, 2009.
- [BHK62] Joseph Battle, Frank Harary, and Yukihiro Kodama. Additivity of the genus of a graph. *Bulletin of the American Mathematical Society*, 68(6):565–568, 11 1962.
- [BJS14] Henning Bruhn, Felix Joos, and Oliver Schaudt. Long cycles through prescribed vertices have the Erdős-Pósa property. *ArXiv e-prints*, December 2014.
- [BKRT15a] Jarosław Błasiok, Marcin Kamiński, Jean-Florent Raymond, and Théophile Trunck. Induced minors and well-quasi-ordering. *arXiv preprint arXiv:1510.07135*, October 2015.
- [BKRT15b] Jarosław Błasiok, Marcin Kamiński, Jean-Florent Raymond, and Théophile Trunck. Induced minors and well-quasi-ordering. *Electronic Notes in Discrete Mathematics*, 49:197 – 201, 2015. The Eight European Conference on Combinatorics, Graph Theory and Applications, EuroComb 2015.
- [Bod93] Hans L. Bodlaender. On linear time minor tests with depth-first search. *J. Algorithms*, 14(1):1–23, 1993.
- [Bou13] Nicolas Bousquet. *Hitting sets: VC-dimension and Multicut*. PhD thesis, Montpellier 2, 2013.
- [BR00] Claude Berge and Bruce Reed. Optimal packings of edge-disjoint odd cycles. *Discrete Mathematics*, 211(1–3):197 – 202, 2000.

- [BRST91] Dan Bienstock, Neil Robertson, Paul Seymour, and Robin Thomas. Quickly excluding a forest. *J. Combin. Theory Ser. B*, 52(2):274–283, 1991.
- [BT04] Hans L. Bodlaender and Dimitrios M. Thilikos. Computing small search numbers in linear time. In *Proceedings of the First International Workshop on Parameterized and Exact Computation (IWPEC 2004)*, volume 3162 of *LNCS*, pages 37–48. Springer, 2004.
- [BT15] Nicolas Bousquet and Stéphan Thomassé. Vc-dimension and Erdős–Pósa property. *Discrete Mathematics*, 338(12):2302 – 2317, 2015.
- [BTV07] Cristina Bazgan, Zsolt Tuza, and Daniel Vanderpooten. Efficient algorithms for decomposing graphs under degree constraints. *Discrete Applied Mathematics*, 155(8):979 – 988, 2007.
- [BTY11] Hans L. Bodlaender, Stéphan Thomassé, and Anders Yeo. Kernel bounds for disjoint cycles and disjoint paths. *Theoretical Computer Science*, 412(35):4570–4578, 2011.
- [Bur69] Rod M Burstall. Proving properties of programs by structural induction. *The Computer Journal*, 12(1):41–48, 1969.
- [Bur78] Michael Burstein. Kuratowski-pontrjagin theorem on planar graphs. *Journal of Combinatorial Theory, Series B*, 24(2):228 – 232, 1978.
- [BvLTT97a] Hans L. Bodlaender, Jan van Leeuwen, Richard Tan, and Dimitrios M. Thilikos. On interval routing schemes and treewidth. *Information and Computation*, 139(1):92–109, 1997.
- [BvLTT97b] Hans L. Bodlaender, Jan van Leeuwen, Richard B. Tan, and Dimitrios M. Thilikos. On interval routing schemes and treewidth. *Inf. Comput.*, 139(1):92–109, 1997.
- [CC13] Chandra Chekuri and Julia Chuzhoy. Large-treewidth graph decompositions and applications. *CoRR*, abs/1304.1577, 2013.
- [CC14] Chandra Chekuri and Julia Chuzhoy. Polynomial bounds for the grid-minor theorem. In *Proceedings of the 46th Annual ACM Symposium on Theory of Computing*, pages 60–69. ACM, 2014.
- [CFS12] Maria Chudnovsky, Alexandra Fradkin, and Paul Seymour. Tournament immersion and cutwidth. *Journal of Combinatorial Theory, Series B*, 102(1):93 – 101, 2012.
- [CGG<sup>+</sup>06] Maria Chudnovsky, Jim Geelen, Bert Gerards, Luis Goddyn, Michael Lohman, and Paul Seymour. Packing non-zero a-paths in group-labelled graphs. *Combinatorica*, 26(5):521–532, 2006.

- [Che11] Gregory Cherlin. Forbidden substructures and combinatorial dichotomies: Wqo and universality. *Discrete Math.*, 311(15):1543–1584, August 2011.
- [Chu15] Julia Chuzhoy. Excluded grid theorem: Improved and simplified. In *Proceedings of the 47th Annual ACM on Symposium on Theory of Computing (STOC)*, pages 645–654, New York, NY, USA, 2015. ACM.
- [Chu16] J. Chuzhoy. Improved Bounds for the Excluded Grid Theorem. *ArXiv e-prints*, February 2016.
- [Cou90] Bruno Courcelle. The monadic second-order logic of graphs. i. recognizable sets of finite graphs. *Information and Computation*, 85(1):12 – 75, 1990.
- [CRST15a] Dimitris Chatzidimitriou, Jean-Florent Raymond, Ignasi Sau, and Dimitrios M. Thilikos. An  $O(\log \text{OPT})$ -Approximation for Covering/Packing Minor Models of  $\theta_r$ . In Laura Sanità and Martin Skutella, editors, *Approximation and Online Algorithms: 13th International Workshop, WAOA 2015, Patras, Greece, September 17-18, 2015. Revised Selected Papers*, pages 122–132, Cham, 2015. Springer International Publishing.
- [CRST15b] Dimitris Chatzidimitriou, Jean-Florent Raymond, Ignasi Sau, and Dimitrios M. Thilikos. An  $O(\log \text{OPT})$ -approximation for covering and packing minor models of  $\theta_r$ . *ArXiv e-prints*, October 2015.
- [CRST15c] Dimitris Chatzidimitriou, Jean-Florent Raymond, Ignasi Sau, and Dimitrios M. Thilikos. Minors in graphs of large  $\theta_r$ -girth. *ArXiv e-prints*, October 2015.
- [CS05] L. Sunil Chandran and C.R. Subramanian. Girth and treewidth. *Journal of Combinatorial Theory, Series B*, 93(1):23 – 32, 2005.
- [CS11] Maria Chudnovsky and Paul Seymour. A well-quasi-order for tournaments. *Journal of Combinatorial Theory, Series B*, 101(1):47–53, 2011.
- [Dam90] Peter Damaschke. Induced subgraphs and well-quasi-ordering. *Journal of Graph Theory*, 14(4):427–435, 1990.
- [DH08] Erik D. Demaine and Mohammad Taghi Hajiaghayi. Linearity of grid minors in treewidth with applications through bidimensionality. *Combinatorica*, 28(1):19–36, 2008.
- [DHJ<sup>+</sup>13] Vida Dujmovic, Daniel J. Harvey, Gwenaël Joret, Bruce A. Reed, and David R. Wood. A linear-time algorithm for finding a complete graph minor in a dense graph. *SIAM Journal of Discrete Mathematics*, 27(4):1770–1774, 2013.

- [DHK09] Erik D. Demaine, Mohammad Taghi Hajiaghayi, and Ken-ichi Kawarabayashi. Algorithmic graph minor theory: Improved grid minor bounds and Wagner’s contraction. *Algorithmica*, 54(2):142–180, 2009.
- [Die05] Reinhard Diestel. *Graph Theory*, volume 173 of *Graduate Texts in Mathematics*. Springer-Verlag, Heidelberg, third edition, 2005.
- [Din] Guoli Ding. Webpage of Guoli Ding. <https://www.math.lsu.edu/~ding/1interests.html>.
- [Din92] Guoli Ding. Subgraphs and well-quasi-ordering. *J. Graph Theory*, 16(5):489–502, November 1992.
- [Din98] Guoli Ding. Chordal graphs, interval graphs, and wqo. *Journal of Graph Theory*, 28(2):105–114, 1998.
- [Din09] Guoli Ding. On canonical antichains. *Discrete Mathematics*, 309(5):1123 – 1134, 2009.
- [DJGT99a] Reinhard Diestel, Tommy R. Jensen, Konstantin Yu. Gorbunov, and Carsten Thomassen. Highly connected sets and the excluded grid theorem. *J. Combin. Theory Ser. B*, 75(1):61–73, 1999.
- [DJGT99b] Reinhard Diestel, Tommy R. Jensen, Konstantin Yu. Gorbunov, and Carsten Thomassen. Highly connected sets and the excluded grid theorem. *Journal of Combinatorial Theory, Series B*, 75(1):61–73, 1999.
- [DKW10] Reinhard Diestel, Ken-ichi Kawarabayashi, and Paul Wollan. The Erdős-Pósa property for clique minors in highly connected graphs. *ArXiv e-prints*, March 2010.
- [DL88] I. J. Dejter and V. Neumann Lara. Unboundedness for generalized odd cyclic transversality, 1988.
- [DO96] Guoli Ding and Bogdan Oporowski. On tree-partitions of graphs. *Discrete Mathematics*, 149(1–3):45 – 58, 1996.
- [DRT10] Jean Daligault, Michael Rao, and Stéphan Thomassé. Well-quasi-order of relabel functions. *Order*, 27(3):301–315, 2010.
- [DXZ03] Guoli Ding, Zhenzhen Xu, and Wenan Zang. Packing cycles in graphs, II. *Journal of Combinatorial Theory, Series B*, 87(2):244 – 253, 2003.
- [DZ02] Guoli Ding and Wenan Zang. Packing cycles in graphs. *Journal of Combinatorial Theory, Series B*, 86(2):381 – 407, 2002.
- [EP65] Paul Erdős and Louis Pósa. On independent circuits contained in a graph. *Canadian Journal of Mathematics*, 17:347–352, 1965.

- [ER52] P. Erdős and R. Rado. Sets having divisor property, solution to problem 4358. *Amer. Math. Monthly*, 59:255–257, 1952.
- [ES87] P. Erdős and G. Szekeres. A combinatorial problem in geometry. In Ira Gessel and Gian-Carlo Rota, editors, *Classic Papers in Combinatorics*, Modern Birkhäuser Classics, pages 49–56. Birkhäuser Boston, 1987.
- [FH14] Samuel Fiorini and Audrey Herinckx. A tighter Erdős-Pósa function for long cycles. *Journal of Graph Theory*, 77(2):111–116, 2014.
- [FHR12] Michael R. Fellows, Danny Hermelin, and Frances A. Rosamond. Well quasi orders in subclasses of bounded treewidth graphs and their algorithmic applications. *Algorithmica*, 64(1):3–18, 2012.
- [FHRV05] Samuel Fiorini, Nadia Hardy, Bruce Reed, and Adrian Vetta. Approximate min-max relations for odd cycles in planar graphs. In *Integer Programming and Combinatorial Optimization*, pages 35–50. Springer, 2005.
- [FJP10] Samuel Fiorini, Gwenaël Joret, and Ugo Pietropaoli. Hitting diamonds and growing cacti. In *Proceedings of Integer Programming and Combinatorial Optimization*, volume 6080 of *Lecture Notes in Computer Science*, pages 191–204. Springer Berlin Heidelberg, 2010.
- [FJS13] Samuel Fiorini, Gwenaël Joret, and Ignasi Sau. Optimal Erdős-Pósa property for pumpkins. Manuscript, 2013.
- [FJW13a] Samuel Fiorini, Gwenaël Joret, and David R. Wood. Excluded forest minors and the Erdős-Pósa property. *Combinatorics, Probability & Computing*, 22(5):700–721, 2013.
- [FJW13b] Samuel Fiorini, Gwenaël Joret, and David R. Wood. Excluded forest minors and the Erdős-Pósa property. *Combinatorics, Probability & Computing*, 22(5):700–721, 2013.
- [FKO09] Nikolaos Fountoulakis, Daniela Kühn, and Deryk Osthus. Minors in random regular graphs. *Random Structures & Algorithms*, 35(4):444–463, 2009.
- [FL94] Michael R. Fellows and Michael A. Langston. On search, decision, and the efficiency of polynomial-time algorithms. *J. Comput. System Sci.*, 49(3):769–779, 1994.
- [FLM<sup>+</sup>13] Fedor V. Fomin, Daniel Lokshtanov, Neeldhara Misra, Geevarghese Philip, and Saket Saurabh. Quadratic upper bounds on the Erdős-Pósa property for a generalization of packing and covering cycles. *Journal of Graph Theory*, To appear in 2013.

- [FLM<sup>+</sup>16] Fedor V. Fomin, Daniel Lokshtanov, Neeldhara Misra, Geevarghese Philip, and Saket Saurabh. Hitting forbidden minors: Approximation and kernelization. *SIAM Journal on Discrete Mathematics*, 30(1):383–410, 2016.
- [FLMS12] F. V. Fomin, D. Lokshtanov, N. Misra, and S. Saurabh. Planar  $\mathcal{F}$ -deletion: Approximation, kernelization and optimal fpt algorithms. In *Foundations of Computer Science (FOCS), 2012 IEEE 53rd Annual Symposium on*, pages 470–479, Oct 2012.
- [FLS12] Fedor V. Fomin, Daniel Lokshtanov, and Saket Saurabh. Bidimensionality and geometric graphs. In *23st ACM–SIAM Symposium on Discrete Algorithms (SODA 2012)*. ACM-SIAM, San Francisco, California, 2012.
- [FS13] Alexandra Fradkin and Paul Seymour. Tournament pathwidth and topological containment. *Journal of Combinatorial Theory, Series B*, 103(3):374 – 384, 2013.
- [FST11] Fedor V. Fomin, Saket Saurabh, and Dimitrios M. Thilikos. Strengthening Erdős–Pósa property for minor-closed graph classes. *Journal of Graph Theory*, 66(3):235–240, 2011.
- [Gal64] Tibor Gallai. Maximum-minimum sätze und verallgemeinerte faktoren von graphen. *Acta Mathematica Hungarica*, 12, 03 1964.
- [GGRW03] Jim F. Geelen, A. M. H. Gerards, Niel Robertson, and G. P. Whittle. On the excluded minors for the matroids of branch-width  $k$ . *Journal of Combinatorial Theory, Series B*, 88:261–265, July 2003.
- [GJ79] Michael R Garey and S David Johnson. *Computers and intractability: a guide to the theory of NP-completeness*. WH Free. Co., San Fr, 1979.
- [GK09] Jim Geelen and Kasper Kabell. The Erdős–Pósa property for matroid circuits. *Journal of Combinatorial Theory, Series B*, 99(2):407–419, March 2009.
- [GKRT16] Archontia Giannopoulou, O-joung Kwon, Jean-Florent Raymond, and Dimitrios M. Thilikos. Packing and covering immersion models of planar subcubic graphs. In *Graph-Theoretic Concepts in Computer Science: 42nd International Workshop, WG 2016, Istanbul, Turkey, June 22–24, 2016, Revised Selected Papers*, volume 9941, page 74. Springer, 2016.
- [GKS15] Robert Ganian, Eun Jung Kim, and Stefan Szeider. Algorithmic applications of tree-cut width. In *Mathematical foundations of computer science 2015. Part II*, volume 9235 of *Lecture Notes in Comput. Sci.*, pages 348–360. Springer, Heidelberg, 2015.

- [GL69] András Gyárfás and Jenő Lehel. A Helly-type problem in trees. In *Combinatorial Theory and its applications*, pages 571–584. Colloquia Mathematica Societatis János Bolyai, 1969.
- [GOB06] Sylvain Gravier and Cécile Ouvrier-Buffet. Maths à modeler: research-situations for teaching mathematics. In *Challenging Mathematics in and beyond the Classroom*. ICMI STUDY 16, 2006.
- [Gol15] Petr A. Golovach, 2015. Personal communication.
- [GPR<sup>+</sup>16a] Archontia C. Giannopoulou, Michał Pilipczuk, Jean-Florent Raymond, Dimitrios M. Thilikos, and Marcin Wrochna. Cutwidth: obstructions and algorithmic aspects. *ArXiv e-prints*, June 2016.
- [GPR<sup>+</sup>16b] Archontia C. Giannopoulou, Michał Pilipczuk, Jean-Florent Raymond, Dimitrios M. Thilikos, and Marcin Wrochna. Linear kernels for edge deletion problems to immersion-closed graph classes. *ArXiv preprint arXiv:1609.07780*, 2016.
- [Grü38] Tibor Grünwald. Ein neuer beweis eines mengerschen satzes. *Journal of the London Mathematical Society*, 1(3):188–192, 1938.
- [GT10] Bertrand Guenin and Robin Thomas. Packing directed circuits exactly. *ArXiv e-prints*, December 2010.
- [Hal91] Rudolf Halin. Tree-partitions of infinite graphs. *Discrete Mathematics*, 97(1–3):203 – 217, 1991.
- [Hax99] P.E. Haxell. Packing and covering triangles in graphs. *Discrete Mathematics*, 195(1–3):251 – 254, 1999.
- [Hig52] Graham Higman. Ordering by divisibility in abstract algebras. *Proceedings of the London Mathematical Society*, s3-2(1):326–336, 1952.
- [HJ07] Illya V. Hicks and Nolan B. McMurray Jr. The branchwidth of graphs and their cycle matroids. *Journal of Combinatorial Theory, Series B*, 97(5):681–692, 2007.
- [HJW16] Tony Huynh, Felix Joos, and Paul Wollan. A unified Erdős–Pósa theorem for constrained cycles. *ArXiv preprints arXiv:1605.07082*, May 2016.
- [HK88] E. P. Haxell and Y. Kohayakawa. Packing and covering triangles in tripartite graphs. *Graphs and Combinatorics*, 14(1):1–10, 1988.
- [HKT11] Penny Haxell, Alexandr Kostochka, and Stéphan Thomassé. Packing and covering triangles in  $k$  4-free planar graphs. *Graphs and Combinatorics*, 28(5):653–662, 2011.
- [HL14] Chun Hung Liu. *Graph Structures and Well-Quasi-Ordering*. PhD thesis, Georgia Tech, 2014.

- [HM13] Frédéric Havet and Ana Karolinna Maia. On disjoint directed cycles with prescribed minimum lengths. Research Report RR-8286, INRIA, April 2013.
- [HNP06] Winfried Hochstättler, Robert Nickel, and Britta Peis. Two disjoint negative cycles in a signed graph. *Electronic Notes in Discrete Mathematics*, 25:107–111, 2006.
- [HS58] András Hajnal and János Surányi. Über die auflösung von graphen in vollständige teilgraphen. *Annales Universitatis Scientarium Budapestinensis de Rolando Eötvös Nominatae, Sectio Mathematica*, 1:113–121, 1958.
- [iKK12] Ken ichi Kawarabayashi and Yusuke Kobayashi. Linear min-max relation between the treewidth of H-minor-free graphs and its largest grid. In Christoph Dürr and Thomas Wilke, editors, *29th International Symposium on Theoretical Aspects of Computer Science (STACS 2012)*, volume 14 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 278–289, Dagstuhl, Germany, 2012. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik.
- [Joh87] David S Johnson. The np-completeness column: An ongoing guide. *Journal of Algorithms*, 8(2):285 – 303, 1987.
- [Joo14] Felix Joos. Parity linkage and the Erdős-Pósa property of odd cycles through prescribed vertices in highly connected graphs. *ArXiv e-prints*, November 2014.
- [JPS<sup>+</sup>11] Gwenaël Joret, Christophe Paul, Ignasi Sau, Saket Saurabh, and Stéphan Thomassé. Hitting and harvesting pumpkins. In Camil Demetrescu and MagnúsM. Halldórsson, editors, *Algorithms – ESA 2011*, volume 6942 of *Lecture Notes in Computer Science*, pages 394–407. Springer Berlin Heidelberg, 2011.
- [JPS<sup>+</sup>14] Gwenaël Joret, Christophe Paul, Ignasi Sau, Saket Saurabh, and Stéphan Thomassé. Hitting and harvesting pumpkins. *SIAM Journal on Discrete Mathematics*, 28(3):1363–1390, 2014.
- [JRST01] Thor Johnson, Neil Robertson, Paul D. Seymour, and Robin Thomas. Directed tree-width. *Journal of Combinatorial Theory, Series B*, 82(1):138 – 154, 2001.
- [JW13] Gwenaël Joret and David R. Wood. Complete graph minors and the graph minor structure theorem. *Journal of Combinatorial Theory, Series B*, 103(1):61–74, 2013.
- [Kai97] Tomás Kaiser. Transversals of d-intervals. *Discrete & Computational Geometry*, 18(2):195–203, 1997.

- [Kan96] Goos Kant. Drawing planar graphs using the canonical ordering. *Algorithmica*, 16(1):4–32, 1996.
- [KiKM11] Naonori Kakimura, Ken ichi Kawarabayashi, and Dániel Marx. Packing cycles through prescribed vertices. *Journal of Combinatorial Theory, Series B*, 101(5):378 – 381, 2011.
- [KK12] Ken-Ichi Kawarabayashi and Yusuke Kobayashi. Edge-disjoint odd cycles in 4-edge-connected graphs. In *29th International Symposium on Theoretical Aspects of Computer Science (STACS)*, pages 206–217, 2012.
- [KK13] Naonori Kakimura and Ken-Ichi Kawarabayashi. Half-integral packing of odd cycles through prescribed vertices. *Combinatorica*, 33(5):549–572, 2013.
- [KKK12] Naonori Kakimura, Ken-ichi Kawarabayashi, and Yusuke Kobayashi. Erdős-Pósa property and its algorithmic applications: Parity constraints, subset feedback set, and subset packing. In *Proceedings of the Twenty-third Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1726–1736. SIAM, 2012.
- [KKKK13] Ken-ichi Kawarabayashi, Daniel Kral, M Krčál, and Stephan Kreutzer. Packing directed cycles through a specified vertex set. In *Proceedings of the Twenty-Fourth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 365–377. Society for Industrial and Applied Mathematics, 2013.
- [KL11a] Nicholas Korpelainen and Vadim Lozin. Two forbidden induced subgraphs and well-quasi-ordering. *Discrete Mathematics*, 311(16):1813 – 1822, 2011.
- [KL11b] Nicholas Korpelainen and Vadim V. Lozin. Bipartite induced subgraphs and well-quasi-ordering. *Journal of Graph Theory*, 67(3):235–249, 2011.
- [KLL02a] T. Kloks, C.M. Lee, and Jiping Liu. New algorithms for  $k$ -face cover,  $k$ -feedback vertex set, and  $k$ -disjoint cycles on plane and planar graphs. In *Proceedings of Graph-Theoretic Concepts in Computer Science*, volume 2573 of *Lecture Notes in Computer Science*, pages 282–295. Springer Berlin Heidelberg, 2002.
- [KLL02b] Ton Kloks, C. M. Lee, and Jiping Liu. New algorithms for  $k$ -face cover,  $k$ -feedback vertex set, and  $k$ -disjoint cycles on plane and planar graphs. In *28th International Workshop on Graph Theoretic Concepts in Computer Science (WG 2002)*, volume 2573 of *LNCS*, pages 282–295. Springer, Berlin, 2002.
- [Klo94a] Tom Kloks. *Treewidth. Computations and Approximations*, volume 842. Springer, LNCS, 1994.

- [Klo94b] Ton Kloks. *Treewidth. Computations and Approximations*, volume 842 of LNCS. Springer, 1994.
- [KM15] O-jong Kwon and Dániel Marx. Erdős–pósa property of planar- $h$ -minor models with prescribed vertex sets. Manuscript, 2015.
- [KN07] Ken-Ichi Kawarabayashi and Atsuhiro Nakamoto. The erdős–pósa property for vertex- and edge-disjoint odd cycles in graphs on orientable surfaces. *Discrete Mathematics*, 307(6):764 – 768, 2007.
- [KNS<sup>+</sup>07] Michael Krivelevich, Zeev Nutov, Mohammad R. Salavatipour, Jacques V. Yuster, and Raphael Yuster. Approximation algorithms and hardness results for cycle packing problems. *ACM Transactions on Algorithms*, 3(4), November 2007.
- [KNY05] Michael Krivelevich, Zeev Nutov, and Raphael Yuster. Approximation algorithms for cycle packing problems. In *Proceedings of the Sixteenth Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 556–561, Philadelphia, PA, USA, 2005. SIAM.
- [KO03] Daniela Kühn and Deryk Osthus. Minors in graphs of large girth. *Random Structures & Algorithms*, 22(2):213–225, 2003.
- [KO04a] Daniela Kühn and Deryk Osthus. Complete minors in  $K_{s,s}$ -free graphs. *Combinatorica*, 25(1):49–64, 2004.
- [KO04b] Daniela Kühn and Deryk Osthus. Induced subdivisions in  $k_{s,s}$ -free graphs of large average degree. *Combinatorica*, 24(2):287–304, 2004.
- [Kön31] Denés Kőnig. *Gráfok és mátrixok*, 1931.
- [Kos84] Alexandr V. Kostochka. Lower bound of the hadwiger number of graphs by their average degree. *Combinatorica*, 4(4):307–316, 1984.
- [KR09] Ken-Ichi Kawarabayashi and Bruce Reed. Highly parity linked graphs. *Combinatorica*, 29(2):215–225, 2009.
- [Kri95] Michael Krivelevich. On a conjecture of tuza about packing and covering of triangles. *Discrete Mathematics*, 142(1–3):281 – 286, 1995.
- [KRT14] M. Kamiński, J.-F. Raymond, and T. Trunck. Multigraphs without large bonds are well-quasi-ordered by contraction. *ArXiv e-prints*, December 2014.
- [KRT16] Marcin Kamiński, Jean-Florent Raymond, and Théophile Trunck. Well-quasi-ordering  $H$ -contraction-free graphs. *ArXiv e-prints*, February 2016.
- [Kru60] J. B. Kruskal. Well-quasi-ordering, the tree theorem, and Vazsonyi’s conjecture. *Transactions of the American Mathematical Society*, 95:210–225, 1960.

- [Kru72] Joseph B Kruskal. The theory of well-quasi-ordering: A frequently discovered concept. *Journal of Combinatorial Theory, Series A*, 13(3):297 – 305, 1972.
- [Kur30] Casimir Kuratowski. Sur le probleme des courbes gauches en topologie. *Fundamenta mathematicae*, 15(1):271–283, 1930.
- [KV04] Daniel Král’ and Heinz-Jürgen Voss. Edge-disjoint odd cycles in planar graphs. *Journal of Combinatorial Theory, Series B*, 90(1):107 – 120, 2004.
- [KW05] Ken-Ichi Kawarabayashi and Paul Wollan. Non-zero disjoint cycles in highly connected group labeled graphs. *Electronic Notes in Discrete Mathematics*, 22:271 – 275, 2005. 7th International Colloquium on Graph Theory.
- [Liu15] Chun-Hung Liu. Packing and Covering Immersions in 4-Edge-Connected Graphs. *ArXiv e-prints*, May 2015.
- [Lov06] László Lovász. Graph minor theory. *Bulletin of the American Mathematical Society*, 43(1):75–86, 2006.
- [LPW14] Chun-Hung Liu, Luke Postle, and Paul Wollan. Erdős-pósa property for the topological minors. Manuscript, 2014.
- [LS15] Alexander Leaf and Paul Seymour. Tree-width and planar minors. *Journal of Combinatorial Theory, Series B*, 111:38 – 53, 2015.
- [LY78] Claudio L. Lucchesi and D. H. Younger. A minimax theorem for directed graphs. *Journal of the London Mathematical Society (2)*, 17(3):369–374, 1978.
- [Mad78a] Wolfgang Mader. Über die maximalzahl kantendisjunkter  $a$ -wege. *Archiv der Mathematik*, 30(1):325–336, 1978.
- [Mad78b] Wolfgang Mader. Über die maximalzahl kreuzungsfreier  $h$ -wege. *Archiv der Mathematik*, 31(1):387–402, 1978.
- [Mar04] Klas Markström. Complete minors in cubic graphs with few short cycles and random cubic graphs. *Ars Combinatorica*, 70, 2004.
- [Men27] Karl Menger. Zur allgemeinen kurventheorie. *Fundamenta Mathematicae*, 1(10):96–115, 1927.
- [MNŠW16] Franck Mousset, Andreas Noever, Nemanja Škorić, and Felix Weissenberger. A tight Erdős-Pósa function for long cycles. *ArXiv e-prints*, March 2016.
- [MNT88] Jiří Matoušek, Jaroslav Nešetřil, and Robin Thomas. On polynomial time decidability of induced-minor-closed classes. *Commentationes Mathematicae Universitatis Carolinae*, 29(4):703–710, 1988.

- [Mod] Maths à Modeler. La chasse à la bête. <http://mathsamodeler.ujf-grenoble.fr>.
- [Mor94] M. Morgenstern. Existence and explicit constructions of  $q + 1$  regular ramanujan graphs for every prime power  $q$ . *Journal of Combinatorial Theory, Series B*, 62(1):44 – 62, 1994.
- [Mos06] Yiannis Moschovakis. *Notes on set theory*. Springer Science & Business Media, 2006.
- [MP67] John McCarthy and James Painter. Correctness of a compiler for arithmetic expressions. *Mathematical aspects of computer science*, 1, 1967.
- [MT01] Bojan Mohar and Carsten Thomassen. *Graphs on surfaces*. Johns Hopkins University Press, Baltimore, MD, 2001.
- [MW15] Dániel Marx and Paul Wollan. An exact characterization of tractable demand patterns for maximum disjoint path problems. In *Proceedings of the Twenty-Sixth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 642–661. SIAM, 2015.
- [MYZ13] Jie Ma, Xingxing Yu, and Wenan Zang. Approximate min-max relations on plane graphs. *Journal of Combinatorial Optimization*, 26(1):127–134, 2013.
- [NW63] C. St. J. A. Nash-Williams. On well-quasi-ordering finite trees. *Proceedings of the Cambridge Philosophical Society*, 59:833–835, 1963.
- [OOT93] Bogdan Oporowski, James Oxley, and Robin Thomas. Typical subgraphs of 3- and 4-connected graphs. *Journal of Combinatorial Theory, Series B*, 57(2):239–257, March 1993.
- [Oum08] Sang-il Oum. Rank-width and well-quasi-ordering. *SIAM Journal on Discrete Mathematics*, 22(2):666–682, 2008.
- [Oxl92] James G. Oxley. *Matroid Theory*. Oxford University Press, New York, 1992.
- [Pet02] Marko Petkovšek. Letter graphs and well-quasi-order by induced subgraphs. *Discrete Mathematics*, 244(1–3):375 – 388, 2002. Algebraic and Topological Methods in Graph Theory.
- [Pit14] Leonidas S Pitsoulis. *Topics in matroid theory*. Springer, 2014.
- [Pon91] I.N. Ponomarenko. The isomorphism problem for classes of graphs closed under contraction. *Journal of Soviet Mathematics*, 55(2):1621–1643, 1991.

- [Pou72] Maurice Pouzet. Un bel ordre d’abritement et ses rapports avec les bornes d’une multirelation. *CR Acad. Sci. Paris Sér. AB*, 274:A1677–A1680, 1972.
- [Pou85] Maurice Pouzet. Applications of well quasi-ordering and better quasi-ordering. In Ivan Rival, editor, *Graphs and Order: The Role of Graphs in the Theory of Ordered Sets and Its Applications*, pages 503–519. Springer Netherlands, Dordrecht, 1985.
- [Pro89] A. Proskurowski. *Maximal Graphs of Path-width  $K$  Or Searching a Partial  $K$ -caterpillar*. University of Oregon. Dept. of Computer and Information Science, 1989.
- [PW12] Matteo Pontecorvi and Paul Wollan. Disjoint cycles intersecting a set of vertices. *Journal of Combinatorial Theory, Series B*, 102(5):1134 – 1141, 2012.
- [Ray16] Jean-Florent Raymond. Hitting minors, subdivisions, and immersions in tournaments. *ArXiv preprint arXiv:1605.08366*, May 2016.
- [Ree97] Bruce A. Reed. *Tree width and tangles: A new connectivity measure and some applications*, pages 87–162. Cambridge University Press, 1997.
- [Ree99] Bruce Reed. Mangoes and blueberries. *Combinatorica*, 19(2):267–296, 1999.
- [RR01] Dieter Rautenbach and Bruce Reed. The Erdős–Pósa property for odd cycles in highly connected graphs. *Combinatorica*, 21(2):267–278, 2001.
- [RRST96a] Bruce Reed, Neil Robertson, Paul Seymour, and Robin Thomas. Packing directed circuits. *Combinatorica*, 16(4):535–554, 1996.
- [RRST96b] Bruce A. Reed, Neil Robertson, Paul D. Seymour, and Robin Thomas. Packing directed circuits. *Combinatorica*, 16(4):535–554, 1996.
- [RS86] Neil Robertson and Paul D. Seymour. Graph Minors. V. Excluding a planar graph. *Journal of Combinatorial Theory, Series B*, 41(2):92–114, 1986.
- [RS91] Neil Robertson and Paul D. Seymour. Graph Minors. X. Obstructions to Tree-decomposition. *Journal of Combinatorial Theory, Series B*, 52(2):153–190, 1991.
- [RS93] Neil Roberston and Paul D. Seymour. *Graph Structure Theory: Proceedings of the Joint Summer Research Conference on Graph Minors, Held June 22 to July 5, 1991, at the University of Washington, Seattle, with Support from the National Science Foundation and the Office of Naval Research*, volume 147. AMS Bookstore, 1993.
- [RS95] Neil Robertson and Paul D Seymour. Graph Minors. XIII. The disjoint paths problem. *Journal of combinatorial theory, Series B*, 63(1):65–110, 1995.

- [RS96] Bruce A. Reed and Bruce F. Shepherd. The Gallai-Younger conjecture for planar graphs. *Combinatorica*, 16(4):555–566, 1996.
- [RS03] Neil Robertson and Paul D Seymour. Graph Minors. XVI. Excluding a non-planar graph. *Journal of Combinatorial Theory, Series B*, 89(1):43–76, 2003.
- [RS04] Neil Robertson and Paul D. Seymour. Graph Minors. XX. Wagner’s conjecture. *Journal of Combinatorial Theory, Series B*, 92(2):325 – 357, 2004.
- [RS10] Neil Robertson and Paul Seymour. Graph Minors XXIII. Nash-Williams’ immersion conjecture. *Journal of Combinatorial Theory, Series B*, 100(2):181–205, March 2010.
- [RST94a] Neil Robertson, Paul D. Seymour, and Robin Thomas. Quickly excluding a planar graph. *J. Combin. Theory Ser. B*, 62(2):323–348, 1994.
- [RST94b] Neil Robertson, Paul D. Seymour, and Robin Thomas. Quickly excluding a planar graph. *Journal of Combinatorial Theory, Series B*, 62(2):323–348, 1994.
- [RST16] Jean-Florent Raymond, Ignasi Sau, and Dimitrios M. Thilikos. An edge variant of the Erdős-Pósa property. *Discrete Mathematics*, 339(8):2027 – 2035, 2016.
- [RT13a] J.-F. Raymond and D. M. Thilikos. Polynomial Gap Extensions of the Erdős-Pósa Theorem. *ArXiv e-prints*, May 2013.
- [RT13b] Jean-Florent Raymond and Dimitrios M. Thilikos. Polynomial gap extensions of the Erdős-Pósa theorem. In Jaroslav Nešetřil and Marco Pellegrini, editors, *The Seventh European Conference on Combinatorics, Graph Theory and Applications: EuroComb 2013*, pages 13–18, Pisa, 2013. Scuola Normale Superiore.
- [RT15] Jean-Florent Raymond and Dimitrios M. Thilikos. Low polynomial exclusion of planar graph patterns. *Journal of Graph Theory*, pages n/a–n/a, 2015.
- [RT16] Jean-Florent Raymond and Dimitrios M. Thilikos. Recent techniques and results on the Erdős-Pósa property. *ArXiv e-prints*, March 2016.
- [Sch01] Alexander Schrijver. A short proof of mader’s s-paths theorem. *Journal of Combinatorial Theory, Series B*, 82(2):319–321, 2001.
- [See85] Detlef Seese. Tree-partite graphs and the complexity of algorithms. In *Proceedings of Fundamentals of Computation Theory*, volume 199 of *Lecture Notes in Computer Science*, pages 412–421. Springer Berlin Heidelberg, 1985.

- [Sey96] Paul D. Seymour. Packing circuits in eulerian digraphs. *Combinatorica*, 16(2):223–231, 1996.
- [Sim67] Miklós Simonovits. A new proof and generalizations of a theorem of erdős and pósa on graphs without  $k + 1$  independent circuits. *Acta Mathematica Academiae Scientiarum Hungarica*, 18(1):191–206, 1967.
- [SS04] András Sebő and László Szegő. The path-packing structure of graphs. In Daniel Bienstock and George Nemhauser, editors, *Integer Programming and Combinatorial Optimization: 10th International IPCO Conference, New York, NY, USA, June 7-11, 2004. Proceedings*, pages 256–270, Berlin, Heidelberg, 2004. Springer Berlin Heidelberg.
- [Sti96] Michael Stiebitz. Decomposing graphs under degree constraints. *Journal of Graph Theory*, 23(3):321–324, 1996.
- [SV05] Mohammad R. Salavatipour and Jacques Verstraete. Disjoint cycles: Integrality gap, hardness, and approximation. In *Proceedings of Integer Programming and Combinatorial Optimization*, volume 3509 of *Lecture Notes in Computer Science*, pages 51–65. Springer Berlin Heidelberg, 2005.
- [Tho83a] A. Thomason. An extremal function for contractions of graphs. *Mathematical Proceedings of the Cambridge Philosophical Society*, 95:261, 1983.
- [Tho83b] Carsten Thomassen. Girth in graphs. *Journal of Combinatorial Theory, Series B*, 35(2):129–141, 1983.
- [Tho85] Robin Thomas. Graphs without  $K_4$  and well-quasi-ordering. *Journal of Combinatorial Theory, Series B*, 38(3):240 – 247, 1985.
- [Tho88] Carsten Thomassen. On the presence of disjoint subgraphs of a specified type. *Journal of Graph Theory*, 12(1):101–111, 1988.
- [Tho01a] Andrew Thomason. The extremal function for complete minors. *Journal of Combinatorial Theory, Series B*, 81(2):318 – 338, 2001.
- [Tho01b] Carsten Thomassen. The Erdős-Pósa property for odd cycles in graphs of large connectivity. *Combinatorica*, 21(2):321–333, 2001.
- [Tur41] Paul Turán. On an extremal problem in graph theory. *Mat. Fiz. Lapok*, 48(436-452):137, 1941.
- [Tut61] W. T. Tutte. A theory of 3-connected graphs. *Indagationes Mathematicae*, 23(1961):441–455, 1961.
- [Tuz90] Zsolt Tuza. A conjecture on triangles of graphs. *Graphs and Combinatorics*, 6(4):373–380, 1990.
- [Vos68] Heinz-Jürgen Voss. Some properties of graphs containing  $k$  independent circuits. In *Proceedings of Colloquium Tihany*, pages 321–334, 1968.

- [Wol11] Paul Wollan. Packing cycles with modularity constraints. *Combinatorica*, 31(1):95–126, 2011.
- [Wol15] Paul Wollan. The structure of graphs not admitting a fixed immersion. *Journal of Combinatorial Theory, Series B*, 110:47–66, 2015.
- [YBdFT99] K. Yamazaki, H. L. Bodlaender, B. de Fluiter, and D. M. Thilikos. Isomorphism for graphs of bounded distance width. *Algorithmica*, 24(2):105–127, 1999.



---

## LIST OF FIGURES

1	Quelques fruits ordonnés par désirabilité. . . . .	11
2	Subdivisions de $K_5$ . . . . .	12
3	Les graphes de Kuratowski $K_5$ et $K_{3,3}$ (de gauche à droite). . . . .	13
4	Deux situations où le choix est difficile. . . . .	14
5	Les cycles forment un ensemble infini de graphes deux à deux incomparables pour l'ordre de sous-graphe. . . . .	16
6	Rats et pièges. . . . .	16
7	La chasse au rat. . . . .	17
8	Une solution pour rats n'est pas toujours une solution pour serpents. . . . .	18
9	Un mur d'Escher de hauteur 4. . . . .	19
1.1	Some fruits ordered by desirability. . . . .	23
1.2	Subdivisions of $K_5$ . . . . .	24
1.3	Kuratowski's graphs $K_5$ and $K_{3,3}$ (from left to right). . . . .	25
1.4	Two situation where the choice is hard. . . . .	26
1.5	Cycles form an infinite set of graphs that are pairwise not comparable for the subgraph ordering. . . . .	28
1.6	Rats and traps. . . . .	28
1.7	Hunting rats. . . . .	29
1.8	A solution for rats is not always a solution for snakes. . . . .	29
1.9	An Escher wall of height 4. . . . .	30
2.1	The graphs $\theta_5$ (left) and $D_4$ (right). . . . .	36
2.2	The $(4 \times 4)$ -grid (left) and the 4-wall (right). . . . .	37
2.3	A wheel of order six (left) and a double wheel of order 6 (right). . . . .	37
2.4	The yurt graph of order 5. . . . .	38
2.5	The contraction of the edge $e$ creates a double edge. . . . .	38
2.6	Dissolution and subdivision as complementary operations. . . . .	39
2.7	A $K_4$ -minor model in the $3 \times 3$ grid. . . . .	40
2.8	A $K_4$ -immersion model in a graph that has no $K_4$ -minor model. . . . .	41
2.9	Connections between common orderings of graphs. . . . .	44

2.10	An example of a tree-partition of a graph. . . . .	46
2.11	An example of a tree-cut decomposition of a graph. . . . .	47
3.1	A 3-colorable graph that contains $K_4$ as minor. . . . .	50
3.2	The graphs $\hat{K}_4$ , gem, and $D_2$ . . . . .	53
3.3	The elements of a canonical antichain for the subgraph relation. . . . .	55
3.4	Two infinite antichains for multigraph contractions: multiedges and edgeless graphs. . . . .	55
3.5	Thomas' alternating double wheels. . . . .	63
3.6	The infinite antichain $\mathcal{A}_M$ of Matoušek, Nešetřil, and Thomas. . . . .	64
3.7	A planar drawing of the third graph of $\mathcal{A}_M$ . . . . .	65
3.8	An illustration of the intervals in $T_6$ . . . . .	65
3.9	Antiholes antichain. . . . .	66
3.10	Nested lozenges. . . . .	66
3.11	The antichain $\mathcal{A}_K$ . . . . .	67
3.12	The multiedges antichain. . . . .	67
3.13	Forbidden configurations in the proof of Lemma 3.53. . . . .	74
3.14	Graphs used in Lemma 3.55 and in Lemma 3.56. . . . .	76
3.15	Two different cases in the proof of Lemma 3.57 . . . . .	77
3.16	The prism. . . . .	77
3.17	The graph $H$ of Lemma 3.59 and the graph obtained after deletion of $y$ . . . . .	78
3.18	The rooted diamond. . . . .	87
3.19	Induced minor-minimal counterexamples in Lemma 3.78. . . . .	90
3.20	Models of Gem in graphs from Figure 3.19. . . . .	91
3.21	A clique-cactus graph. . . . .	96
3.22	Graphs of $\mathcal{D}$ . . . . .	96
3.23	A bond of size 3 in the house graph. . . . .	105
3.24	Attaching $H$ to vertices $(u, v)$ of $G$ . . . . .	105
3.25	Two infinite antichains for multigraph contractions: multiedges and edgeless graphs. . . . .	113
4.1	An example of a distance-decomposition of a graph. . . . .	118
4.2	An unimportant path in a tree. . . . .	122
4.3	A wheel of order six and a double wheel of order 6. . . . .	138
4.4	The graph $\Xi_5$ . . . . .	143
4.5	Example for Lemma 4.54. . . . .	146
4.6	The yurt graph of order 5. . . . .	148
4.7	The graph $\widehat{W}_5$ . . . . .	153
4.8	Finding $\Gamma_5$ as a strong immersion in $\widehat{W}_5$ . . . . .	153
4.9	Swapping branch vertices. . . . .	155
5.1	Unavoidable patterns of graphs of large tree-partition width. . . . .	171
5.2	The gadget $\Gamma_{4,3}$ used in Lemma 5.42. . . . .	178
5.3	The "thickened" $K_5$ for $k = 3$ . . . . .	179

5.4 A 2-connected graph  $H$  for which  $\mathcal{I}(H)$  does not have the (vertex or edge) Erdős–Pósa property. . . . . 182

5.5 The host graph  $G$ . . . . . 182

6.1 Gluing graphs together:  $G = \mathbf{G}_1 \oplus \mathbf{G}_2$ . . . . . 197

6.2 Cutting a graph. . . . . 198

6.3 An  $e\text{-}\mathcal{T}(H)$ -packing in  $\mathbf{G}$  . . . . . 200

6.4 A partial subdivision from the packing of Figure 6.3. . . . . 200

6.5 The compression of the partial packing of Figure 6.4. . . . . 201



---

# LIST OF TABLES

2.1	Requirements for containment models . . . . .	40
2.2	Containment relations defined in terms of local operations. . . . .	43
3.1	Status of Question 3.2 for the common orderings of graphs. . . . .	51
3.2	If $(\text{Excl}_{\leq_{i.m.}}(H), \leq_{i.m.})$ is a wqo, then $\overline{H}$ belongs to this table. . . . .	71
3.3	If $(\text{Excl}_{\leq_{i.m.}}(H), \leq_{i.m.})$ is a wqo, then $H$ belongs to this table. . . . .	71
7.1	Summary of the exclusion results of Chapter 4. . . . .	212



---

# INDEX

- $A^*$ , 33
- $A_x$ , 159
- $C_n$ , 36
- $D_n$ , 36
- $E(\cdot)$ , 34
- $E(\cdot, \cdot)$ , 35
- $G \setminus X$ , 34
- $G[X]$ , 34
- $H \times G$ , 34
- $K_n$ , 36
- $N_G(\cdot)$ , 35
- $P_n$ , 36
- $V(\cdot)$ , 34
- $W_k$ , 37
- $\Delta(\cdot)$ , 35
- $\Gamma_{k,r}$ , 37
- $\mathcal{I}(\cdot)$ , 43
- $\mathcal{T}(\cdot)$ , 43
- $\deg_G(\cdot)$ , 35
- $\delta(\cdot)$ , 35
- desc, 36
- diam( $\cdot$ ), 35
- dist( $\cdot, \cdot$ ), 35
- Excl, 57
- Incl, 57
- lab, 57
- lca $_T(\cdot, \cdot)$ , 36
- $\preceq^*$ , 56
- $\preceq^P$ , 56
- $\preceq_A \cup \preceq_B$ , 56
- $\preceq_A \times \preceq_B$ , 56
- $\mathcal{M}(\cdot)$ , 43
- mdeg $_G(\cdot)$ , 35
- mult, 34
- mult $_G(\cdot, \cdot)$ , 35
- $\mathring{u}Tv$ , 35
- $uT\mathring{v}$ , 35
- $uTv$ , 35
- pw( $\cdot$ ), 45
- e, 159
- $\langle \cdot \rangle$ , 33
- v, 159
- x, 159
- tcw( $\cdot$ ), 47
- $\theta_n$ , 36
- tpw( $\cdot$ ), 46
- tw( $\cdot$ ), 44
- x-cover, 159
- x-pack, 159
- $m(\cdot)$ , 34
- $n(\cdot)$ , 34
- ecc $_G(\cdot)$ , 35
- p, 36
- $|\cdot|$ , 34
- antichain, 34
  - canonical antichain, 57
  - fundamental antichain, 57
- bag, 43, 46
  - torso of a bag, 44

basic induced subgraph, 68  
 block, 35  
  
 cartesian product of graphs, 34  
 child, 36  
 closure, 39, 57  
 cograph, 36  
 connected  
      $k$ -connected graph, 35  
     2-connected component, 35  
     component, 35  
     graph, 35  
 contraction, 41  
 cover, 159  
     covering number, 159  
 cut, 35  
     cut-set, 35  
     width of a cut, 35  
  
 degree, 35  
 descendant, 36  
 diamond, 36, 53  
 distance, 35  
  
 eccentricity, 35  
 edge, 34  
     edge contraction, 38  
     edge subdivision, 39  
     incidence, 35  
 epi, *see* epimorphism  
 epimorphism, 56  
 Erdős–Pósa property, 159  
  
 $\mathcal{C}$ -free, 34  
  
 gap, 159  
 gem, 53  
 girth, 35  
 good pair, 34  
 graph, 34  
     labeled graph, 57  
     rooted graph, 58  
     simple graph, 34  
     subcubic, 36  
     underlying simple graph, 34  
 grid, 37  
  
 height, 36  
  
 immersion, 42  
     expansion, 43  
     strong immersion, 42  
  
 join, 84  
  
 leaf, 35  
 least common ancestor, 36  
 lift, 39  
 linear forest, 36  
  
 major, 43  
 minor, 41  
     induced minor, 41  
     topological minor, 42  
 model, 40  
 monotone function, 56  
  
 near-partition, 46  
 neighborhood, 35  
 non-comparable, 34  
  
 order  
     quasi, 33  
     well-quasi-, 34  
  
 packing, 159  
     packing number, 159  
 parent, 36  
 path decomposition, 45  
     nice path decomposition, 45  
 pathwidth, 45  
 prism, 36  
  
 root, 36  
  
 separation, 35  
 sequence, 33  
     bad sequence, 34  
 subdivision, 43  
 subgraph, 41  
      $\mathcal{C}$ -subgraph, 34  
     induced subgraph, 34, 41  
  
 tree, 35  
     rooted tree, 36

- tree partition, 46
  - rooted tree partition, 46
  - width of a tree partition, 46
- tree partition width, 46
- tree-cut decomposition, 46
  - nice tree-cut decomposition, 48
  - rooted tree-cut decomposition, 47
- tree-cut width, 47
- tree-decomposition, 43
  - nice tree-decomposition, 45
  - width of a tree-decomposition, 44
- treewidth, 44
  
- uniquely defined, 57
  
- vertex, 34
  - adjacence, 35
  - deletion, 38
  - vertex dissolution, 39
  - vertex subdivision, 39
  
- wall, 37
  - horizontal paths, 37
  - vertical paths, 37
- wheel, 37
  - cycle and center of a wheel, 37
  - double wheel, 37
  
- yurt, 38

---

## Résumé

Le thème central à cette thèse est l'étude des propriétés des classes de graphes définies par sous-structures interdites et leurs applications.

La première direction que nous suivons a trait aux beaux ordres. À l'aide de théorèmes de décomposition dans les classes de graphes interdisant une sous-structure, nous identifions celles qui sont bellement-ordonnées. Les ordres et sous-structures considérés sont ceux associés aux notions de contraction et mineur induit.

Ensuite, toujours en considérant des classes de graphes définies par sous-structures interdites, nous obtenons des bornes sur des invariants comme le degré, la largeur arborescente, la tree-cut width et un nouvel invariant généralisant la maille.

La troisième direction est l'étude des relations entre les invariants combinatoires liés aux problèmes de packing et de couverture de graphes. Dans cette direction, nous établissons de nouvelles relations entre ces invariants pour certaines classes de graphes. Nous présentons également des applications algorithmiques de ces résultats.

**Mots clefs :** théorie des graphes structurelle, sous-structures interdites, beaux ordres, propriété d'Erdős-Pósa.

---

## Abstract

The central theme of this thesis is the study of the properties of the classes of graphs defined by forbidden substructures and their applications.

The first direction that we follow concerns well-quasi-orders. Using decomposition theorems on graph classes forbidding one substructure, we identify those that are well-quasi-ordered. The orders and substructures that we consider are those related to the notions of contraction and induced minor.

Then, still considering classes of graphs defined by forbidden substructures, we obtain bounds on invariants such as degree, treewidth, tree-cut width, and a new invariant generalizing the girth.

The third direction is the study of the links between the combinatorial invariants related to problems of packing and covering of graphs. In this direction, we establish new connections between these invariants for some classes of graphs. We also present algorithmic applications of the results.

**Keywords:** structural Graph Theory, forbidden substructures, well-quasi-orders, Erdős-Pósa property.

---